

B70

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 : A61K 39/00, G01N 33/53, C12P 15/09		A2	(11) International Publication Number: WO 00/57903
			(43) International Publication Date: 5 October 2000 (05.10.00)
(21) International Application Number: PCT/US00/07525		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 22 March 2000 (22.03.00)		Published With declaration under Article 17(2)(a); without abstract; title not checked by the International Searching Authority.	
(30) Priority Data: 60/126,595 26 March 1999 (26.03.99) US 60/171,549 22 December 1999 (22.12.99) US			
(71) Applicant (for all designated States except US): HUMAN GENOME SCIENCES, INC. [US/US]; 9410 Key West Avenue, Rockville, MD 20850 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): ROSEN, Craig, A. [US/US]; 22400 Rolling Hill Road, Laytonsville, MD 20882 (US). RUBEN, Steven, M. [US/US]; 18528 Heritage Hills Drive, Laytonsville, MD 20882 (US). KOMAT-SOULIS, George [US/US]; 9518 Garwood Street, Silver Spring, MD 20901 (US).			
(74) Agents: HOOVER, Kenley, K. et al.; Human Genome Sciences, Inc., 9410 Key West Avenue, Rockville, MD 20850 (US).			
(54) Title: 48 HUMAN SECRETED PROTEINS			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

48 Human Secreted Proteins

Field of the Invention

This invention relates to newly identified polynucleotides and the polypeptides encoded by these polynucleotides, uses of such polynucleotides and polypeptides, and their production.

Background of the Invention

Unlike bacterium, which exist as a single compartment surrounded by a membrane, human cells and other eucaryotes are subdivided by membranes into many functionally distinct compartments. Each membrane-bounded compartment, or organelle, contains different proteins essential for the function of the organelle. The cell uses "sorting signals," which are amino acid motifs located within the protein, to target proteins to particular cellular organelles.

One type of sorting signal, called a signal sequence, a signal peptide, or a leader sequence, directs a class of proteins to an organelle called the endoplasmic reticulum (ER). The ER separates the membrane-bounded proteins from all other types of proteins. Once localized to the ER, both groups of proteins can be further directed to another organelle called the Golgi apparatus. Here, the Golgi distributes the proteins to vesicles, including secretory vesicles, the cell membrane, lysosomes, and the other organelles.

Proteins targeted to the ER by a signal sequence can be released into the extracellular space as a secreted protein. For example, vesicles containing secreted proteins can fuse with the cell membrane and release their contents into the extracellular space - a process called exocytosis. Exocytosis can occur constitutively or after receipt of a triggering signal. In the latter case, the proteins are stored in secretory vesicles (or secretory granules) until exocytosis is triggered. Similarly, proteins residing on the cell membrane can also be secreted into the extracellular space by proteolytic cleavage of a "linker" holding the protein to the membrane.

Despite the great progress made in recent years, only a small number of genes encoding human secreted proteins have been identified. These secreted proteins include the commercially valuable human insulin, interferon, Factor VIII, human

growth hormone, tissue plasminogen activator, and erythropoietin. Thus, in light of the pervasive role of secreted proteins in human physiology, a need exists for identifying and characterizing novel human secreted proteins and the genes that encode them. This knowledge will allow one to detect, to treat, and to prevent
5 medical diseases, disorders, and/or conditions by using secreted proteins or the genes that encode them.

Summary of the Invention

The present invention relates to novel polynucleotides and the encoded
10 polypeptides. Moreover, the present invention relates to vectors, host cells, antibodies, and recombinant and synthetic methods for producing the polypeptides and polynucleotides. Also provided are diagnostic methods for detecting diseases, disorders, and/or conditions related to the polypeptides and polynucleotides, and therapeutic methods for treating such diseases, disorders, and/or conditions. The
15 invention further relates to screening methods for identifying binding partners of the polypeptides.

Detailed Description

Definitions

20 The following definitions are provided to facilitate understanding of certain terms used throughout this specification.

In the present invention, "isolated" refers to material removed from its original environment (e.g., the natural environment if it is naturally occurring), and thus is altered "by the hand of man" from its natural state. For example, an isolated
25 polynucleotide could be part of a vector or a composition of matter, or could be contained within a cell, and still be "isolated" because that vector, composition of matter, or particular cell is not the original environment of the polynucleotide. The term "isolated" does not refer to genomic or cDNA libraries, whole cell total or mRNA preparations, genomic DNA preparations (including those separated by
30 electrophoresis and transferred onto blots), sheared whole cell genomic DNA

preparations or other compositions where the art demonstrates no distinguishing features of the polynucleotide/sequences of the present invention.

In the present invention, a "secreted" protein refers to those proteins capable
5 of being directed to the ER, secretory vesicles, or the extracellular space as a result of a signal sequence, as well as those proteins released into the extracellular space without necessarily containing a signal sequence. If the secreted protein is released into the extracellular space, the secreted protein can undergo extracellular processing to produce a "mature" protein. Release into the extracellular space can occur by many
10 mechanisms, including exocytosis and proteolytic cleavage.

In specific embodiments, the polynucleotides of the invention are at least 15, at least 30, at least 50, at least 100, at least 125, at least 500, or at least 1000 continuous nucleotides but are less than or equal to 300 kb, 200 kb, 100 kb, 50 kb, 15 kb, 10 kb, 7.5 kb, 5 kb, 2.5 kb, 2.0 kb, or 1 kb, in length. In a further embodiment,
15 polynucleotides of the invention comprise a portion of the coding sequences, as disclosed herein, but do not comprise all or a portion of any intron. In another embodiment, the polynucleotides comprising coding sequences do not contain coding sequences of a genomic flanking gene (i.e., 5' or 3' to the gene of interest in the genome). In other embodiments, the polynucleotides of the invention do not contain
20 the coding sequence of more than 1000, 500, 250, 100, 50, 25, 20, 15, 10, 5, 4, 3, 2, or 1 genomic flanking gene(s).

As used herein, a "polynucleotide" refers to a molecule having a nucleic acid sequence contained in SEQ ID NO:X or the cDNA contained within the clone deposited with the ATCC. For example, the polynucleotide can contain the
25 nucleotide sequence of the full length cDNA sequence, including the 5' and 3' untranslated sequences, the coding region, with or without the signal sequence, the secreted protein coding region, as well as fragments, epitopes, domains, and variants of the nucleic acid sequence. Moreover, as used herein, a "polypeptide" refers to a molecule having the translated amino acid sequence generated from the
30 polynucleotide as broadly defined.

In the present invention, the full length sequence identified as SEQ ID NO:X was often generated by overlapping sequences contained in multiple clones (contig analysis). A representative clone containing all or most of the sequence for SEQ ID NO:X was deposited with the American Type Culture Collection ("ATCC"). As shown in Table 1, each clone is identified by a cDNA Clone ID (Identifier) and the ATCC Deposit Number. The ATCC is located at 10801 University Boulevard, Manassas, Virginia 20110-2209, USA. The ATCC deposit was made pursuant to the terms of the Budapest Treaty on the international recognition of the deposit of microorganisms for purposes of patent procedure.

A "polynucleotide" of the present invention also includes those polynucleotides capable of hybridizing, under stringent hybridization conditions, to sequences contained in SEQ ID NO:X, the complement thereof, or the cDNA within the clone deposited with the ATCC. "Stringent hybridization conditions" refers to an overnight incubation at 42 degree C in a solution comprising 50% formamide, 5x SSC (750 mM NaCl, 75 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65 degree C.

Also contemplated are nucleic acid molecules that hybridize to the polynucleotides of the present invention at lower stringency hybridization conditions. Changes in the stringency of hybridization and signal detection are primarily accomplished through the manipulation of formamide concentration (lower percentages of formamide result in lowered stringency); salt conditions, or temperature. For example, lower stringency conditions include an overnight incubation at 37 degree C in a solution comprising 6X SSPE (20X SSPE = 3M NaCl; 0.2M NaH_2PO_4 ; 0.02M EDTA, pH 7.4), 0.5% SDS, 30% formamide, 100 µg/ml salmon sperm blocking DNA; followed by washes at 50 degree C with 1XSSPE, 0.1% SDS. In addition, to achieve even lower stringency, washes performed following stringent hybridization can be done at higher salt concentrations (e.g. 5X SSC).

Note that variations in the above conditions may be accomplished through the inclusion and/or substitution of alternate blocking reagents used to suppress

background in hybridization experiments. Typical blocking reagents include Denhardt's reagent, BLOTTO, heparin, denatured salmon sperm DNA, and commercially available proprietary formulations. The inclusion of specific blocking reagents may require modification of the hybridization conditions described above,
5 due to problems with compatibility.

Of course, a polynucleotide which hybridizes only to polyA+ sequences (such as any 3' terminal polyA+ tract of a cDNA shown in the sequence listing), or to a complementary stretch of T (or U) residues, would not be included in the definition of "polynucleotide," since such a polynucleotide would hybridize to any nucleic acid
10 molecule containing a poly (A) stretch or the complement thereof (e.g., practically any double-stranded cDNA clone generated using oligo dT as a primer).

The polynucleotide of the present invention can be composed of any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. For example, polynucleotides can be composed of
15 single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, the polynucleotide can be composed of
20 triple-stranded regions comprising RNA or DNA or both RNA and DNA. A polynucleotide may also contain one or more modified bases or DNA or RNA backbones modified for stability or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications can be made to DNA and RNA; thus, "polynucleotide" embraces
25 chemically, enzymatically, or metabolically modified forms.

The polypeptide of the present invention can be composed of amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres, and may contain amino acids other than the 20 gene-encoded amino acids. The polypeptides may be modified by either natural processes, such as
30 posttranslational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts and in more

detailed monographs, as well as in a voluminous research literature. Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be branched, for example, as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched, and branched cyclic polypeptides may result from posttranslation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cysteine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, pegylation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination. (See, for instance, PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York (1993); POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS, B. C. Johnson, Ed., Academic Press, New York, pgs. 1-12 (1983); Seifter et al., Meth Enzymol 182:626-646 (1990); Rattan et al., Ann NY Acad Sci 663:48-62 (1992).)

"SEQ ID NO:X" refers to a polynucleotide sequence while "SEQ ID NO:Y" refers to a polypeptide sequence, both sequences identified by an integer specified in Table 1.

"A polypeptide having biological activity" refers to polypeptides exhibiting activity similar, but not necessarily identical to, an activity of a polypeptide of the present invention, including mature forms, as measured in a particular biological assay, with or without dose dependency. In the case where dose dependency does exist, it need not be identical to that of the polypeptide, but rather substantially similar

to the dose-dependence in a given activity as compared to the polypeptide of the present invention (i.e., the candidate polypeptide will exhibit greater activity or not more than about 25-fold less and, preferably, not more than about tenfold less activity, and most preferably, not more than about three-fold less activity relative to the polypeptide of the present invention.)

Many proteins (and translated DNA sequences) contain regions where the amino acid composition is highly biased toward a small subset of the available residues. For example, membrane spanning domains and signal peptides (which are also membrane spanning) typically contain long stretches where Leucine (L), Valine (V), Alanine (A), and Isoleucine (I) predominate. Poly-Adenosine tracts (polyA) at the end of cDNAs appear in forward translations as poly-Lysine (poly-K) and poly-Phenylalanine (poly-F) when the reverse complement is translated. These regions are often referred to as "low complexity" regions.

Such regions can cause database similarity search programs such as BLAST to find high-scoring sequence matches that do not imply true homology. The problem is exacerbated by the fact that most weight matrices (used to score the alignments generated by BLAST) give a match between any of a group of hydrophobic amino acids (L, V and I) that are commonly found in certain low complexity regions almost as high a score as for exact matches.

In order to compensate for this, BLASTX.2 (version 2.0a5MP-WashU) employs two filters ("seg" and "xnu") which "mask" the low complexity regions in a particular sequence. These filters parse the sequence for such regions, and create a new sequence in which the amino acids in the low complexity region have been replaced with the character "X". This is then used as the input sequence (sometimes referred to herein as "Query" and/or "Q") to the BLASTX program. While this regime helps to ensure that high-scoring matches represent true homology, there is a negative consequence in that the BLASTX program uses the query sequence that has been masked by the filters to draw alignments.

Thus, a stretch of "X"s in an alignment shown in the following application does not necessarily indicate that either the underlying DNA sequence or the translated protein sequence is unknown or uncertain. Nor is the presence of such

stretches meant to indicate that the sequence is identical or not identical to the sequence disclosed in the alignment of the present invention. Such stretches may simply indicate that the BLASTX program masked amino acids in that region due to the detection of a low complexity region, as defined above. In all cases, the reference sequence(s) (sometimes referred to herein as "Subject", "Sbjct", and/or "S") indicated in the specification, sequence table (Table 1), and/or the deposited clone is (are) the definitive embodiment(s) of the present invention, and should not be construed as limiting the present invention to the partial sequence shown in an alignment, unless specifically noted otherwise herein.

Polynucleotides and Polypeptides of the Invention

FEATURES OF PROTEIN ENCODED BY GENE NO: 1

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gi1196433 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "unknown protein [Homo sapiens]." A partial alignment demonstrating the observed homology is shown immediately below.

```
>gi1196433 unknown protein [Homo sapiens] >sp|Q14288|Q14288 HYPOTHETICAL
PROTEIN (FRAGMENT).
Length = 641
```

Minus Strand HSPs:

```
Score = 743 (261.5 bits), Expect = 6.6e-115, Sum P(3) = 6.6e-115
Identities = 170/327 (51%), Positives = 218/327 (66%), Frame = -1
```

```
30 Q: 1236 SMYKNQ*HFYTPITSKLESQMKNITLFTIAIKKTKYLEIHLTKEVKDLYKEN*ETLLKEI 1057
    ++ K+Q YT + ESQ+ + + FTIA K+ KYL I LT++VKDL+KEN + LLKEI
S: 100 NVQKSQAFLYTN-NRQTESQIMSELPFTIASKRIKYLGIQLTRDVKDLFKENYKPLLKEI 158

35 Q: 1056 TDDTDKWKNI PCSWITRINIVKMAILLKAIYRFNVIPIK*PVSFFTELEKTILKFIWNQK 877
    +DT+KWKNI PCSW+ RINIVKMAIL K IYRFN IPIK P++FFTELEKT LKFIWNQK
S: 159 KEDTNKWKNI PCSWVGRINIVKMAILPKVIYRFNAIPIKLPM TFFTELEKTTLKFIWNQK 218

40 Q: 876 RTQITKAIPSKKNKAEASHY--LTLYYTIKLQ*PKQHGTTGKADKQTKGTELRTQK*SCT 703
    R +I K+I S+KNKA LYY + + + "Q" RT+
S: 219 RARIASILSQKNKAGGITLPDFKLYKATVTKTAWYQNRDIDQWN----RTEPSEIM 274

Q: 702 PTATCSLT--KSTKNRQWGKDSLFSKWC*DSWLAICKRMKQRFILW*KDPYLSPTYKINS 529
    P L K KN+QWGKDSLFSKWC ++WLAIC+++K DP+L+PTYKINS
S: 275 PHIYNYLIFDKPEKNQWGKDSLFSKWCWENWLAICRKLK-----DPFLTPYTKINS 327
```


Q: 528 RWIKYLSVRPQTIRILQENLGNTILDIGSGKEFMTKXXXXXXXXXXXXXXXXXELLHYK 349
 RWIK L+VRP+TI+ L+ENLG TI DIG GK+FM+K +L+ K
 S: 328 RWIKDLNVRPKTIKLEENLGITIEDIGVGKDFMSK----TPKAMATKAKIDKWDLIK 383
 Q: 348 RNFQWSKQTTYKI-----EKM FANYASDKGLTSRIY 256
 +F +K+TT ++ EK+FA Y+SDKGL SRIY
 S: 384 -SFCTAKETTIRVNRQPTTWEKIFATYSSDKGLISRIY 420

10 The segment of gill 196433 that is shown as "S" above is set out in the sequence listing as SEQ ID NO. 107.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out in the sequence listing as SEQ ID NO. 108 which corresponds to the "Q" sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Anergic T-cell; Human Microvascular Endothelial Cells, fract. A.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:11 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1612 of SEQ ID NO:11, b is an integer of 15 to 1626, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:11, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 2

30 The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gi1572819 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "similar to the RAS gene family [Caenorhabditis elegans]." A partial alignment demonstrating the observed homology is shown immediately below.

>gi1572819 similar to the RAS gene family [Caenorhabditis elegans]
Length = 625

Plus Strand HSPs:

Score = 540 (190.1 bits), Expect = 8.5e-111, Sum P(3) = 8.5e-111
Identities = 119/286 (41%), Positives = 172/286 (60%), Frame = +2

5 Q: 728 LFIQGRHETTWITLRRFGYSDALELTADYLSPLIHVPPGCSTELNHLGYQFVQRVFEKH 907
 LFI+RGRHETTW +LR+FGY +L+L+ DYL P I +P GCSTEL+ G QFV +FEK+
10 S: 261 LFIERGRHETTWAVLRKFGYETSLKLSedylyPRITIPVGCSTELSPQGVQFVSALFEKY 320

 Q: 908 DQDRDGALSPVELQSLFSVFPAAPWGPPELPRVTRT-EAGRLPLHGYLCQWTLVTYLDVRS 1084
 D+D+DG LSP ELQ+LFSV P + + T + G L +GY+ W + T +++
 S: 321 DEDKDGCLSPSELQNLFSVCPVPVITKDNILALETNQRGWLTYNGYMAWNNMTTLINLTQ 380

15 Q: 1085 CLGHLGYLGYPT-LCEQDQA----HAITVTREKRLDQEKQGTQKXLLCKVVGARGVGKS 1249
 L YLG+P +A +I VTRE++ D E T R C VVGA+ GK+
 S: 381 TFEQLAYLGFVGRSGPGRAGNTLDSIRVTRERKKDLENHGTDRKVFQCLVVGAKDAGKT 440

20 Q: 1250 AFLQAFGLGRGLGH--QDTREQPPGYAIDTVQVNGQEKYLILCEVGT----DGLLATSLDA 1411
 F+Q+ GRG+ "Q" R P + I+ V+V + KYL+L EV D L S +
 S: 441 VFMQSLAGRMADVAQIGRRHSP-FVINRVVKEESKYLLREVDVLSPODAL--GSGET 497

 Q: 1412 TCDVACLMDGSDPKSFAHCASVYKHHYMDGQTPCLFVSSKADLPEVSR 1558
 + DV ++D S+P SFA CA+VY+ ++ +TPC+ +++K + EV +
25 S: 498 SADVVAFLYDISNPDSFAFCATVYQKYFYRTKTPCVMIATKVEREEVDQ 546

The segment of gill572819 that is shown as "S" above is set out in the sequence listing as SEQ ID NO. 109.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid
30 sequence set out in the sequence listing as SEQ ID NO. 110 which corresponds to the "Q"
sequence in the alignment shown above (gaps introduced in a sequence by the computer are,
of course, removed).

It has been discovered that this gene is expressed primarily in the following
tissues/cDNA libraries: Primary Dendritic Cells, lib 1; Soares infant brain INIB and to a
35 lesser extent in breast lymph node CDNA library; Soares_pregnant_uterus_NbHPU; Human
adult testis, large s; Human Synovial Sarcoma; Healing groin wound, 7.5 hours post incision;
Colon Normal II; Human Testes Tumor; Human Cerebellum; Soares ovary tumor NbHOT;
Human Tonsils, lib I; Human Pituitary, subtracted; Human Neutrophils, Activated, re-
excision; Smooth Muscle Serum Treated, Norm; Human Primary Breast Cancer; Human
40 Soleus; B Cell lymphoma; Alzheimers, spongy change; Human Osteoclastoma Stromal Cells
- unamplified; Human Whole Brain #2 - Oligo dT > 1.5Kb; Salivary Gland, Lib 2; Healing
groin wound, 6.5 hours post incision; Fetal Liver, subtraction II; Breast Cancer Cell line,
angiogenic; Soares_pineal_gland_N3HPG; Human Brain, Striatum; Human Uterine Cancer;
Human Pancreas Tumor; Soares adult brain N2b5HB55Y; Resting T-Cell Library,II; H.
45 Frontal cortex,epileptic,re-excision; Human Eosinophils; Early Stage Human Brain; Human
Neutrophil, Activated; Human Adult Pulmonary,re-excision; Colon Normal III; Human

Microvascular Endothelial Cells, fract. A; Bone Marrow Cell Line (RS4,11); T cell helper II; Soares fetal liver spleen INFLS; Soares_pineal_gland_N3HPG and Stratagene corneal stroma (#937222).

The tissue distribution in immune cells indicates the polynucleotides and polypeptides corresponding to this gene would be useful for the diagnosis and treatment of a variety of immune system disorders. Representative uses are described in the "Immune Activity" and "Infectious Disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. Involvement in the regulation of cytokine production, antigen presentation, or other processes suggests a usefulness for treatment of cancer (e.g. by boosting immune responses). Expression in cells of lymphoid origin, indicates the natural gene product would be involved in immune functions. Therefore it would also be useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Furthermore, the protein may also be used to determine biological activity, raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:12 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2383 of SEQ ID NO:12, b is an integer of 15 to 2397, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:12, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 3

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Soares fetal liver spleen 1NFLS and to a lesser extent in Salivary Gland, Lib 2; Soares ovary tumor NbHOT; Pancreas Islet Cell Tumor; Osteoblasts; Soares_fetal_heart_NbHH19W; Soares_fetal_liver_spleen_1NFLS_S1; Soares_senescent_fibroblasts_NbHSF; Human Thymus Tumor; Soares_multiple_sclerosis_2NbHMSP; Weizmann Olfactory Epithelium; prostate-edited; Human Fetal Spleen; Smooth muscle, control, re-excision; H. Whole Brain #2, re-excision; H Female Bladder, Adult; Healing groin wound, 7.5 hours post incision; Human Fetal Epithelium (Skin); H. Kidney Cortex, subtracted; Healing groin wound, 6.5 hours post incision; Breast Cancer Cell line, angiogenic; Human Brain, Striatum; Human Fetal Dura Mater; Ulcerative Colitis; Hepatocellular Tumor, re-excision; Stratagene liver (#937224); Human Gall Bladder; Soares_fetal_heart_NbHH19W; H. Frontal cortex,epileptic,re-excision; Brain frontal cortex; Human Fetal Lung III and Human Microvascular Endothelial Cells, fract. A.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 61 as residues: Gln-25 to Arg-31.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:13 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the

present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 800 of SEQ ID NO:13, b is an integer of 15 to 814, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:13, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 4

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Soares_pregnant_uterus_NbHPU and to a lesser extent in LNCAP prostate cell line; Human Stomach, re-excision; Soares_parathyroid_tumor_NbHPA; Nine Week Old Early Stage Human; 12 Week Old Early Stage Human, II; Early Stage Human Brain; Human Microvascular Endothelial Cells, fract. A; Soares_pineal_gland_N3HPG and Smooth muscle, control.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:14 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 627 of SEQ ID NO:14, b is an integer of 15 to 641, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:14, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 5

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gnl|PID|e1266315 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "membrane glycoprotein gp36 [Homo sapiens]." A partial alignment demonstrating the observed homology is shown immediately below.

```
>gnl|PID|e1266315 (AJ225022) membrane glycoprotein gp36 [Homo sapiens]
```

```

5      Score = 665 (234.1 bits), Expect = 1.0e-64, P = 1.0e-64
      Identities = 137/159 (86%), Positives = 137/159 (86%), Frame = +1

10     Q:   193 MWKVSALLFVLGSASLWVLAEGASTGQPEDDTETTGLEGGVAMPGAEDDVVTPGTSEDY 372
      S:     1 MWKVSALLFVLGSASLWVLAEGASTGQPEDDTETTGLEGGVAMPGAEDDVVTPGTSEDY 60

15     Q:   373 KSGLTTLVATSVNSVTGIRIEDLPTSESTVHAQEQSPSATASNVATSHSTEKVDGDTQTT 552
      S:     61 KSGLTTLVATSVNSVTGIRIEDLPTSESTVHAQEQSPSATASNVATSHSTEKVDGDTQTT 120

      Q:   553 VEKDGLSTVTXXXXXXXXXXXXXXXXXXXXXXXXMRKMSGR 669
      S:     121 VEKDGLSTVT VLGIIIVGVLLAIGFIGGIIIVVMRKMSGR 159

```

25 Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out in the sequence listing as SEQ ID NO. 112 which corresponds to the "Q" sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 63 as residues: Met-1 to Arg-9.

40 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:15 and may have been publicly available prior to conception of the present invention.

Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2149 of SEQ ID NO:15, b is an integer of 15 to 2163, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:15, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 6

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Human Thyroid; human tonsils; Human Microvascular Endothelial Cells, fract. A.

The tissue distribution suggests that the protein product of this clone is useful for the diagnosis and/or treatment of disorders involving the vasculature. Elevated expression of this gene product by endothelial cells suggests that it may play vital roles in the regulation of endothelial cell function; secretion; proliferation; or angiogenesis. Alternately, this may represent a gene product expressed by the endothelium and transported to distant sites of action on a variety of target organs. Expression of this gene product by hematopoietic cells also suggests involvement in the proliferation; survival; activation; or differentiation of all blood cell lineages.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:16 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1298 of SEQ ID NO:16, b is an integer of 15 to 1312, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:16, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 7

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: NTERA2 teratocarcinoma cell line+retinoic acid (14 days); Jurkat T-

Cell, S phase; NTERA2, control; Stratagene NT2 neuronal precursor 937230; Human Microvascular Endothelial Cells, fract. A; Keratinocyte; Primary Dendritic Cells, lib 1.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:17 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1712 of SEQ ID NO:17, b is an integer of 15 to 1726, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:17, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 8

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gnl|PID|e303266 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "hypothetical protein [Bos taurus]." A partial alignment demonstrating the observed homology is shown immediately below.

```

>gnl|PID|e303266 hypothetical protein [Bos taurus] >sp|O18975|O18975
  HYPOTHETICAL 16.6 KD PROTEIN (FRAGMENT).
  Length = 145

Plus Strand HSPs:

Score = 231 (81.3 bits), Expect = 1.1e-17, P = 1.1e-17
Identities = 46/64 (71%), Positives = 47/64 (73%), Frame = +1

Q:      7 AEELKRNAETGNLPHSYRLIXXXXXXXXXXXXXXXXXXDVYDIKKQAWFTYNDLEVSKIQE 186
        AEELKRNAETGNLPHSYRLI                                DVYDIKKQAWFTYNDLEVSKIQE
S:      80 AEELKRNAETGNLPHSYRLISVSHIGSTSSSGHYISDVYDIKKQAWFTYNDLEVSKIQE 139

Q:      187 AAVQ 198
        A+VQ
S:      140 ASVQ 143
  
```

The segment of gnl|PID|e303266 that is shown as "S" above is set out in the sequence listing as SEQ ID NO. 113.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out in the sequence listing as SEQ ID NO. 114 which corresponds to the "Q" sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

5 It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Human Microvascular Endothelial Cells, fract. A; H. Whole Brain #2, re-excision.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 66 as residues: Arg-48 to Cys-57.

10 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:18 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly,
15 preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1992 of SEQ ID NO:18, b is an integer of 15 to 2006, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:18, and where b is greater than or equal to a + 14.

20

FEATURES OF PROTEIN ENCODED BY GENE NO: 9

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Human Microvascular Endothelial Cells, fract. A and to a lesser extent in Soares_fetal_liver_spleen_1NFLS_S1; Stratagene lung carcinoma 937218;
25 Soares_fetal_heart_NbHH19W; Human Primary Breast Cancer, re-excision; CHME Cell Line, untreated; Stomach cancer (human), re-excision; Stromal cell TF274; Human adult testis, large s; Human Testes Tumor; human tonsils; NCI_CGAP_GCB1; Stratagene NT2 neuronal precursor 937230; T Cell helper I; Hodgkin's Lymphoma II; Soares fetal liver spleen INFLS and Primary Dendritic Cells, lib 1.

30 When tested against U937 Myeloid cell lines, supernatants removed from cells containing this gene activated the GAS assay. Thus, it is likely that this gene activates myeloid cells through the Jak-STAT signal transduction pathway. The gamma activating sequence (GAS) is a promoter element found upstream of many genes which are involved in the Jak-STAT pathway. The Jak-STAT pathway is a large, signal transduction pathway

involved in the differentiation and proliferation of cells. Therefore, activation of the Jak-STAT pathway, reflected by the binding of the GAS element, can be used to indicate proteins involved in the proliferation and differentiation of cells.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:19 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1593 of SEQ ID NO:19, b is an integer of 15 to 1607, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:19, and where b is greater than or equal to a + 14.

15 FEATURES OF PROTEIN ENCODED BY GENE NO: 10

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Bone Marrow Cell Line (RS4,11) and to a lesser extent in NTERA2, control; Smooth muscle.control; Soares_senescent_fibroblasts_NbHSF; Stratagene pancreas (#937208); Primary Dendritic Cells, lib 1; Soares infant brain 1NIB; LNCAP untreated; Human colon carcinoma (HCC) cell line, remake; Smooth Muscle Serum Treated, Norm; Human Lung; Smooth Muscle- HASTE normalized; Jurkat T-Cell, S phase; Brain Frontal Cortex, re-excision; Human Testes Tumor, re-excision; Hemangiopericytoma; NCI_CGAP_Lu5; NCI_CGAP_GCB1; Early Stage Human Brain; Human Microvascular Endothelial Cells, fract. A; Human Bone Marrow, treated; Keratinocyte and Human 8 Week Whole Embryo.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 68 as residues: Gly-18 to Gly-23.

Preferred polypeptides comprise the following amino acid sequence:

MPDAEVLWVAVVEAVFLRTQGSSSPQLQAGQGEEDVELQEAKGSSSESREDSAGR
LMGSFHGMAPASSDWDQLGWEQEQTALPNWCGTSTSEFLNSLRGTWLLALELHTQ
TISLKGRSLFRVPEVPIKSPSCQLWCLICGSVCCCRFGRDGWAS (SEQ ID NO:) and
MAALLLTAKRSSSLCQWAAERACDYWPVTCRGTVLPSWIQRPWETLRSSPTVSPKS

AAAYGPTNETPKLTGWTFNGHFWDPEETSSLQAYCLSVKFQSKPCSSKGIQEFRA

SPTPVR (SEQ ID NO:).

Polynucleotides encoding these polypeptides are also encompassed by the invention.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:20 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1388 of SEQ ID NO:20, b is an integer of 15 to 1402, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:20, and where b is greater than or equal to a + 14.

15 FEATURES OF PROTEIN ENCODED BY GENE NO: 11

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Human OB MG63 control fraction I.

Preferred polypeptides comprise the following amino acid sequence:

MDRVGIFREHPASEAAAAPSSGITLPSPTLHWGVSYCASCWRTQLGAEAVGSERRLE

20 KVCGAVCENS GARSLLEKPKRKSYILNHSVQFREPCQCVSFSSPKTLLPQVIVGR

(SEQ ID NO:) and

MSWGS GGGGKGRDDLPREKPTTCPVFDRLFDIFAKIPLVESQASCAGIGIAASHWRL

DCSVDSCKWSCLSLILIARGFQMKRKLKEEWPSFSKFCGSSKDFYVHTLSSLVLGTFY

LICLGSACFPGCSLQTWNSVNPRLTDLEPEVSFILMV (SEQ ID NO:). Polynucleotides

25 encoding these polypeptides are also encompassed by the invention.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:21 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the

present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1207 of SEQ ID NO:21, b is an integer of 15 to 1221, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:21, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 12

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Soares infant brain 1NIB and to a lesser extent in Brain Amygdala Depression; Human OB MG63 treated (10 nM E2) fraction I; Human Fetal Bone; Human Bone Marrow Stromal Fibroblast; Macrophage-oxLDL, re-excision; Brain frontal cortex; Smooth muscle, control; Monocyte activated; Soares_NhHMPu_S1; Keratinocyte; Soares fetal liver spleen 1NFLS and Primary Dendritic Cells, lib 1.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:22 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1241 of SEQ ID NO:22, b is an integer of 15 to 1255, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:22, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 13

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gn|PID|d1035774 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "HRIHFB2018 [Homo sapiens]." A partial alignment demonstrating the observed homology is shown immediately below.

```
>gnl|PID|d1035774 (AB015332) HRIHFB2018 [Homo sapiens] >gnl|PID|d1035774
```

(AB015332) HRIHFB2018 [Homo sapiens] >sp|D1035774|D1035774
 HRIHFB2018 PROTEIN (FRAGMENT).
 Length = 373

5 Plus Strand HSPs:

Score = 1144 (402.7 bits), Expect = 2.5e-122, Sum P(2) = 2.5e-122
 Identities = 220/287 (76%), Positives = 220/287 (76%), Frame = +1

10 Q: 67 EQQKPASSDVVLPATMSYTGFGVQGGSETTLQSTYSDTSAQPTCDYGYGTWNSXXXXXXXXXX 246
 EQQKPASSDVVLPATMSYTGFGVQGGSETTLQSTYSDTSAQPTCDYGYGTWNS
 S: 1 EQQKPASSDVVLPATMSYTGFGVQGGSETTLQSTYSDTSAQPTCDYGYGTWNSGTNRGYEGY 60

15 Q: 247 XXXXXXXXXXXXXXXXXXXXMATSHSWEMPXXXXXXXXXXXXXXXXXXVLSRINQRLDMVPHL 426
 MATSHSWEMP VLSRINQRLDMVPHL
 S: 61 GYGYGYGDNTTNYGYGMATSHSWEMPSSDTNANTSASGSASADSVLSRINQRLDMVPHL 120

20 Q: 427 ETDMMQGGVYGSGGERYSYSCDSRAVLSELDYRSGYDYSELDPPEMAYEGQYDAYR 606
 ETDMMQGGVYGSGGERYSYSCDSRAVLSELDYRSGYDYSELDPPEMAYEGQYDAYR
 S: 121 ETDMMQGGVYGSGGERYSYSCDSRAVLSELDYRSGYDYSELDPPEMAYEGQYDAYR 180

25 Q: 607 DQFRMRGNDTFGPRAQWARDARSGRPMASGYGRMWEDPMGARGQCMGASRLPSLFSQN 786
 DQFRMRGNDTFGPRAQWARDARSGRPMASGYGRMWEDPMGARGQCMGASRLPSLFSQN
 S: 181 DQFRMRGNDTFGPRAQWARDARSGRPMASGYGRMWEDPMGARGQCMGASRLPSLFSQN 240

Q: 787 I IPEYGMXXXXXXXXXXXXXXXXXXXXXXXXXMKQMRRTWKTWTTADFR 927
 I IPEYGM MKQMRRTWKTWTTADFR
 S: 241 I IPEYGMFQGMRGGAFFPGGSRFGFGNGMKQMRRTWKTWTTADFR 287

30 The segment of gnlIPIDid1035774 that is shown as "S" above is set out in the
 sequence listing as SEQ ID NO. 115.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid
 sequence set out in the sequence listing as SEQ ID NO. 116 which corresponds to the "Q"
 sequence in the alignment shown above (gaps introduced in a sequence by the computer are,
 35 of course, removed).

It has been discovered that this gene is expressed primarily in the following
 tissues/cDNA libraries: Soares adult brain N2b5HB55Y and to a lesser extent in Human
 Osteoarthritic Cartilage Fraction III; Human Manic Depression Tissue; Brain Frontal Cortex,
 re-excision; T-Cell PHA 16 hrs; Ulcerative Colitis; NTERA2, control; Human Eosinophils;
 40 Colon Normal II; breast lymph node CDNA library; Human Bone Marrow, treated; Primary
 Dendritic Cells, lib 1 and Soares infant brain 1NIB.

Many polynucleotide sequences, such as EST sequences, are publicly available and
 accessible through sequence databases. Some of these sequences are related to SEQ ID
 NO:23 and may have been publicly available prior to conception of the present invention.
 45 Preferably, such related polynucleotides are specifically excluded from the scope of the
 present invention. To list every related sequence would be cumbersome. Accordingly,
 preferably excluded from the present invention are one or more polynucleotides comprising a

nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1628 of SEQ ID NO:23, b is an integer of 15 to 1642, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:23, and where b is greater than or equal to a + 14.

5

FEATURES OF PROTEIN ENCODED BY GENE NO: 14

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gil971762 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "60S ribosomal protein L11 [Sus scrofa]." A partial alignment demonstrating the observed homology is shown immediately below.

15

>gi|971762 60S ribosomal protein L11 [Sus scrofa]
Length = 165

Minus Strand HSPs:

20

Score = 187 (65.8 bits), Expect = 6.9e-14, P = 6.9e-14
Identities = 42/62 (67%), Positives = 49/62 (79%), Frame = -1

25

Q: 509 TPCYPLPAARYTVRSFGIRRNEKIAVHCTVRGAKAEIILEKGLKVSLIP*WSDIDQHSFS 330
TP + ARYTVRSFGIRRNEKIAVHCTVRGAKAEIILEKGLKV ++ +++FS
S: 46 TPVFS--KARYTVRSFGIRRNEKIAVHCTVRGAKAEIILEKGLKVREY---ELRKNNFS 99
Q: 329 NT 324
+T
S: 100 DT 101

30

The segment of gil971762 that is shown as "S" above is set out in the sequence listing as SEQ ID NO. 117. Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein. Assays for determining such activities are also known in the art, some of which have been described elsewhere herein.

35

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out in the sequence listing as SEQ ID NO. 118 which corresponds to the "Q" sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

40

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Hemangiopericytoma; Bone Marrow Cell Line (RS4,11) and to a lesser extent in H.Leukocytes, normalized cot 5B; Kidney cancer; Human White Adipose;

Human Adult Pulmonary; Human Manic Depression Tissue; human ovarian cancer; Human Substantia Nigra; Activated T-Cell (12hs)/Thiouridine labelled Eco; Smooth muscle, control; Spleen, Chronic lymphocytic leukemia and Human 8 Week Whole Embryo.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:24 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 918 of SEQ ID NO:24, b is an integer of 15 to 932, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:24, and where b is greater than or equal to a + 14.

15 **FEATURES OF PROTEIN ENCODED BY GENE NO: 15**

It has been discovered that this gene is expressed primarily in Human Manic Depression Tissue.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:25 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1050 of SEQ ID NO:25, b is an integer of 15 to 1064, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:25, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 16

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Human Hippocampus, subtracted; Soares_fetal_heart_NbHH19W and to a lesser extent in Human Manic Depression Tissue; T-Cell PHA 16 hrs and Resting T-Cell Library, II.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:26 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1266 of SEQ ID NO:26, b is an integer of 15 to 1280, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:26, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 17

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Soares infant brain 1NIB and to a lesser extent in Soares breast 3NbHBst; Human Manic Depression Tissue; H. Frontal cortex.epileptic, re-excision; Human Eosinophils; Human Endometrial Tumor; Soares fetal liver spleen 1NFLS; Human Synovium; Soares_NhHMPu_S1; Stratagene HeLa cell s3 937216; Synovial hypoxia-RSF subtracted; Soares_fetal_lung_NbHL19W; H. Kidney Medulla, re-excision; Human Uterine Cancer; Human Pancreas Tumor; Human Ovarian Cancer Reexcision; Human Fetal Brain; Human Whole Six Week Old Embryo; Stratagene colon (#937204); Stratagene HeLa cell s3 937216; NCI_CGAP_Co3; NCI_CGAP_Pr2; Colon Normal II; Human Fetal Heart; Monocyte activated; Human B Cell Lymphoma; Hodgkin's Lymphoma II and Human Cerebellum.

Preferred polypeptides comprise the following amino acid sequence:

MLLAEISSVAHQDGSFCPIVMCGDFNSVPGSPLYSFKEGKLNYEGLPIGKVSGQEQ
SSRGQRILSIWPPNLGISQNCVYEVQQVPKVEKTDSDLTQTQLKQTEVLVTAEKLS
NLQHHFSLSSVYSHYFPDTGIPEVTTCHSRSAITVDYIFYSAEKEDVAGHPGAEVALV
GGLKLLARLSPLTEQDLWTVNGLPNENNSSDHLPLAKFRLEL (SEQ ID NO:).

Polynucleotides encoding these polypeptides are also encompassed by the invention.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:27 and may have been publicly available prior to conception of the present invention.

Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 3606 of SEQ ID NO:27, b is an integer of 15 to 3620, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:27, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 18

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gi54806 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "TIS7 protein [Mus sp.]," which is a gene induced by the tumor promoter TPA. A partial alignment demonstrating the observed homology is shown immediately below.

```
>gi|54806 TIS7 protein (AA 1-449) [Mus sp.] >pir|A44989|A44989
interferon-related protein TIS7 - mouse >sp|P19182|PC4_MOUSE
INTERFERON-RELATED PROTEIN PC4 (TPA INDUCED SEQUENCE 7) (TIS7
PROTEIN).
Length = 449
```

Plus Strand HSPs:

Score = 1375 (484.0 bits), Expect = 1.8e-190, Sum P(3) = 1.8e-190
Identities = 266/281 (94%), Positives = 277/281 (98%), Frame = +1

```
Q: 7 SEDGPEVLDEEGTQEDLEYKLGKGLIDLTLDKSAKTRQAALLEGIKNALASKMLYEFILERR 186
+EDGPEVLDEEGTQEDLEYKLGKGLIDLTLDKSAKTRQAALLEG+KNAL-SK+LYEF+LERR
S: 62 AEDGPEVLDEEGTQEDLEYKLGKGLIDLTLDKSAKTRQAALLEGVKNALSSKVLVEFVLERR 121

Q: 187 MTLTDSIERCLKKGSDEQRAAAALASVLCIQLGPGIESEEILKTLGPILKKIICDGSAS 366
MTLTDSIERCLKKGSDEQRAAAA+ASVLCIQLGPG ESEEILKTLGPILKKIICD+AS
S: 122 MTLTDSIERCLKKGSDEQRAAAAVASVLCIQLGPGFESEEILKTLGPILKKIICDGAAS 181

Q: 367 MQARQTCATCFGVCCFIATDDITELYSTLECLNIFTKSYLKEKDTTVICSTPNTVLHIS 546
+QARQTCATCFGVCCFIATDDITELYSTLEC ENIFTKSYLKEKDT V CSTPNTVLHIS
S: 182 IQARQTCATCFGVCCFIATDDITELYSTLECFENIFTKSYLKEKDTNVTCTPNTVLHIS 241

Q: 547 SLLAWTLLLTICPINEVKKKLEMHFKLPSLLSCDDVMRIAAGESLALLFELARGIESD 726
SLLAWTLLLTICPINEVKKKLE+HFHKLPSLLSCDDVMRIAAGESLALLFELARG+ESD
S: 242 SLLAWTLLLTICPINEVKKKLELHFHKLPSLLSCDDVMRIAAGESLALLFELARGMESD 301

Q: 727 FFYEDMESLTQMLRALATDGNKHKRAKVDKQRQSVFRDVL 849
FFYEDM+SLTQMLRALATDGNKHKRAKVDKQRQSVFRDVL
S: 302 FFYEDMDSLTQMLRALATDGNKHKRAKVDKQRQSVFRDVL 342
```

Score = 348 (122.5 bits), Expect = 1.8e-190, Sum P(3) = 1.8e-190
Identities = 65/74 (87%), Positives = 68/74 (91%), Frame = +3

Q: 831 FQRCPEAVEERDFPTETIKFGPERMYIDCWVKKHTYDTFKEVLGSGMQYHLQSNFLRNV 1010
 F+ AVEERDFPTET+KFGPERMYID WVKKHTYDTFKEVLGSGMQYHLQ+NEFLRNV
 S: 337 FRDVLRAVEERDFPTETVKFGPERMYIDSWVKKHTYDTFKEVLGSGMQYHLQTNEFLRNV 396
 5 Q: 1011 FELGPPVMLDAARL 1052
 FELGPPVMLDAA L
 S: 397 FELGPPVMLDAATL 410
 10 Score = 168 (59.1 bits), Expect = 1.8e-190, Sum P(3) = 1.8e-190
 Identities = 35/40 (87%), Positives = 36/40 (90%), Frame = +2
 Q: 1046 TLKR*RFS-FERHLYNSAAFKARTKARSKCRDKRADVGEF 1162
 TLK + S FERHLYNSAAFKARTKARSKCRDKRADVGEF
 15 S: 409 TLKTMKISRFERHLYNSAAFKARTKARSKCRDKRADVGEF 448

The segments of gil54806 that are shown as "S" above are set out in the sequence listing as SEQ ID NO. 119, SEQ ID NO. 121 and SEQ ID NO. 123. Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein.
 20 Assays for determining such activities are also known in the art, some of which have been described elsewhere herein. Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out in the sequence listing as SEQ ID NO. 120, SEQ ID NO. 122 and/or SEQ ID NO. 124 which correspond to the "Q" sequences in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

25 It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Human Osteoclastoma Stromal Cells - unamplified and to a lesser extent in Soares placenta Nb2HP; Human Gall Bladder; Soares_fetal_liver_spleen_1NFLS_S1; L428 cell line; Soares_placenta_8to9weeks_2NbHP8to9W; Human Adipose; Bone Marrow Stromal Cell,
 30 untreated; NTERA2, control; Stratagene lung (#937210); Stratagene colon (#937204); Soares_parathyroid_tumor_NbHPA; Soares_multiple_sclerosis_2NbHMSP; Bone marrow; Soares fetal liver spleen 1NFLS; Soares infant brain 1NIB; Normal Prostate; Human Prostate, subtracted; Healing Abdomen wound, 70&90 min post incision; Human Placenta; Early Stage Human Lung, subtracted; Pancreas Tumor PCA4 Tu; H Female Bladder, Adult; Human
 35 Adipose Tissue, re-excision; Human Osteoclastoma, re-excision; Synovial hypoxia; Human Manic Depression Tissue; Prostate BPH; KMH2; T-Cell PHA 16 hrs; Human Pancreas Tumor; Soares_pregnant_uterus_NbHPU; Soares breast 2NbHBst; Pancreas Islet Cell Tumor; Macrophage-oxLDL, re-excision; H. Frontal cortex, epileptic, re-excision; Human Eosinophils; Human Substantia Nigra; Adipocytes; Dendritic cells, pooled; Human Fetal
 40 Heart; Human Neutrophil, Activated; Human Osteoclastoma; T Cell helper I; Smooth

muscle, control; Human Bone Marrow, treated; Human Testes; T cell helper II; Keratinocyte and Human Cerebellum.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:28 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1882 of SEQ ID NO:28, b is an integer of 15 to 1896, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:28, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 19

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gil439877 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "reverse transcriptase [Homo sapiens]." A partial alignment demonstrating the observed homology is shown immediately below.

```

>gil439877 reverse transcriptase [Homo sapiens]
      Length = 361

      Minus Strand HSPs:

Score = 138 (48.6 bits), Expect = 4.2e-26, Sum P(4) = 4.2e-26
Identities = 37/92 (40%), Positives = 51/92 (55%), Frame = -1

Q:   431 KVGKNNRHRFSKEEIQIAKKDKKRCSTLSVIMKMQMKTTRHCFTPNMGWKQKVREKQVM 255
      K K+ NRHFSKE+I AKK K+CS I +MQ+KTT R+ TP + + +K
S:   157 KWAKDMNRHFSKEDIYAAKHKMKCSPSLAIREMQIKTTMRYHLTPV---RMAIIKKSGN 213

Q:   254 AKMWRNQN SQG----CRKGREMVQTLGKAGWRLK 162
      + WR G C ++VQ L K+ WR L+
S:   214 NRCWRGCGEIGTLLHCWNNCLVQPLWKSVMRFLR 248
  
```

The segment of gil439877 that is shown as "S" above is set out in the sequence listing as SEQ ID NO. 125. Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein. Assays for determining such activities are also known in the art, some of which have been described elsewhere herein.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out in the sequence listing as SEQ ID NO. 126 which corresponds to the "Q" sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

5 It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Human Manic Depression Tissue; 12 Week Old Early Stage Human; Human T-Cell Lymphoma.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID
10 NO:29 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between
15 1 to 1347 of SEQ ID NO:29, b is an integer of 15 to 1361, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:29, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 20

20 It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Soares infant brain 1NIB and to a lesser extent in Soares_parathyroid_tumor_NbHPA; H. Frontal cortex,epileptic.re-excision; Spleen, Chronic lymphocytic leukemia; Soares adult brain N2b4HB55Y; Alzhcimers, spongy change; Glioblastoma; LNCAP prostate cell line; Healing groin wound, 6.5 hours post incision;
25 Human Manic Depression Tissue; Soares_fetal_heart_NbHH19W; Human Heart; Brain frontal cortex; Primary Dendritic cells,frac 2; human tonsils and Human Amygdala.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 78 as residues: Pro-4 to Gly-13, Ala-42 to Ser-50.

Many polynucleotide sequences, such as EST sequences, are publicly available and
30 accessible through sequence databases. Some of these sequences are related to SEQ ID NO:30 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a

nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1309 of SEQ ID NO:30, b is an integer of 15 to 1323, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:30, and where b is greater than or equal to a + 14.

5

FEATURES OF PROTEIN ENCODED BY GENE NO: 21

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. pirlS72481|S72481 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "probable transposase - human transposable element MER37." A partial alignment demonstrating the observed homology is shown immediately below.

```

15  >pirlS72481|S72481 probable transposase - human transposable element MER37
    >pirlS72486|S72486 putative transposase - human transposon
MER37
      (fragment) (SUB 177-349)
      Length = 454

20  Minus Strand HSPs:

      Score = 340 (119.7 bits), Expect = 5.3e-54, Sum P(3) = 5.3e-54
      Identities = 86/203 (42%), Positives = 110/203 (54%), Frame = -2

25  Q:  729 SQVVNAMKSS*RLKVLHRTHE**GKH--LISDIERVLWVWIEDQISHNVSLRS--QS 562
      SQVVNA +   ++K          K  LI+D+E+VL+VWIEDQ SHN+ LS+S  QS
S:  48  SQVVNAKEKFLKEIKSATPVNTRMIRKQNSLIADMEKVLVWVWIEDQTSJNIPLSQSLIQS 107

30  Q:  561 KALTLFNSLXXXXXXXXXXXXXXXXXWFMRFKRSYLRNIKVLXXXXXXXXXXXXXYPE 382
      KALTLFNS+          WFMRFK+RS+L NIKV          YPE
S:  108 KALTLFNSMKAERGEEAAEEKLEASRGWFMRFKERSHLHNIKVQGEAASADVEAAASYPE 167

35  Q:  381 DLAGIIDKAXYTKQHIFTVNQXAIEXRCCLGLS*LESTRHRLFLKLERTS*PLV-GANA 205
      DLA IID+ YTKQ IF V++ A   +           +   K +   L+ GANA
S:  168 DLAKIIDEGGYTKQIFNVDETAFYWKMPKPSRTFIAREEKSMPGFKASKDRLTLLLGANA 227

40  Q:  204 ASDTKLKLVLIFPSQNPTALQNY 136
      A D KLK +LI+ S+NP AL+NY
S:  228 AGDFKLKPMLIYHSENPRALKNY 250

```

The segment of pirlS72481|S72481 that is shown as "S" above is set out in the sequence listing as SEQ ID NO. 127.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out in the sequence listing as SEQ ID NO. 128 which corresponds to the "Q" sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

45

It has been discovered that this gene is expressed primarily in Human Manic Depression Tissue.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 79 as residues: Pro-43 to Cys-48.

5 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:31 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly,
10 preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2069 of SEQ ID NO:31, b is an integer of 15 to 2083, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:31, and where b is greater than or equal to a + 14.

15

FEATURES OF PROTEIN ENCODED BY GENE NO: 22

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Human Manic Depression Tissue; Human Fetal Brain.

20 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:32 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly,
25 preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1724 of SEQ ID NO:32, b is an integer of 15 to 1738, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:32, and where b is greater than or equal to a + 14.

30 FEATURES OF PROTEIN ENCODED BY GENE NO: 23

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Stratagene lung (#937210) and to a lesser extent in Human Prostate Cancer, Stage B2; Human Manic Depression Tissue; Stratagene HeLa cell s3 937216 and T cell helper II.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:33 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 903 of SEQ ID NO:33, b is an integer of 15 to 917, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:33, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 24

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Soares infant brain 1NIB and to a lesser extent in Human Hypothalamus, schizophrenia, re-excision; Human Brain, Striatum; Soares_NhHMPu_S1; Human Cerebellum; Soares fetal liver spleen 1NFLS; Human Amygdala, re-excision; Human Manic Depression Tissue; Soares_multiple_sclerosis_2NbHMSP; Human Adrenal Gland Tumor; H. Frontal cortex, epileptic, re-excision and Human Substantia Nigra.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 82 as residues: Gln-13 to Tyr-20.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:34 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1758 of SEQ ID NO:34, b is an integer of 15 to 1772, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:34, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 25

It has been discovered that this gene is expressed primarily in H. Meningioma, M6.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:35 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 785 of SEQ ID NO:35, b is an integer of 15 to 799, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:35, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 26

It has been discovered that this gene is expressed primarily in H. Meningima, M1.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 84 as residues: Thr-12 to Glu-21.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:36 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1716 of SEQ ID NO:36, b is an integer of 15 to 1730, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:36, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 27

It has been discovered that this gene is expressed primarily in H. Meningima, M1.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:37 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a

nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1006 of SEQ ID NO:37, b is an integer of 15 to 1020, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:37, and where b is greater than or equal to a + 14.

5

FEATURES OF PROTEIN ENCODED BY GENE NO: 28

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Activated T-cell(12h)/Thiouridine-re-excision and to a lesser extent in Human Colon Carcinoma (HCC) cell line; H. Meningima, M1; Stratagene fetal spleen
10 (#937205); H. Frontal cortex,epileptic,re-excision; Brain frontal cortex and T cell helper II.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:38 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the
15 present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 943 of SEQ ID NO:38, b is an integer of 15 to 957, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:38, and where b is greater than or
20 equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 29

It has been discovered that this gene is expressed primarily in H. Meningima, M1.

Many polynucleotide sequences, such as EST sequences, are publicly available and
25 accessible through sequence databases. Some of these sequences are related to SEQ ID NO:39 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a
30 nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1020 of SEQ ID NO:39, b is an integer of 15 to 1034, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:39, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 30

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gnl|PID|e314281 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "Pro-Pol-dUTPase polyprotein [Murine endogenous retrovirus]." A partial alignment demonstrating the observed homology is shown immediately below.

```

10  >gnl|PID|e314281 Pro-Pol-dUTPase polyprotein [Murine endogenous
    retrovirus]
    >bbs|132484 fibulin gene homolog [mice, Peptide Partial, 78
    aa]
        [Mus sp.] (SUB 380-457)
        Length = 1182
15  Minus Strand HSPs:
    Score = 245 (86.2 bits), Expect = 4.7e-19, Sum P(2) = 4.7e-19
    Identities = 73/211 (34%), Positives = 108/211 (51%), Frame = -3
20  Q: 700 PMAPWGFLHRQLTE*KRSRKRAL*GLG*HD-----MLA*AKKMDWCCTKDTP--KITLG 545
    P+A WG + QLTE +++R G + A + + KDT K +
    S: 598 PIASWGPYPDQLTEEEKTRAWFTDGSARYAGTTQKWTAALQPLSGTTLKDTGERKSSQW 657
25  Q: 544 ANSSPV--VICFVSGEKRPKVRIYDF*VVENGLVAWSRASREQH*RIETKHV*GRGMWL 371
    A V V+ FV +K P VR++TD V NGL WS ++ + +I K + GR MW+
    S: 658 AELRAVHMLVLFVCKKKWPDVRLFTDSWAVANGLAGWSGTWKDHNWKIGEKDIWGRSMWI 717
30  Q: 370 EL*EWAHNVLIFVSHTRAHQRRARVAEEALNKHMDRMTGLMNACQPLSSAISILTKGP*-- 197
    ++ +WA +V IFVSH AHQ+ AEE N +D+MT +++ "Q" LS AI ++ +
    S: 718 DISKWAKDVKIFVSHVNAHQKVTSAEEEFNNQVDKMTSVDS-QTLSPAIPVIAQWAHEQ 776
    Q: 196 -----MNGLVVMA*NDGLSLSKADLATTTTEC 116
    +G A GL L+KADLAT +C
35  S: 777 SGHGGRDGGYPWAQQHGLPLTKADLATAAAD 808

```

The segment of gnl|PID|e314281 that is shown as "S" above is set out in the sequence listing as SEQ ID NO. 129.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out in the sequence listing as SEQ ID NO. 130 which corresponds to the "Q" sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: H. Meningima, M1; Human Heart.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:40 and may have been publicly available prior to conception of the present invention.

Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1747 of SEQ ID NO:40, b is an integer of 15 to 1761, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:40, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 31

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: H. Meningima, M1; Epithelial-TNF α and INF induced.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:41 and may have been publicly available prior to conception of the present invention.

Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 602 of SEQ ID NO:41, b is an integer of 15 to 616, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:41, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 32

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gi439877 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "reverse transcriptase [Homo sapiens]." A partial alignment demonstrating the observed homology is shown immediately below.

```
>gi439877 reverse transcriptase [Homo sapiens]
      Length = 361
```

Plus Strand HSPs:

```
Score = 237 (83.4 bits), Expect = 2.3e-19, P = 2.3e-19
Identities = 46/97 (47%), Positives = 62/97 (63%), Frame = +1
```

Q: 280 EKSNNHPPADPTSNTGSSYNST*DLVGTQIQTISVLNIIIIIREMQIKTTIRCHLTLVQMAFI 459
 +K+N+P + + S D+ + + IREMQIKTT+R HLT V+MA I
 S: 149 KKTNNPIKKWAKDMNRHFSKEDIYAAKHKMKCSPSLAIREMQIKTTMRVHLTPVRMAII 208
 Q: 460 QKTGNKNCWQECGEKGTLIHISWWSKSVQPLWKTWV 570
 +K+GNN+CW+ CGE GTL+H WW K VQPLWK+VW
 S: 209 KKSNNRCWRGCGEIGTLLHC-WWNCCLVQPLWKSVM 244

10 The segment of gil439877 that is shown as "S" above is set out in the sequence listing as SEQ ID NO. 131. Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein. Assays for determining such activities are also known in the art, some of which have been described elsewhere herein.

15 Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out in the sequence listing as SEQ ID NO. 132 which corresponds to the "Q" sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

 It has been discovered that this gene is expressed primarily in H. Meningima, M1.

20 Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 90 as residues: Phe-11 to Glu-18.

 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:42 and may have been publicly available prior to conception of the present invention.

25 Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 559 of SEQ ID NO:42, b is an integer of 15 to 573, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:42, and where b is greater than or
 30 equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 33

35 It has been discovered that this gene is expressed primarily in Spleen metastatic melanoma.

 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID

NO:43 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a
5 nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1370 of SEQ ID NO:43, b is an integer of 15 to 1384, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:43, and where b is greater than or equal to $a + 14$.

10 FEATURES OF PROTEIN ENCODED BY GENE NO: 34

It has been discovered that this gene is expressed primarily in Spleen metastatic melanoma.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID
15 NO:44 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between
20 1 to 692 of SEQ ID NO:44, b is an integer of 15 to 706, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:44, and where b is greater than or equal to $a + 14$.

FEATURES OF PROTEIN ENCODED BY GENE NO: 35

25 It has been discovered that this gene is expressed primarily in Spleen metastatic melanoma.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 93 as residues: Lys-39 to Lys-48.

Many polynucleotide sequences, such as EST sequences, are publicly available and
30 accessible through sequence databases. Some of these sequences are related to SEQ ID NO:45 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a

nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 405 of SEQ ID NO:45, b is an integer of 15 to 419, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:45, and where b is greater than or equal to a + 14.

5

FEATURES OF PROTEIN ENCODED BY GENE NO: 36

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Stratagene ovary (#937217); Spleen metastatic melanoma.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:46 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1011 of SEQ ID NO:46, b is an integer of 15 to 1025, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:46, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 37

It has been discovered that this gene is expressed primarily in Spleen metastatic melanoma.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:47 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 769 of SEQ ID NO:47, b is an integer of 15 to 783, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:47, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 38

It has been discovered that this gene is expressed primarily in Spleen metastatic melanoma.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:48 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 895 of SEQ ID NO:48, b is an integer of 15 to 909, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:48, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 39

It has been discovered that this gene is expressed primarily in Spleen metastatic melanoma.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:49 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 752 of SEQ ID NO:49, b is an integer of 15 to 766, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:49, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 40

It has been discovered that this gene is expressed primarily in Spleen metastatic melanoma.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 98 as residues: Arg-24 to Arg-35.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID

NO:50 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a
5 nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 698 of SEQ ID NO:50, b is an integer of 15 to 712, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:50, and where b is greater than or equal to a + 14.

10 FEATURES OF PROTEIN ENCODED BY GENE NO: 41

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: CD34 positive cells (cord blood).re-ex; Healing groin wound, 7.5 hours post incision; Synovial hypoxia-RSF subtracted; Spleen metastatic melanoma; CD34 depleted Buffy Coat (Cord Blood).

15 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:51 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly,
20 preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 666 of SEQ ID NO:51, b is an integer of 15 to 680, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:51, and where b is greater than or equal to a + 14.

25

FEATURES OF PROTEIN ENCODED BY GENE NO: 42

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Human Testes Tumor; Spleen metastatic melanoma.

Many polynucleotide sequences, such as EST sequences, are publicly available and
30 accessible through sequence databases. Some of these sequences are related to SEQ ID NO:52 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a

nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 990 of SEQ ID NO:52, b is an integer of 15 to 1004, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:52, and where b is greater than or equal to a + 14.

5

FEATURES OF PROTEIN ENCODED BY GENE NO: 43

It has been discovered that this gene is expressed primarily in Spleen metastatic melanoma.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 101 as residues: Pro-41 to Leu-47.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:53 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1291 of SEQ ID NO:53, b is an integer of 15 to 1305, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:53, and where b is greater than or equal to a + 14.

15
20

FEATURES OF PROTEIN ENCODED BY GENE NO: 44

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gi403460 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "transformation-related protein [Homo sapiens]." A partial alignment demonstrating the observed homology is shown immediately below.

25

>gi|403460 transformation-related protein [Homo sapiens] >sp|Q15662|Q15662
TRANSFORMATION-RELATED PROTEIN (FRAGMENT).
Length = 368

30

35

Plus Strand HSPs:

Score = 145 (51.0 bits), Expect = 6.2e-15, Sum P(2) = 6.2e-15
Identities = 30/59 (50%), Positives = 38/59 (64%), Frame = +3

Q: 123 WDYRHMPPRLRIFVFSVETGSHHAA*ADLELLS*TDPPASASQNTTRTTGVSHRAWPSLA 299
 WDYRH+PP ++ FVFSVETG H A A LELL+ + PP SA +AWP+L+
 S: 39 WDYRHVPPRQVHFVFSVETGFHRAGQAGLELLTSSVPPTSAFFPKCWDYRRDDQAWPTLS 97
 5 Score = 109 (38.4 bits), Expect = 6.2e-15, Sum P(2) = 6.2e-15
 Identities = 21/26 (80%), Positives = 24/26 (92%), Frame = +1
 Q: 10 DRISLLSPRLECNGVILANYNLRLPG 87
 DR+SLLSPRLECNG+ILA+ LRLPG
 10 S: 1 DRLSLLSPRLECNGMILAHCKLRLPG 26

The segments of gil403460 that are shown as "S" above are set out in the sequence listing as SEQ ID NO. 133 and SEQ ID NO. 135. Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein. Assays for determining such activities are also known in the art, some of which have been described elsewhere herein. Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out in the sequence listing as SEQ ID NO. 134 and/or SEQ ID NO. 136 which correspond to the "Q" sequences in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Spleen metastatic melanoma; Primary Dendritic Cells, lib 1.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:54 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 799 of SEQ ID NO:54, b is an integer of 15 to 813, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:54, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 45

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gil1710216 (all information available through the recited accession number is incorporated herein by reference) which is described

therein as "unknown [Homo sapiens]." A partial alignment demonstrating the observed homology is shown immediately below.

```

5      >gi|1710216 unknown [Homo sapiens]
          Length = 139

          Minus Strand HSPs:

10      Score = 111 (39.1 bits), Expect = 9.0e-09, Sum P(2) = 9.0e-09
          Identities = 23/34 (67%), Positives = 26/34 (76%), Frame = -1

          Q: 1595 AH*NLVLPGSSNPLTSASQVAGTTGTCHQTRLIF 1494
              AH NL LPGSSN SAS+VAGT GTC + +LIF
15      S: 71 AHCNLCLPGSSNSPASASRVAGTAGTCRRAQLIF 104

          Score = 57 (20.1 bits), Expect = 9.0e-09, Sum P(2) = 9.0e-09
          Identities = 11/18 (61%), Positives = 14/18 (77%), Frame = -3

20      Q: 1644 ETGSHSVAQVECSGASSS 1591
              ET SHSV ++ECSS S-
          S: 54 ETQSHSVTRLECSGTISA 71
  
```

The segments of gi1710216 that are shown as "S" above are set out in the sequence listing as SEQ ID NO. 137 and SEQ ID NO. 139.

25 Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out in the sequence listing as SEQ ID NO. 138 and/or SEQ ID NO. 140 which correspond to the "Q" sequences in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following

30 tissues/cDNA libraries: Spleen metastatic melanoma; Human Amygdala.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:55 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the

35 present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1680 of SEQ ID NO:55, b is an integer of 15 to 1694, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:55, and where b is greater than or

40 equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 46

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gi1809140 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "death domain containing protein CRADD [Homo sapiens]." A partial alignment demonstrating the observed homology is shown immediately below.

```

>gi|1809140 death domain containing protein CRADD [Homo sapiens]
>gi|1785557
    death adaptor molecule RAIDD [Homo sapiens]
    Length = 199

    Plus Strand HSPs:

Score = 525 (184.8 bits), Expect = 6.5e-50, P = 6.5e-50
Identities = 100/100 (100%), Positives = 100/100 (100%), Frame = +3

Q:   66 GDRLTGIPSHILNSSPSDRQINQLAQRIGPEWEPMVLSLGLSQTDIYRCKANHPPHNVSQ 245
      GDRLTGIPSHILNSSPSDRQINQLAQRIGPEWEPMVLSLGLSQTDIYRCKANHPPHNVSQ
S:  100 GDRLTGIPSHILNSSPSDRQINQLAQRIGPEWEPMVLSLGLSQTDIYRCKANHPPHNVSQ 199

Q:   246 VVEAFIRWRQRFQKQATFQSLHNGLRAVEVDPSSLHLMLE 365
      VVEAFIRWRQRFQKQATFQSLHNGLRAVEVDPSSLHLMLE
S:  160 VVEAFIRWRQRFQKQATFQSLHNGLRAVEVDPSSLHLMLE 199

```

The segment of gi1809140 that is shown as "S" above is set out in the sequence listing as SEQ ID NO. 141. Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein. Assays for determining such activities are also known in the art, some of which have been described elsewhere herein.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out in the sequence listing as SEQ ID NO. 142 which corresponds to the "Q" sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Human Striatum Depression, re-rescue; Human Macrophage; Aorta endothelial cells + TNF- α ; Soares_NhHMPu_S1; Human Activated T-Cells; 12 Week Old Early Stage Human; Soares melanocyte 2NbHM; Human Testes; Soares infant brain 1NIB.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:56 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the

present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 851 of SEQ ID NO:56, b is an integer of 15 to 865, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:56, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 47

It has been discovered that this gene is expressed primarily in Human Macrophage. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 105 as residues: Asn-19 to Thr-33.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:57 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 323 of SEQ ID NO:57, b is an integer of 15 to 337, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:57, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 48

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gil950650 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "immune-responsive gene 1 [Mus musculus]." This gene appears to be a human homolog of the murine gene. A partial alignment demonstrating the observed homology is shown immediately below.

```
>gi|950650 immune-responsive gene 1 [Mus musculus] >pir|I54546|I54546
immune-responsive gene 1 - mouse (fragment)
Length = 646
```

Plus Strand HSPs:

Score = 1629 (573.4 bits), Expect = 7.0e-167, P = 7.0e-167
 Identities = 325/451 (72%), Positives = 366/451 (81%), Frame = +2

```

5  Q:   11 SMDFDDTWHPATHPSGAVLPVLTALAEALPRSPKFSGLDLLLAFNVGIEVQGRLLHFAKE 190
    S:  182 SMDFDDTWHPATHPSGAVLPVLTALAEALPRSPKFSGLDLLLAFNVGIEVQGRLLHFAKE 241
10  Q:  191 ANDMPKRFPSPVVGTLGSAASAKFLGLSSTKCREALAIIVSHAGAPMANAATQTKPLH 370
    S:  242 AKDIPKRFPSPVVGTLGSAASAKFLGLSSTKCREALAIIVSHAGAPMANAATQTKPLH 301
15  Q:  371 IGNAAKHGIEAFLAMLGLQGNKQVLDLEAGFGAFYANYSPKVLPSIASYSWLLDQQDVA 550
    S:  302 IGNAAKHGMEATFLAMLGLQGNKQVLDLEAGFGAFYANYSPEDLPSLDLHIWLLDQQDVA 361
20  Q:  551 FKRFPAHLSTHWVADAAA+VRKHLV-PERALLPTDYIKRIVLRIPNVQYVNRFPVSEHE 727
    S:  362 FKRFPAHLSTHWVADAAA+VRKHLV-PERALLPTDYIKRIVLRIPNVQYVNRFPVSEHE 421
25  Q:  728 ARHSFQYVACAMLLDGGITVPSFHECQINRPQVRELLSKVELEYPPDNLPSPFNILYCEIS 907
    S:  422 ARHSFQYVACAMLLDGGITVPSFHECQINRPQVRELLSKVELEYPPDNLPSPFNILYCEIS 481
30  Q:  908 VTLKDGATFTDRSDTFYGHWRKPLSQEDLEEKFRANASKMLSWDTVESLIKIVKNLED*K 1087
    S:  482 VTLKDGATFTDRSDTFYGHWRKPLSQEDLEEKFRANASKMLSWDTVESLIKIVKNLED*K 541
35  Q: 1088 TVLC*LHFSKDLSTGRSFKLSSM*FYHKSLLRLTNI*MTLHLGRFNDLVCKARVCCCLVF 1267
    S:  542 TVLC*LHFSKDLSTGRSFKLSSM*FYHKSLLRLTNI*MTLHLGRFNDLVCKARVCCCLVF 1267
    Q: 1268 PGKMNKDGESPETELHISGRSLLLKILQDSST 1363
    S:  598 PGKMNKDGESPETELHISGRSLLLKILQDSST 1363
    Q: 1363 PGKMNKDGESPETELHISGRSLLLKILQDSST 1363
    S:  598 PGKMNKDGESPETELHISGRSLLLKILQDSST 1363
  
```

The segment of gil950650 that is shown as "S" above is set out in the sequence listing as SEQ ID NO. 143. Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art, some of which are described elsewhere herein. Assays for determining such activities are also known in the art, some of which have been described elsewhere herein.

Preferred polypeptides of the invention comprise a polypeptide having the amino acid sequence set out in the sequence listing as SEQ ID NO. 144 which corresponds to the "Q" sequence in the alignment shown above (gaps introduced in a sequence by the computer are, of course, removed).

The gene encoding the disclosed cDNA is believed to reside on chromosome 13. Accordingly, polynucleotides related to this invention are useful as a marker in linkage analysis for chromosome 13.

It has been discovered that this gene is expressed primarily in the following tissues/cDNA libraries: Activated T-Cells, 4 hrs, subtracted; Human Activated Monocytes.

The tissue distribution in immune cells and homology to immune-responsive gene I indicates the polynucleotides and polypeptides corresponding to this gene would be useful for the diagnosis and treatment of a variety of immune system disorders. Representative uses are described in the "Immune Activity" and "Infectious Disease" sections below, in Example 11, 13, 14, 16, 18, 19, 20, and 27, and elsewhere herein. Briefly, the expression indicates a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. Involvement in the regulation of cytokine production, antigen presentation, or other processes suggests a usefulness for treatment of cancer (e.g. by boosting immune responses). Expression in cells of lymphoid origin, indicates the natural gene product would be involved in immune functions. Therefore it would also be useful as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's disease, and scleroderma. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. Thus, this gene product is thought to be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Furthermore, the protein may also be used to determine biological activity, raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:58 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1763 of SEQ ID NO:58, b is an integer of 15 to 1777, where both a and b correspond to

the positions of nucleotide residues shown in SEQ ID NO:58, and where b is greater than or equal to $a + 14$.

Table 1

Gene No.	cDNA Clone ID	ATCC Deposit NO:Z and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
1	HMEIH57	203957 4/26/99	Lambda ZAP II	11	1626	1	1621	121	121	59	1	22	23	50
2	HMEIJ21	203957 4/26/99	Lambda ZAP II	12	2397	1	1525		553	60	1			20
3	HFEB A88	203957 4/26/99	Uni-ZAP XR	13	814	205	814	385	385	61	1	29	30	57
4	HMEIX79	203957 4/26/99	Lambda ZAP II	14	641	1	641	235	235	62	1			11
5	HMEJC96	203957 4/26/99	Lambda ZAP II	15	2163	401	2163	533	533	63	1			15
6	HMEJD36	203957 4/26/99	Lambda ZAP II	16	1312	1	1312	124	124	64	1	29	30	36
7	HMEKH55	203957 4/26/99	Lambda ZAP II	17	1726	1	1726	245	245	65	1	24	25	38
8	HMI:KW44	203957 4/26/99	Lambda ZAP II	18	2006	1	2006	26	26	66	1	33	34	58
9	HMELM75	203957 4/26/99	Lambda ZAP II	19	1607	1	1607	113	113	67	1	18	19	93
10	HMEW26	203957 4/26/99	Lambda ZAP II	20	1402	1	1339	92	92	68	1	18	19	23
11	HMGBT32	203957 4/26/99	Uni-ZAP XR	21	1221	1	1221	243	243	69	1			21

Gene No.	cDNA Clone ID	ATCC Deposit NO:Z and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
12	HMHBP74	203957 4/26/99	Uni-ZAP XR	22	1255	190	1255	208	208	70	1	22	23	33
13	HMIAC52	203957 4/26/99	Uni-ZAP XR	23	1642	58	1642	251	251	71	1	41	42	52
14	HMIAG55	203957 4/26/99	Uni-ZAP XR	24	932	1	932	113	113	72	1	23	24	35
15	HMIAK10	203957 4/26/99	Uni-ZAP XR	25	1064	1	1064	195	195	73	1	22	23	31
16	HIMIAL39	203957 4/26/99	Uni-ZAP XR	26	1280	1	1280	212	212	74	1	27	28	41
17	HMIAO82	203957 4/26/99	Uni-ZAP XR	27	3620	1	3620	129	129	75	1			24
18	HMIAR42	203957 4/26/99	Uni-ZAP XR	28	1896	1	1896	236	236	76	1			8
19	HMIAV33	203957 4/26/99	Uni-ZAP XR	29	1361	1	1361		360	77	1	17	18	24
20	HMIBD93	203957 4/26/99	Uni-ZAP XR	30	1323	734	1323		983	78	1	27	28	65
21	HMIBE95	203957 4/26/99	Uni-ZAP XR	31	2083	1	2083	31	31	79	1	24	25	90
22	HMIBF07	203957 4/26/99	Uni-ZAP XR	32	1738	1	1738	229	229	80	1			6
23	HMIBG57	203957 4/26/99	Uni-ZAP XR	33	917	1	917	130	130	81	1	23	24	33

Gene No.	cDNA Clone ID	A'CC Deposit NO:Z and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
24	HMIC180	203957 4/26/99	Uni-ZAP XR	34	1772	1	1772		1149	82	1	12	13	32
25	HMJAK70	203957 4/26/99	pSport1	35	799	1	799	273	273	83	1			10
26	HMKBA33	203957 4/26/99	pSport1	36	1730	1	1730	87	87	84	1	19	20	34
27	HMKCK32	203957 4/26/99	pSport1	37	1020	1	1020	60	60	85	1	20	21	37
28	HMKCP81	203957 4/26/99	pSport1	38	957	1	957	131	131	86	1	17	18	30
29	HMKCY49	203957 4/26/99	pSport1	39	1034	1	1034	269	269	87	1			2
30	HMKDG69	203957 4/26/99	pSport1	40	1761	1	1761		286	88	1			12
31	HMKDM80	203957 4/26/99	pSport1	41	616	1	616		386	89	1	14	15	35
32	HMKEG88	203957 4/26/99	pSport1	42	573	1	573	142	142	90	1			19
33	HMMAA09	203957 4/26/99	pSport1	43	1384	1	1384	127	127	91	1	30	31	34
34	HMMAK92	203957 4/26/99	pSport1	44	706	1	706	120	120	92	1	30	31	34
35	HMMAL32	203957 4/26/99	pSport1	45	419	1	419	276	276	93	1	35	36	48

Gene No.	cDNA Clone ID	ATCC Deposit NO:Z and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
36	HMMBD19	203957 4/26/99	pSportI	46	1025	1	1025	290	290	94	1			9
37	HMMBH91	203957 4/26/99	pSportI	47	783	1	783	234	234	95	1	18	19	22
38	HMMBH94	203957 4/26/99	pSportI	48	909	1	909	86	86	96	1			20
39	HMMBK55	203957 4/26/99	pSportI	49	766	1	766	260	260	97	1	20	21	41
40	HMMBQ31	203957 4/26/99	pSportI	50	712	1	712	85	85	98	1	21	22	37
41	HMMBR63	203957 4/26/99	pSportI	51	680	27	680	271	271	99	1	26	27	44
42	HMMBS55	203957 4/26/99	pSportI	52	1004	1	1004	103	103	100	1	22	23	40
43	HMMBT47	203957 4/26/99	pSportI	53	1305	1	1305	43	43	101	1	23	24	50
44	HMMCD35	203957 4/26/99	pSportI	54	813	1	813		308	102	1	23	24	42
45	HMMCD95	203957 4/26/99	pSportI	55	1694	1	1694	236	236	103	1			17
46	HMPAB26	203957 4/26/99	pBluescript	56	865	1	865		563	104	1			6
47	HMPAP48	203957 4/26/99	pBluescript	57	337	1	337	48	48	105	1	17	18	33

Gene No.	48	cDNA Clone ID	HMQAI38	ATCC Deposit NO:Z and Date	203957 4/26/99	Vector	NT SEQ ID NO: X	Total NT Seq.	1777	5' NT of Clone Seq.	1	3' NT of Clone Seq.	1777	5' NT of Start Codon	24	5' NT of First AA of Signal Pep	24	AA SEQ ID NO: Y	106	First AA of Sig Pep	1	Last AA of Sig Pep		First AA of Secreted Portion		Last AA of ORF	20
----------	----	---------------	---------	----------------------------	-------------------	--------	-----------------	---------------	------	---------------------	---	---------------------	------	----------------------	----	---------------------------------	----	-----------------	-----	---------------------	---	--------------------	--	------------------------------	--	----------------	----

Table 1 summarizes the information corresponding to each "Gene No." described above. The nucleotide sequence identified as "NT SEQ ID NO:X" was assembled from partially homologous ("overlapping") sequences obtained from the "cDNA clone ID" identified in Table 1 and, in some cases, from additional related DNA clones. The overlapping sequences were assembled into a single contiguous sequence of high redundancy (usually three to five overlapping sequences at each nucleotide position), resulting in a final sequence identified as SEQ ID NO:X.

The cDNA Clone ID was deposited on the date and given the corresponding deposit number listed in "ATCC Deposit No:Z and Date." Some of the deposits contain multiple different clones corresponding to the same gene. "Vector" refers to the type of vector contained in the cDNA Clone ID.

"Total NT Seq." refers to the total number of nucleotides in the contig identified by "Gene No." The deposited clone may contain all or most of these sequences, reflected by the nucleotide position indicated as "5' NT of Clone Seq." and the "3' NT of Clone Seq." of SEQ ID NO:X. The nucleotide position of SEQ ID NO:X of the putative start codon (methionine) is identified as "5' NT of Start Codon." Similarly, the nucleotide position of SEQ ID NO:X of the predicted signal sequence is identified as "5' NT of First AA of Signal Pep."

The translated amino acid sequence, beginning with the methionine, is identified as "AA SEQ ID NO:Y," although other reading frames can also be easily translated using known molecular biology techniques. The polypeptides produced by these alternative open reading frames are specifically contemplated by the present invention.

The first and last amino acid position of SEQ ID NO:Y of the predicted signal peptide is identified as "First AA of Sig Pep" and "Last AA of Sig Pep." The predicted first amino acid position of SEQ ID NO:Y of the secreted portion is identified as "Predicted First AA of Secreted Portion." Finally, the amino acid position of SEQ ID NO:Y of the last amino acid in the open reading frame is identified as "Last AA of ORF."

SEQ ID NO:X (where X may be any of the polynucleotide sequences disclosed in the sequence listing) and the translated SEQ ID NO:Y (where Y may be

any of the polypeptide sequences disclosed in the sequence listing) are sufficiently accurate and otherwise suitable for a variety of uses well known in the art and described further below. For instance, SEQ ID NO:X is useful for designing nucleic acid hybridization probes that will detect nucleic acid sequences contained in SEQ ID NO:X or the cDNA contained in the deposited clone. These probes will also hybridize to nucleic acid molecules in biological samples, thereby enabling a variety of forensic and diagnostic methods of the invention. Similarly, polypeptides identified from SEQ ID NO:Y may be used, for example, to generate antibodies which bind specifically to proteins containing the polypeptides and the secreted proteins encoded by the cDNA clones identified in Table 1.

Nevertheless, DNA sequences generated by sequencing reactions can contain sequencing errors. The errors exist as misidentified nucleotides, or as insertions or deletions of nucleotides in the generated DNA sequence. The erroneously inserted or deleted nucleotides cause frame shifts in the reading frames of the predicted amino acid sequence. In these cases, the predicted amino acid sequence diverges from the actual amino acid sequence, even though the generated DNA sequence may be greater than 99.9% identical to the actual DNA sequence (for example, one base insertion or deletion in an open reading frame of over 1000 bases).

Accordingly, for those applications requiring precision in the nucleotide sequence or the amino acid sequence, the present invention provides not only the generated nucleotide sequence identified as SEQ ID NO:X and the predicted translated amino acid sequence identified as SEQ ID NO:Y, but also a sample of plasmid DNA containing a human cDNA of the invention deposited with the ATCC, as set forth in Table 1. The nucleotide sequence of each deposited clone can readily be determined by sequencing the deposited clone in accordance with known methods. The predicted amino acid sequence can then be verified from such deposits. Moreover, the amino acid sequence of the protein encoded by a particular clone can also be directly determined by peptide sequencing or by expressing the protein in a suitable host cell containing the deposited human cDNA, collecting the protein, and determining its sequence.

The present invention also relates to the genes corresponding to SEQ ID NO:X, SEQ ID NO:Y, or the deposited clone. The corresponding gene can be isolated in accordance with known methods using the sequence information disclosed herein. Such methods include preparing probes or primers from the disclosed
5 sequence and identifying or amplifying the corresponding gene from appropriate sources of genomic material.

Also provided in the present invention are allelic variants, orthologs, and/or species homologs. Procedures known in the art can be used to obtain full-length genes, allelic variants, splice variants, full-length coding portions, orthologs, and/or
10 species homologs of genes corresponding to SEQ ID NO:X, SEQ ID NO:Y, or a deposited clone, using information from the sequences disclosed herein or the clones deposited with the ATCC. For example, allelic variants and/or species homologs may be isolated and identified by making suitable probes or primers from the sequences provided herein and screening a suitable nucleic acid source for allelic variants and/or
15 the desired homologue.

The polypeptides of the invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides
20 are well understood in the art.

The polypeptides may be in the form of the secreted protein, including the mature form, or may be a part of a larger protein, such as a fusion protein (see below). It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, pro-sequences, sequences which aid in purification,
25 such as multiple histidine residues, or an additional sequence for stability during recombinant production.

The polypeptides of the present invention are preferably provided in an isolated form, and preferably are substantially purified. A recombinantly produced version of a polypeptide, including the secreted polypeptide, can be substantially
30 purified using techniques described herein or otherwise known in the art, such as, for example, by the one-step method described in Smith and Johnson, Gene 67:31-40

(1988). Polypeptides of the invention also can be purified from natural, synthetic or recombinant sources using techniques described herein or otherwise known in the art, such as, for example, antibodies of the invention raised against the secreted protein.

The present invention provides a polynucleotide comprising, or alternatively
5 consisting of, the nucleic acid sequence of SEQ ID NO:X, and/or a cDNA contained in ATCC deposit Z. The present invention also provides a polypeptide comprising, or alternatively, consisting of, the polypeptide sequence of SEQ ID NO:Y and/or a polypeptide encoded by the cDNA contained in ATCC deposit Z. Polynucleotides encoding a polypeptide comprising, or alternatively consisting of the polypeptide
10 sequence of SEQ ID NO:Y and/or a polypeptide sequence encoded by the cDNA contained in ATCC deposit Z are also encompassed by the invention.

Signal Sequences

The present invention also encompasses mature forms of the polypeptide
15 having the polypeptide sequence of SEQ ID NO:Y and/or the polypeptide sequence encoded by the cDNA in a deposited clone. Polynucleotides encoding the mature forms (such as, for example, the polynucleotide sequence in SEQ ID NO:X and/or the polynucleotide sequence contained in the cDNA of a deposited clone) are also encompassed by the invention. According to the signal hypothesis, proteins secreted
20 by mammalian cells have a signal or secretory leader sequence which is cleaved from the mature protein once export of the growing protein chain across the rough endoplasmic reticulum has been initiated. Most mammalian cells and even insect cells cleave secreted proteins with the same specificity. However, in some cases, cleavage of a secreted protein is not entirely uniform, which results in two or more
25 mature species of the protein. Further, it has long been known that cleavage specificity of a secreted protein is ultimately determined by the primary structure of the complete protein, that is, it is inherent in the amino acid sequence of the polypeptide.

Methods for predicting whether a protein has a signal sequence, as well as the
30 cleavage point for that sequence, are available. For instance, the method of McGeoch, Virus Res. 3:271-286 (1985), uses the information from a short N-terminal

charged region and a subsequent uncharged region of the complete (uncleaved) protein. The method of von Heinje, *Nucleic Acids Res.* 14:4683-4690 (1986) uses the information from the residues surrounding the cleavage site, typically residues -13 to +2, where +1 indicates the amino terminus of the secreted protein. The accuracy of predicting the cleavage points of known mammalian secretory proteins for each of these methods is in the range of 75-80%. (von Heinje, *supra.*) However, the two methods do not always produce the same predicted cleavage point(s) for a given protein.

In the present case, the deduced amino acid sequence of the secreted polypeptide was analyzed by a computer program called SignalP (Henrik Nielsen et al., *Protein Engineering* 10:1-6 (1997)), which predicts the cellular location of a protein based on the amino acid sequence. As part of this computational prediction of localization, the methods of McGeoch and von Heinje are incorporated. The analysis of the amino acid sequences of the secreted proteins described herein by this program provided the results shown in Table 1.

As one of ordinary skill would appreciate, however, cleavage sites sometimes vary from organism to organism and cannot be predicted with absolute certainty. Accordingly, the present invention provides secreted polypeptides having a sequence shown in SEQ ID NO:Y which have an N-terminus beginning within 5 residues (i.e., + or - 5 residues) of the predicted cleavage point. Similarly, it is also recognized that in some cases, cleavage of the signal sequence from a secreted protein is not entirely uniform, resulting in more than one secreted species. These polypeptides, and the polynucleotides encoding such polypeptides, are contemplated by the present invention.

Moreover, the signal sequence identified by the above analysis may not necessarily predict the naturally occurring signal sequence. For example, the naturally occurring signal sequence may be further upstream from the predicted signal sequence. However, it is likely that the predicted signal sequence will be capable of directing the secreted protein to the ER. Nonetheless, the present invention provides the mature protein produced by expression of the polynucleotide sequence of SEQ ID NO:X and/or the polynucleotide sequence contained in the cDNA of a deposited

clone, in a mammalian cell (e.g., COS cells, as described below). These polypeptides, and the polynucleotides encoding such polypeptides, are contemplated by the present invention.

5 **Polynucleotide and Polypeptide Variants**

The present invention is directed to variants of the polynucleotide sequence disclosed in SEQ ID NO:X, the complementary strand thereto, and/or the cDNA sequence contained in a deposited clone.

10 The present invention also encompasses variants of the polypeptide sequence disclosed in SEQ ID NO:Y and/or encoded by a deposited clone.

"Variant" refers to a polynucleotide or polypeptide differing from the polynucleotide or polypeptide of the present invention, but retaining essential properties thereof. Generally, variants are overall closely similar, and, in many regions, identical to the polynucleotide or polypeptide of the present invention.

15 The present invention is also directed to nucleic acid molecules which comprise, or alternatively consist of, a nucleotide sequence which is at least 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to, for example, the nucleotide coding sequence in SEQ ID NO:X or the complementary strand thereto, the nucleotide coding sequence contained in a deposited cDNA clone or the
20 complementary strand thereto, a nucleotide sequence encoding the polypeptide of SEQ ID NO:Y, a nucleotide sequence encoding the polypeptide encoded by the cDNA contained in a deposited clone, and/or polynucleotide fragments of any of these nucleic acid molecules (e.g., those fragments described herein).
Polynucleotides which hybridize to these nucleic acid molecules under stringent
25 hybridization conditions or lower stringency conditions are also encompassed by the invention, as are polypeptides encoded by these polynucleotides.

The present invention is also directed to polypeptides which comprise, or alternatively consist of, an amino acid sequence which is at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% identical to, for example, the polypeptide sequence
30 shown in SEQ ID NO:Y, the polypeptide sequence encoded by the cDNA contained

in a deposited clone, and/or polypeptide fragments of any of these polypeptides (e.g., those fragments described herein).

By a nucleic acid having a nucleotide sequence at least, for example, 95% "identical" to a reference nucleotide sequence of the present invention, it is intended
5 that the nucleotide sequence of the nucleic acid is identical to the reference sequence except that the nucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence encoding the polypeptide. In other words, to obtain a nucleic acid having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the
10 reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. The query sequence may be an entire sequence shown in Table 1, the ORF (open reading frame), or any fragment specified as described herein.

15 As a practical matter, whether any particular nucleic acid molecule or polypeptide is at least 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to a nucleotide sequence of the present invention can be determined conventionally using known computer programs. A preferred method for determining the best overall match between a query sequence (a sequence of the present invention) and a
20 subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag et al. (Comp. App. Biosci. 6:237-245(1990)). In a sequence alignment the query and subject sequences are both DNA sequences. An RNA sequence can be compared by converting U's to T's. The result of said global sequence alignment is in percent
25 identity. Preferred parameters used in a FASTDB alignment of DNA sequences to calculate percent identity are: Matrix=Unitary, k-tuple=4, Mismatch Penalty=1, Joining Penalty=30, Randomization Group Length=0, Cutoff Score=1, Gap Penalty=5, Gap Size Penalty 0.05, Window Size=500 or the length of the subject nucleotide sequence, whichever is shorter.

30 If the subject sequence is shorter than the query sequence because of 5' or 3' deletions, not because of internal deletions, a manual correction must be made to the

results. This is because the FASTDB program does not account for 5' and 3' truncations of the subject sequence when calculating percent identity. For subject sequences truncated at the 5' or 3' ends, relative to the query sequence, the percent identity is corrected by calculating the number of bases of the query sequence that are 5' and 3' of the subject sequence, which are not matched/aligned, as a percent of the total bases of the query sequence. Whether a nucleotide is matched/aligned is determined by results of the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This corrected score is what is used for the purposes of the present invention. Only bases outside the 5' and 3' bases of the subject sequence, as displayed by the FASTDB alignment, which are not matched/aligned with the query sequence, are calculated for the purposes of manually adjusting the percent identity score.

For example, a 90 base subject sequence is aligned to a 100 base query sequence to determine percent identity. The deletions occur at the 5' end of the subject sequence and therefore, the FASTDB alignment does not show a matched/alignment of the first 10 bases at 5' end. The 10 unpaired bases represent 10% of the sequence (number of bases at the 5' and 3' ends not matched/total number of bases in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 bases were perfectly matched the final percent identity would be 90%. In another example, a 90 base subject sequence is compared with a 100 base query sequence. This time the deletions are internal deletions so that there are no bases on the 5' or 3' of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only bases 5' and 3' of the subject sequence which are not matched/aligned with the query sequence are manually corrected for. No other manual corrections are to be made for the purposes of the present invention.

By a polypeptide having an amino acid sequence at least, for example, 95% "identical" to a query amino acid sequence of the present invention, it is intended that the amino acid sequence of the subject polypeptide is identical to the query sequence

except that the subject polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the query amino acid sequence. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a query amino acid sequence, up to 5% of the amino acid residues in the subject
5 sequence may be inserted, deleted, (indels) or substituted with another amino acid. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

10 As a practical matter, whether any particular polypeptide is at least 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to, for instance, an amino acid sequences shown in Table 1 (SEQ ID NO:Y) or to the amino acid sequence encoded by cDNA contained in a deposited clone can be determined conventionally using known computer programs. A preferred method for determining the best overall match
15 between a query sequence (a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag et al. (Comp. App. Biosci. 6:237-245(1990)). In a sequence alignment the query and subject sequences are either both nucleotide sequences or both amino acid sequences. The result of said
20 global sequence alignment is in percent identity. Preferred parameters used in a FASTDB amino acid alignment are: Matrix=PAM 0, k-tuple=2, Mismatch Penalty=1, Joining Penalty=20, Randomization Group Length=0, Cutoff Score=1, Window Size=sequence length, Gap Penalty=5, Gap Size Penalty=0.05, Window Size=500 or the length of the subject amino acid sequence, whichever is shorter.

25 If the subject sequence is shorter than the query sequence due to N- or C-terminal deletions, not because of internal deletions, a manual correction must be made to the results. This is because the FASTDB program does not account for N- and C-terminal truncations of the subject sequence when calculating global percent identity. For subject sequences truncated at the N- and C-termini, relative to the
30 query sequence, the percent identity is corrected by calculating the number of residues of the query sequence that are N- and C-terminal of the subject sequence, which are

not matched/aligned with a corresponding subject residue, as a percent of the total bases of the query sequence. Whether a residue is matched/aligned is determined by results of the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This final percent identity score is what is used for the purposes of the present invention. Only residues to the N- and C-termini of the subject sequence, which are not matched/aligned with the query sequence, are considered for the purposes of manually adjusting the percent identity score. That is, only query residue positions outside the farthest N- and C-terminal residues of the subject sequence.

For example, a 90 amino acid residue subject sequence is aligned with a 100 residue query sequence to determine percent identity. The deletion occurs at the N-terminus of the subject sequence and therefore, the FASTDB alignment does not show a matching/alignment of the first 10 residues at the N-terminus. The 10 unpaired residues represent 10% of the sequence (number of residues at the N- and C-termini not matched/total number of residues in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 residues were perfectly matched the final percent identity would be 90%. In another example, a 90 residue subject sequence is compared with a 100 residue query sequence. This time the deletions are internal deletions so there are no residues at the N- or C-termini of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only residue positions outside the N- and C-terminal ends of the subject sequence, as displayed in the FASTDB alignment, which are not matched/aligned with the query sequence are manually corrected for. No other manual corrections are to be made for the purposes of the present invention.

The variants may contain alterations in the coding regions, non-coding regions, or both. Especially preferred are polynucleotide variants containing alterations which produce silent substitutions, additions, or deletions, but do not alter the properties or activities of the encoded polypeptide. Nucleotide variants produced by silent substitutions due to the degeneracy of the genetic code are preferred.

Moreover, variants in which 5-10, 1-5, or 1-2 amino acids are substituted, deleted, or added in any combination are also preferred. Polynucleotide variants can be produced for a variety of reasons, e.g., to optimize codon expression for a particular host (change codons in the human mRNA to those preferred by a bacterial host such as E. coli).

Naturally occurring variants are called "allelic variants," and refer to one of several alternate forms of a gene occupying a given locus on a chromosome of an organism. (Genes II, Lewin, B., ed., John Wiley & Sons, New York (1985).) These allelic variants can vary at either the polynucleotide and/or polypeptide level and are included in the present invention. Alternatively, non-naturally occurring variants may be produced by mutagenesis techniques or by direct synthesis.

Using known methods of protein engineering and recombinant DNA technology, variants may be generated to improve or alter the characteristics of the polypeptides of the present invention. For instance, one or more amino acids can be deleted from the N-terminus or C-terminus of the secreted protein without substantial loss of biological function. The authors of Ron et al., J. Biol. Chem. 268: 2984-2988 (1993), reported variant KGF proteins having heparin binding activity even after deleting 3, 8, or 27 amino-terminal amino acid residues. Similarly, Interferon gamma exhibited up to ten times higher activity after deleting 8-10 amino acid residues from the carboxy terminus of this protein. (Dobeli et al., J. Biotechnology 7:199-216 (1988).)

Moreover, ample evidence demonstrates that variants often retain a biological activity similar to that of the naturally occurring protein. For example, Gayle and coworkers (J. Biol. Chem 268:22105-22111 (1993)) conducted extensive mutational analysis of human cytokine IL-1a. They used random mutagenesis to generate over 3,500 individual IL-1a mutants that averaged 2.5 amino acid changes per variant over the entire length of the molecule. Multiple mutations were examined at every possible amino acid position. The investigators found that "[m]ost of the molecule could be altered with little effect on either [binding or biological activity]." (See, Abstract.) In fact, only 23 unique amino acid sequences, out of more than 3,500

nucleotide sequences examined, produced a protein that significantly differed in activity from wild-type.

Furthermore, even if deleting one or more amino acids from the N-terminus or C-terminus of a polypeptide results in modification or loss of one or more biological functions, other biological activities may still be retained. For example, the ability of a deletion variant to induce and/or to bind antibodies which recognize the secreted form will likely be retained when less than the majority of the residues of the secreted form are removed from the N-terminus or C-terminus. Whether a particular polypeptide lacking N- or C-terminal residues of a protein retains such immunogenic activities can readily be determined by routine methods described herein and otherwise known in the art.

Thus, the invention further includes polypeptide variants which show substantial biological activity. Such variants include deletions, insertions, inversions, repeats, and substitutions selected according to general rules known in the art so as to have little effect on activity. For example, guidance concerning how to make phenotypically silent amino acid substitutions is provided in Bowie et al., Science 247:1306-1310 (1990), wherein the authors indicate that there are two main strategies for studying the tolerance of an amino acid sequence to change.

The first strategy exploits the tolerance of amino acid substitutions by natural selection during the process of evolution. By comparing amino acid sequences in different species, conserved amino acids can be identified. These conserved amino acids are likely important for protein function. In contrast, the amino acid positions where substitutions have been tolerated by natural selection indicates that these positions are not critical for protein function. Thus, positions tolerating amino acid substitution could be modified while still maintaining biological activity of the protein.

The second strategy uses genetic engineering to introduce amino acid changes at specific positions of a cloned gene to identify regions critical for protein function. For example, site directed mutagenesis or alanine-scanning mutagenesis (introduction of single alanine mutations at every residue in the molecule) can be used.

(Cunningham and Wells, Science 244:1081-1085 (1989).) The resulting mutant molecules can then be tested for biological activity.

As the authors state, these two strategies have revealed that proteins are surprisingly tolerant of amino acid substitutions. The authors further indicate which amino acid changes are likely to be permissive at certain amino acid positions in the protein. For example, most buried (within the tertiary structure of the protein) amino acid residues require nonpolar side chains, whereas few features of surface side chains are generally conserved. Moreover, tolerated conservative amino acid substitutions involve replacement of the aliphatic or hydrophobic amino acids Ala, Val, Leu and Ile; replacement of the hydroxyl residues Ser and Thr; replacement of the acidic residues Asp and Glu; replacement of the amide residues Asn and Gln, replacement of the basic residues Lys, Arg, and His; replacement of the aromatic residues Phe, Tyr, and Trp, and replacement of the small-sized amino acids Ala, Ser, Thr, Met, and Gly.

Besides conservative amino acid substitution, variants of the present invention include (i) substitutions with one or more of the non-conserved amino acid residues, where the substituted amino acid residues may or may not be one encoded by the genetic code, or (ii) substitution with one or more of amino acid residues having a substituent group, or (iii) fusion of the mature polypeptide with another compound, such as a compound to increase the stability and/or solubility of the polypeptide (for example, polyethylene glycol), or (iv) fusion of the polypeptide with additional amino acids, such as, for example, an IgG Fc fusion region peptide, or leader or secretory sequence, or a sequence facilitating purification. Such variant polypeptides are deemed to be within the scope of those skilled in the art from the teachings herein.

For example, polypeptide variants containing amino acid substitutions of charged amino acids with other charged or neutral amino acids may produce proteins with improved characteristics, such as less aggregation. Aggregation of pharmaceutical formulations both reduces activity and increases clearance due to the aggregate's immunogenic activity. (Pinckard et al., Clin. Exp. Immunol. 2:331-340 (1967); Robbins et al., Diabetes 36: 838-845 (1987); Cleland et al., Crit. Rev. Therapeutic Drug Carrier Systems 10:307-377 (1993).)

A further embodiment of the invention relates to a polypeptide which

comprises the amino acid sequence of the present invention having an amino acid sequence which contains at least one amino acid substitution, but not more than 50 amino acid substitutions, even more preferably, not more than 40 amino acid substitutions, still more preferably, not more than 30 amino acid substitutions, and still even more preferably, not more than 20 amino acid substitutions. Of course, in order of ever-increasing preference, it is highly preferable for a peptide or polypeptide to have an amino acid sequence which comprises the amino acid sequence of the present invention, which contains at least one, but not more than 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1 amino acid substitutions. In specific embodiments, the number of additions, substitutions, and/or deletions in the amino acid sequence of the present invention or fragments thereof (e.g., the mature form and/or other fragments described herein), is 1-5, 5-10, 5-25, 5-50, 10-50 or 50-150, conservative amino acid substitutions are preferable.

15 **Polynucleotide and Polypeptide Fragments**

The present invention is also directed to polynucleotide fragments of the polynucleotides of the invention.

In the present invention, a "polynucleotide fragment" refers to a short polynucleotide having a nucleic acid sequence which: is a portion of that contained in a deposited clone, or encoding the polypeptide encoded by the cDNA in a deposited clone; is a portion of that shown in SEQ ID NO:X or the complementary strand thereto, or is a portion of a polynucleotide sequence encoding the polypeptide of SEQ ID NO:Y. The nucleotide fragments of the invention are preferably at least about 15 nt, and more preferably at least about 20 nt, still more preferably at least about 30 nt, and even more preferably, at least about 40 nt, at least about 50 nt, at least about 75 nt, or at least about 150 nt in length. A fragment "at least 20 nt in length," for example, is intended to include 20 or more contiguous bases from the cDNA sequence contained in a deposited clone or the nucleotide sequence shown in SEQ ID NO:X. In this context "about" includes the particularly recited value, a value larger or smaller by several (5, 4, 3, 2, or 1) nucleotides, at either terminus or at both termini. These nucleotide fragments have uses that include, but are not limited to, as

diagnostic probes and primers as discussed herein. Of course, larger fragments (e.g., 50, 150, 500, 600, 2000 nucleotides) are preferred.

Moreover, representative examples of polynucleotide fragments of the invention, include, for example, fragments comprising, or alternatively consisting of,
5 a sequence from about nucleotide number 1-50, 51-100, 101-150, 151-200, 201-250, 251-300, 301-350, 351-400, 401-450, 451-500, 501-550, 551-600, 651-700, 701-750, 751-800, 800-850, 851-900, 901-950, 951-1000, 1001-1050, 1051-1100, 1101-1150, 1151-1200, 1201-1250, 1251-1300, 1301-1350, 1351-1400, 1401-1450, 1451-1500, 1501-1550, 1551-1600, 1601-1650, 1651-1700, 1701-1750, 1751-1800, 1801-1850,
10 1851-1900, 1901-1950, 1951-2000, or 2001 to the end of SEQ ID NO:X, or the complementary strand thereto, or the cDNA contained in a deposited clone. In this context "about" includes the particularly recited ranges, and ranges larger or smaller by several (5, 4, 3, 2, or 1) nucleotides, at either terminus or at both termini. Preferably, these fragments encode a polypeptide which has biological activity. More
15 preferably, these polynucleotides can be used as probes or primers as discussed herein. Polynucleotides which hybridize to these nucleic acid molecules under stringent hybridization conditions or lower stringency conditions are also encompassed by the invention, as are polypeptides encoded by these polynucleotides.

In the present invention, a "polypeptide fragment" refers to an amino acid
20 sequence which is a portion of that contained in SEQ ID NO:Y or encoded by the cDNA contained in a deposited clone. Protein (polypeptide) fragments may be "free-standing," or comprised within a larger polypeptide of which the fragment forms a part or region, most preferably as a single continuous region. Representative examples of polypeptide fragments of the invention, include, for example, fragments
25 comprising, or alternatively consisting of, from about amino acid number 1-20, 21-40, 41-60, 61-80, 81-100, 102-120, 121-140, 141-160, or 161 to the end of the coding region. Moreover, polypeptide fragments can be about 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, or 150 amino acids in length. In this context "about" includes the particularly recited ranges or values, and ranges or values larger or
30 smaller by several (5, 4, 3, 2, or 1) amino acids, at either extreme or at both extremes. Polynucleotides encoding these polypeptides are also encompassed by the invention.

Preferred polypeptide fragments include the secreted protein as well as the mature form. Further preferred polypeptide fragments include the secreted protein or the mature form having a continuous series of deleted residues from the amino or the carboxy terminus, or both. For example, any number of amino acids, ranging from 1-60, can be deleted from the amino terminus of either the secreted polypeptide or the mature form. Similarly, any number of amino acids, ranging from 1-30, can be deleted from the carboxy terminus of the secreted protein or mature form. Furthermore, any combination of the above amino and carboxy terminus deletions are preferred. Similarly, polynucleotides encoding these polypeptide fragments are also preferred.

Also preferred are polypeptide and polynucleotide fragments characterized by structural or functional domains, such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions. Polypeptide fragments of SEQ ID NO:Y falling within conserved domains are specifically contemplated by the present invention. Moreover, polynucleotides encoding these domains are also contemplated.

Other preferred polypeptide fragments are biologically active fragments. Biologically active fragments are those exhibiting activity similar, but not necessarily identical, to an activity of the polypeptide of the present invention. The biological activity of the fragments may include an improved desired activity, or a decreased undesirable activity. Polynucleotides encoding these polypeptide fragments are also encompassed by the invention.

Preferably, the polynucleotide fragments of the invention encode a polypeptide which demonstrates a functional activity. By a polypeptide demonstrating a "functional activity" is meant, a polypeptide capable of displaying one or more known functional activities associated with a full-length (complete) polypeptide of invention protein. Such functional activities include, but are not limited to, biological activity, antigenicity [ability to bind (or compete with a

polypeptide of the invention for binding) to an antibody to the polypeptide of the invention], immunogenicity (ability to generate antibody which binds to a polypeptide of the invention), ability to form multimers with polypeptides of the invention, and ability to bind to a receptor or ligand for a polypeptide of the invention.

5 The functional activity of polypeptides of the invention, and fragments, variants derivatives, and analogs thereof, can be assayed by various methods.

For example, in one embodiment where one is assaying for the ability to bind or compete with full-length polypeptide of the invention for binding to an antibody of the polypeptide of the invention, various immunoassays known in the art can be used, including but not limited to, competitive and non-competitive assay systems using techniques such as radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoradiometric assays, gel diffusion precipitation reactions, immunodiffusion assays, in situ immunoassays (using colloidal gold, enzyme or radioisotope labels, for example), western blots, precipitation reactions, agglutination assays (e.g., gel agglutination assays, hemagglutination assays), complement fixation assays, immunofluorescence assays, protein A assays, and immunoelectrophoresis assays, etc. In one embodiment, antibody binding is detected by detecting a label on the primary antibody. In another embodiment, the primary antibody is detected by detecting binding of a secondary antibody or reagent to the primary antibody. In a further embodiment, the secondary antibody is labeled. Many means are known in the art for detecting binding in an immunoassay and are within the scope of the present invention.

In another embodiment, where a ligand for a polypeptide of the invention is identified, or the ability of a polypeptide fragment, variant or derivative of the invention to multimerize is being evaluated, binding can be assayed, e.g., by means well-known in the art, such as, for example, reducing and non-reducing gel chromatography, protein affinity chromatography, and affinity blotting. See generally, Phizicky, E., et al., 1995, Microbiol. Rev. 59:94-123. In another embodiment, physiological correlates of binding of a polypeptide of the invention to its substrates (signal transduction) can be assayed.

In addition, assays described herein (see Examples) and otherwise known in the art may routinely be applied to measure the ability of polypeptides of the invention and fragments, variants derivatives and analogs thereof to elicit related biological activity related to that of the polypeptide of the invention (either in vitro or in vivo). Other methods will be known to the skilled artisan and are within the scope of the invention.

Epitopes and Antibodies

The present invention encompasses polypeptides comprising, or alternatively consisting of, an epitope of the polypeptide having an amino acid sequence of SEQ ID NO:Y, or an epitope of the polypeptide sequence encoded by a polynucleotide sequence contained in ATCC deposit No. Z or encoded by a polynucleotide that hybridizes to the complement of the sequence of SEQ ID NO:X or contained in ATCC deposit No. Z under stringent hybridization conditions or lower stringency hybridization conditions as defined supra. The present invention further encompasses polynucleotide sequences encoding an epitope of a polypeptide sequence of the invention (such as, for example, the sequence disclosed in SEQ ID NO:X), polynucleotide sequences of the complementary strand of a polynucleotide sequence encoding an epitope of the invention, and polynucleotide sequences which hybridize to the complementary strand under stringent hybridization conditions or lower stringency hybridization conditions defined supra.

The term "epitopes," as used herein, refers to portions of a polypeptide having antigenic or immunogenic activity in an animal, preferably a mammal, and most preferably in a human. In a preferred embodiment, the present invention encompasses a polypeptide comprising an epitope, as well as the polynucleotide encoding this polypeptide. An "immunogenic epitope," as used herein, is defined as a portion of a protein that elicits an antibody response in an animal, as determined by any method known in the art, for example, by the methods for generating antibodies described infra. (See, for example, Geysen et al., Proc. Natl. Acad. Sci. USA 81:3998-4002 (1983)). The term "antigenic epitope," as used herein, is defined as a portion of a protein to which an antibody can immunospecifically bind its antigen as

determined by any method well known in the art, for example, by the immunoassays described herein. Immunospecific binding excludes non-specific binding but does not necessarily exclude cross-reactivity with other antigens. Antigenic epitopes need not necessarily be immunogenic.

5 Fragments which function as epitopes may be produced by any conventional means. (See, e.g., Houghten, Proc. Natl. Acad. Sci. USA 82:5131-5135 (1985), further described in U.S. Patent No. 4,631,211).

10 In the present invention, antigenic epitopes preferably contain a sequence of at least 4, at least 5, at least 6, at least 7, more preferably at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 20, at least 25, at least 30, at least 40, at least 50, and, most preferably, between about 15 to about 30 amino acids. Preferred polypeptides comprising immunogenic or antigenic epitopes are at least 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 amino acid residues in length. Additional non-exclusive preferred antigenic epitopes
15 include the antigenic epitopes disclosed herein, as well as portions thereof. Antigenic epitopes are useful, for example, to raise antibodies, including monoclonal antibodies, that specifically bind the epitope. Preferred antigenic epitopes include the antigenic epitopes disclosed herein, as well as any combination of two, three, four, five or more of these antigenic epitopes. Antigenic epitopes can be used as the target molecules in
20 immunoassays. (See, for instance, Wilson et al., Cell 37:767-778 (1984); Sutcliffe et al., Science 219:660-666 (1983)).

25 Similarly, immunogenic epitopes can be used, for example, to induce antibodies according to methods well known in the art. (See, for instance, Sutcliffe et al., supra; Wilson et al., supra; Chow et al., Proc. Natl. Acad. Sci. USA 82:910-914; and Bittle et al., J. Gen. Virol. 66:2347-2354 (1985)). Preferred immunogenic epitopes include the immunogenic epitopes disclosed herein, as well as any combination of two, three, four, five or more of these immunogenic epitopes. The polypeptides comprising one or more immunogenic epitopes may be presented for eliciting an antibody response together with a carrier protein, such as an albumin, to
30 an animal system (such as rabbit or mouse), or, if the polypeptide is of sufficient length (at least about 25 amino acids), the polypeptide may be presented without a

carrier. However, immunogenic epitopes comprising as few as 8 to 10 amino acids have been shown to be sufficient to raise antibodies capable of binding to, at the very least, linear epitopes in a denatured polypeptide (e.g., in Western blotting).

Epitope-bearing polypeptides of the present invention may be used to induce
5 antibodies according to methods well known in the art including, but not limited to, in vivo immunization, in vitro immunization, and phage display methods. See, e.g., Sutcliffe et al., supra; Wilson et al., supra, and Bittle et al., J. Gen. Virol., 66:2347-2354 (1985). If in vivo immunization is used, animals may be immunized with free peptide; however, anti-peptide antibody titer may be boosted by coupling the peptide
10 to a macromolecular carrier, such as keyhole limpet hemacyanin (KLH) or tetanus toxoid. For instance, peptides containing cysteine residues may be coupled to a carrier using a linker such as maleimidobenzoyl- N-hydroxysuccinimide ester (MBS), while other peptides may be coupled to carriers using a more general linking agent such as glutaraldehyde. Animals such as rabbits, rats and mice are immunized with
15 either free or carrier- coupled peptides, for instance, by intraperitoneal and/or intradermal injection of emulsions containing about 100 µg of peptide or carrier protein and Freund's adjuvant or any other adjuvant known for stimulating an immune response. Several booster injections may be needed, for instance, at intervals of about two weeks, to provide a useful titer of anti-peptide antibody which
20 can be detected, for example, by ELISA assay using free peptide adsorbed to a solid surface. The titer of anti-peptide antibodies in serum from an immunized animal may be increased by selection of anti-peptide antibodies, for instance, by adsorption to the peptide on a solid support and elution of the selected antibodies according to methods well known in the art.

25 As one of skill in the art will appreciate, and as discussed above, the polypeptides of the present invention comprising an immunogenic or antigenic epitope can be fused to other polypeptide sequences. For example, the polypeptides of the present invention may be fused with the constant domain of immunoglobulins (IgA, IgE, IgG, IgM), or portions thereof (CH1, CH2, CH3, or any combination
30 thereof and portions thereof) resulting in chimeric polypeptides. Such fusion proteins may facilitate purification and may increase half-life in vivo. This has been shown

for chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins. See, e.g., EP 394,827; Traunecker et al., *Nature*, 331:84-86 (1988). Enhanced delivery of an antigen across the epithelial barrier to the immune system has been demonstrated for antigens (e.g., insulin) conjugated to an FcRn binding partner such as IgG or Fc fragments (see, e.g., PCT Publications WO 96/22024 and WO 99/04813). IgG Fusion proteins that have a disulfide-linked dimeric structure due to the IgG portion disulfide bonds have also been found to be more efficient in binding and neutralizing other molecules than monomeric polypeptides or fragments thereof alone. See, e.g., Fountoulakis et al., *J. Biochem.*, 270:3958-3964 (1995). Nucleic acids encoding the above epitopes can also be recombined with a gene of interest as an epitope tag (e.g., the hemagglutinin ("HA") tag or flag tag) to aid in detection and purification of the expressed polypeptide. For example, a system described by Janknecht et al. allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht et al., 1991, *Proc. Natl. Acad. Sci. USA* 88:8972-897). In this system, the gene of interest is subcloned into a vaccinia recombination plasmid such that the open reading frame of the gene is translationally fused to an amino-terminal tag consisting of six histidine residues. The tag serves as a matrix binding domain for the fusion protein. Extracts from cells infected with the recombinant vaccinia virus are loaded onto Ni²⁺-nitriloacetic acid-agarose column and histidine-tagged proteins can be selectively eluted with imidazole-containing buffers.

Additional fusion proteins of the invention may be generated through the techniques of gene-shuffling, motif-shuffling, exon-shuffling, and/or codon-shuffling (collectively referred to as "DNA shuffling"). DNA shuffling may be employed to modulate the activities of polypeptides of the invention, such methods can be used to generate polypeptides with altered activity, as well as agonists and antagonists of the polypeptides. See, generally, U.S. Patent Nos. 5,605,793; 5,811,238; 5,830,721; 5,834,252; and 5,837,458, and Patten et al., *Curr. Opin. Biotechnol.* 8:724-33 (1997); Harayama, *Trends Biotechnol.* 16(2):76-82 (1998); Hansson, et al., *J. Mol. Biol.* 287:265-76 (1999); and Lorenzo and Blasco, *Biotechniques* 24(2):308-13

(1998) (each of these patents and publications are hereby incorporated by reference in its entirety). In one embodiment, alteration of polynucleotides corresponding to SEQ ID NO:X and the polypeptides encoded by these polynucleotides may be achieved by DNA shuffling. DNA shuffling involves the assembly of two or more DNA
5 segments by homologous or site-specific recombination to generate variation in the polynucleotide sequence. In another embodiment, polynucleotides of the invention, or the encoded polypeptides, may be altered by being subjected to random mutagenesis by error-prone PCR, random nucleotide insertion or other methods prior to recombination. In another embodiment, one or more components, motifs, sections,
10 parts, domains, fragments, etc., of a polynucleotide encoding a polypeptide of the invention may be recombined with one or more components, motifs, sections, parts, domains, fragments, etc. of one or more heterologous molecules.

Antibodies

15 Further polypeptides of the invention relate to antibodies and T-cell antigen receptors (TCR) which immunospecifically bind a polypeptide, polypeptide fragment, or variant of SEQ ID NO:Y, and/or an epitope, of the present invention (as determined by immunoassays well known in the art for assaying specific antibody-antigen binding). Antibodies of the invention include, but are not limited to,
20 polyclonal, monoclonal, multispecific, human, humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab') fragments, fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies (including, e.g., anti-Id antibodies to antibodies of the invention), and epitope-binding fragments of any of the above. The term "antibody," as used herein, refers to immunoglobulin molecules
25 and immunologically active portions of immunoglobulin molecules, i.e., molecules that contain an antigen binding site that immunospecifically binds an antigen. The immunoglobulin molecules of the invention can be of any type (e.g., IgG, IgE, IgM, IgD, IgA and IgY), class (e.g., IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2) or subclass of immunoglobulin molecule.

30 Most preferably the antibodies are human antigen-binding antibody fragments of the present invention and include, but are not limited to, Fab, Fab' and F(ab')₂, Fd,

single-chain Fvs (scFv), single-chain antibodies, disulfide-linked Fvs (sdFv) and fragments comprising either a VL or VH domain. Antigen-binding antibody fragments, including single-chain antibodies, may comprise the variable region(s) alone or in combination with the entirety or a portion of the following: hinge region, CH1, CH2, and CH3 domains. Also included in the invention are antigen-binding fragments also comprising any combination of variable region(s) with a hinge region, CH1, CH2, and CH3 domains. The antibodies of the invention may be from any animal origin including birds and mammals. Preferably, the antibodies are human, murine (e.g., mouse and rat), donkey, sheep rabbit, goat, guinea pig, camel, horse, or chicken. As used herein, "human" antibodies include antibodies having the amino acid sequence of a human immunoglobulin and include antibodies isolated from human immunoglobulin libraries or from animals transgenic for one or more human immunoglobulin and that do not express endogenous immunoglobulins, as described infra and, for example in, U.S. Patent No. 5,939,598 by Kucherlapati et al.

The antibodies of the present invention may be monospecific, bispecific, trispecific or of greater multispecificity. Multispecific antibodies may be specific for different epitopes of a polypeptide of the present invention or may be specific for both a polypeptide of the present invention as well as for a heterologous epitope, such as a heterologous polypeptide or solid support material. See, e.g., PCT publications WO 93/17715; WO 92/08802; WO 91/00360; WO 92/05793; Tutt, et al., J. Immunol. 147:60-69 (1991); U.S. Patent Nos. 4,474,893; 4,714,681; 4,925,648; 5,573,920; 5,601,819; Kostelny et al., J. Immunol. 148:1547-1553 (1992).

Antibodies of the present invention may be described or specified in terms of the epitope(s) or portion(s) of a polypeptide of the present invention which they recognize or specifically bind. The epitope(s) or polypeptide portion(s) may be specified as described herein, e.g., by N-terminal and C-terminal positions, by size in contiguous amino acid residues, or listed in the Tables and Figures. Antibodies which specifically bind any epitope or polypeptide of the present invention may also be excluded. Therefore, the present invention includes antibodies that specifically bind polypeptides of the present invention, and allows for the exclusion of the same.

Antibodies of the present invention may also be described or specified in terms of their cross-reactivity. Antibodies that do not bind any other analog, ortholog, or homolog of a polypeptide of the present invention are included. Antibodies that bind polypeptides with at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 65%, at least 60%, at least 55%, and at least 50% identity (as calculated using methods known in the art and described herein) to a polypeptide of the present invention are also included in the present invention. In specific embodiments, antibodies of the present invention cross-react with murine, rat and/or rabbit homologs of human proteins and the corresponding epitopes thereof. Antibodies that do not bind polypeptides with less than 95%, less than 90%, less than 85%, less than 80%, less than 75%, less than 70%, less than 65%, less than 60%, less than 55%, and less than 50% identity (as calculated using methods known in the art and described herein) to a polypeptide of the present invention are also included in the present invention. In a specific embodiment, the above-described cross-reactivity is with respect to any single specific antigenic or immunogenic polypeptide, or combination(s) of 2, 3, 4, 5, or more of the specific antigenic and/or immunogenic polypeptides disclosed herein. Further included in the present invention are antibodies which bind polypeptides encoded by polynucleotides which hybridize to a polynucleotide of the present invention under stringent hybridization conditions (as described herein). Antibodies of the present invention may also be described or specified in terms of their binding affinity to a polypeptide of the invention. Preferred binding affinities include those with a dissociation constant or K_d less than 5×10^{-2} M, 10^{-2} M, 5×10^{-3} M, 10^{-3} M, 5×10^{-4} M, 10^{-4} M, 5×10^{-5} M, 10^{-5} M, 5×10^{-6} M, 10^{-6} M, 5×10^{-7} M, 10^{-7} M, 5×10^{-8} M, 10^{-8} M, 5×10^{-9} M, 10^{-9} M, 5×10^{-10} M, 10^{-10} M, 5×10^{-11} M, 10^{-11} M, 5×10^{-12} M, 10^{-12} M, 5×10^{-13} M, 10^{-13} M, 5×10^{-14} M, 10^{-14} M, 5×10^{-15} M, or 10^{-15} M.

The invention also provides antibodies that competitively inhibit binding of an antibody to an epitope of the invention as determined by any method known in the art for determining competitive binding, for example, the immunoassays described herein. In preferred embodiments, the antibody competitively inhibits binding to the

epitope by at least 95%, at least 90%, at least 85 %, at least 80%, at least 75%, at least 70%, at least 60%, or at least 50%.

Antibodies of the present invention may act as agonists or antagonists of the polypeptides of the present invention. For example, the present invention includes
5 antibodies which disrupt the receptor/ligand interactions with the polypeptides of the invention either partially or fully. Preferably, antibodies of the present invention bind an antigenic epitope disclosed herein, or a portion thereof. The invention features both receptor-specific antibodies and ligand-specific antibodies. The invention also features receptor-specific antibodies which do not prevent ligand
10 binding but prevent receptor activation. Receptor activation (i.e., signaling) may be determined by techniques described herein or otherwise known in the art. For example, receptor activation can be determined by detecting the phosphorylation (e.g., tyrosine or serine/threonine) of the receptor or its substrate by immunoprecipitation followed by western blot analysis (for example, as described
15 supra). In specific embodiments, antibodies are provided that inhibit ligand activity or receptor activity by at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, or at least 50% of the activity in absence of the antibody.

The invention also features receptor-specific antibodies which both prevent
20 ligand binding and receptor activation as well as antibodies that recognize the receptor-ligand complex, and, preferably, do not specifically recognize the unbound receptor or the unbound ligand. Likewise, included in the invention are neutralizing antibodies which bind the ligand and prevent binding of the ligand to the receptor, as well as antibodies which bind the ligand, thereby preventing receptor activation, but
25 do not prevent the ligand from binding the receptor. Further included in the invention are antibodies which activate the receptor. These antibodies may act as receptor agonists, i.e., potentiate or activate either all or a subset of the biological activities of the ligand-mediated receptor activation, for example, by inducing dimerization of the receptor. The antibodies may be specified as agonists, antagonists or inverse agonists
30 for biological activities comprising the specific biological activities of the peptides of the invention disclosed herein. The above antibody agonists can be made using

methods known in the art. See, e.g., PCT publication WO 96/40281; U.S. Patent No. 5,811,097; Deng et al., *Blood* 92(6):1981-1988 (1998); Chen et al., *Cancer Res.* 58(16):3668-3678 (1998); Harrop et al., *J. Immunol.* 161(4):1786-1794 (1998); Zhu et al., *Cancer Res.* 58(15):3209-3214 (1998); Yoon et al., *J. Immunol.* 160(7):3170-3179 (1998); Prat et al., *J. Cell. Sci.* 111(Pt2):237-247 (1998); Pitard et al., *J. Immunol. Methods* 205(2):177-190 (1997); Liautard et al., *Cytokine* 9(4):233-241 (1997); Carlson et al., *J. Biol. Chem.* 272(17):11295-11301 (1997); Taryman et al., *Neuron* 14(4):755-762 (1995); Muller et al., *Structure* 6(9):1153-1167 (1998); Bartunek et al., *Cytokine* 8(1):14-20 (1996) (which are all incorporated by reference herein in their entireties).

Antibodies of the present invention may be used, for example, but not limited to, to purify, detect, and target the polypeptides of the present invention, including both in vitro and in vivo diagnostic and therapeutic methods. For example, the antibodies have use in immunoassays for qualitatively and quantitatively measuring levels of the polypeptides of the present invention in biological samples. See, e.g., Harlow et al., *Antibodies: A Laboratory Manual*, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988) (incorporated by reference herein in its entirety).

As discussed in more detail below, the antibodies of the present invention may be used either alone or in combination with other compositions. The antibodies may further be recombinantly fused to a heterologous polypeptide at the N- or C-terminus or chemically conjugated (including covalently and non-covalently conjugations) to polypeptides or other compositions. For example, antibodies of the present invention may be recombinantly fused or conjugated to molecules useful as labels in detection assays and effector molecules such as heterologous polypeptides, drugs, radionuclides, or toxins. See, e.g., PCT publications WO 92/08495; WO 91/14438; WO 89/12624; U.S. Patent No. 5,314,995; and EP 396.387.

The antibodies of the invention include derivatives that are modified, i.e., by the covalent attachment of any type of molecule to the antibody such that covalent attachment does not prevent the antibody from generating an anti-idiotypic response. For example, but not by way of limitation, the antibody derivatives include antibodies that have been modified, e.g., by glycosylation, acetylation, pegylation,

phosphylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to a cellular ligand or other protein, etc. Any of numerous chemical modifications may be carried out by known techniques, including, but not limited to specific chemical cleavage, acetylation, formylation, metabolic synthesis of tunicamycin, etc. Additionally, the derivative may contain one or more non-classical amino acids.

The antibodies of the present invention may be generated by any suitable method known in the art. Polyclonal antibodies to an antigen-of-interest can be produced by various procedures well known in the art. For example, a polypeptide of the invention can be administered to various host animals including, but not limited to, rabbits, mice, rats, etc. to induce the production of sera containing polyclonal antibodies specific for the antigen. Various adjuvants may be used to increase the immunological response, depending on the host species, and include but are not limited to, Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanins, dinitrophenol, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and corynebacterium parvum. Such adjuvants are also well known in the art.

Monoclonal antibodies can be prepared using a wide variety of techniques known in the art including the use of hybridoma, recombinant, and phage display technologies, or a combination thereof. For example, monoclonal antibodies can be produced using hybridoma techniques including those known in the art and taught, for example, in Harlow et al., *Antibodies: A Laboratory Manual*, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988); Hammerling, et al., in: *Monoclonal Antibodies and T-Cell Hybridomas* 563-681 (Elsevier, N.Y., 1981) (said references incorporated by reference in their entireties). The term "monoclonal antibody" as used herein is not limited to antibodies produced through hybridoma technology. The term "monoclonal antibody" refers to an antibody that is derived from a single clone, including any eukaryotic, prokaryotic, or phage clone, and not the method by which it is produced.

Methods for producing and screening for specific antibodies using hybridoma technology are routine and well known in the art and are discussed in detail in the Examples (e.g., Example 16). In a non-limiting example, mice can be immunized with a polypeptide of the invention or a cell expressing such peptide. Once an
5 immune response is detected, e.g., antibodies specific for the antigen are detected in the mouse serum, the mouse spleen is harvested and splenocytes isolated. The splenocytes are then fused by well known techniques to any suitable myeloma cells, for example cells from cell line SP20 available from the ATCC. Hybridomas are selected and cloned by limited dilution. The hybridoma clones are then assayed by
10 methods known in the art for cells that secrete antibodies capable of binding a polypeptide of the invention. Ascites fluid, which generally contains high levels of antibodies, can be generated by immunizing mice with positive hybridoma clones.

Accordingly, the present invention provides methods of generating monoclonal antibodies as well as antibodies produced by the method comprising
15 culturing a hybridoma cell secreting an antibody of the invention wherein, preferably, the hybridoma is generated by fusing splenocytes isolated from a mouse immunized with an antigen of the invention with myeloma cells and then screening the hybridomas resulting from the fusion for hybridoma clones that secrete an antibody able to bind a polypeptide of the invention.

20 Antibody fragments which recognize specific epitopes may be generated by known techniques. For example, Fab and F(ab')₂ fragments of the invention may be produced by proteolytic cleavage of immunoglobulin molecules, using enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab')₂ fragments). F(ab')₂ fragments contain the variable region, the light chain constant region and the
25 CH1 domain of the heavy chain.

For example, the antibodies of the present invention can also be generated using various phage display methods known in the art. In phage display methods, functional antibody domains are displayed on the surface of phage particles which carry the polynucleotide sequences encoding them. In a particular embodiment, such
30 phage can be utilized to display antigen binding domains expressed from a repertoire or combinatorial antibody library (e.g., human or murine). Phage expressing an

antigen binding domain that binds the antigen of interest can be selected or identified with antigen, e.g., using labeled antigen or antigen bound or captured to a solid surface or bead. Phage used in these methods are typically filamentous phage including fd and M13 binding domains expressed from phage with Fab, Fv or disulfide stabilized Fv antibody domains recombinantly fused to either the phage gene III or gene VIII protein. Examples of phage display methods that can be used to make the antibodies of the present invention include those disclosed in Brinkman et al., *J. Immunol. Methods* 182:41-50 (1995); Ames et al., *J. Immunol. Methods* 184:177-186 (1995); Kettleborough et al., *Eur. J. Immunol.* 24:952-958 (1994); Persic et al., *Gene* 187 9-18 (1997); Burton et al., *Advances in Immunology* 57:191-280 (1994); PCT application No. PCT/GB91/01134; PCT publications WO 90/02809; WO 91/10737; WO 92/01047; WO 92/18619; WO 93/11236; WO 95/15982; WO 95/20401; and U.S. Patent Nos. 5,698,426; 5,223,409; 5,403,484; 5,580,717; 5,427,908; 5,750,753; 5,821,047; 5,571,698; 5,427,908; 5,516,637; 5,780,225; 5,658,727; 5,733,743 and 5,969,108; each of which is incorporated herein by reference in its entirety.

As described in the above references, after phage selection, the antibody coding regions from the phage can be isolated and used to generate whole antibodies, including human antibodies, or any other desired antigen binding fragment, and expressed in any desired host, including mammalian cells, insect cells, plant cells, yeast, and bacteria, e.g., as described in detail below. For example, techniques to recombinantly produce Fab, Fab' and F(ab')₂ fragments can also be employed using methods known in the art such as those disclosed in PCT publication WO 92/22324; Mullinax et al., *BioTechniques* 12(6):864-869 (1992); and Sawai et al., *AJRI* 34:26-34 (1995); and Better et al., *Science* 240:1041-1043 (1988) (said references incorporated by reference in their entireties).

Examples of techniques which can be used to produce single-chain Fvs and antibodies include those described in U.S. Patents 4,946,778 and 5,258,498; Huston et al., *Methods in Enzymology* 203:46-88 (1991); Shu et al., *PNAS* 90:7995-7999 (1993); and Skerra et al., *Science* 240:1038-1040 (1988). For some uses, including in vivo use of antibodies in humans and in vitro detection assays, it may be preferable

to use chimeric, humanized, or human antibodies. A chimeric antibody is a molecule in which different portions of the antibody are derived from different animal species, such as antibodies having a variable region derived from a murine monoclonal antibody and a human immunoglobulin constant region. Methods for producing chimeric antibodies are known in the art. See e.g., Morrison, Science 229:1202 (1985); Oi et al., BioTechniques 4:214 (1986); Gillies et al., (1989) J. Immunol. Methods 125:191-202; U.S. Patent Nos. 5,807,715; 4,816,567; and 4,816,397, which are incorporated herein by reference in their entirety. Humanized antibodies are antibody molecules from non-human species antibody that binds the desired antigen having one or more complementarity determining regions (CDRs) from the non-human species and a framework regions from a human immunoglobulin molecule. Often, framework residues in the human framework regions will be substituted with the corresponding residue from the CDR donor antibody to alter, preferably improve, antigen binding. These framework substitutions are identified by methods well known in the art, e.g., by modeling of the interactions of the CDR and framework residues to identify framework residues important for antigen binding and sequence comparison to identify unusual framework residues at particular positions. (See, e.g., Queen et al., U.S. Patent No. 5,585,089; Riechmann et al., Nature 332:323 (1988), which are incorporated herein by reference in their entireties.) Antibodies can be humanized using a variety of techniques known in the art including, for example, CDR-grafting (EP 239,400; PCT publication WO 91/09967; U.S. Patent Nos. 5,225,539; 5,530,101; and 5,585,089), veneering or resurfacing (EP 592,106; EP 519,596; Padlan, Molecular Immunology 28(4/5):489-498 (1991); Studnicka et al., Protein Engineering 7(6):805-814 (1994); Roguska. et al., PNAS 91:969-973 (1994)), and chain shuffling (U.S. Patent No. 5,565,332).

Completely human antibodies are particularly desirable for therapeutic treatment of human patients. Human antibodies can be made by a variety of methods known in the art including phage display methods described above using antibody libraries derived from human immunoglobulin sequences. See also, U.S. Patent Nos. 4,444,887 and 4,716,111; and PCT publications WO 98/46645, WO 98/50433, WO

98/24893, WO 98/16654, WO 96/34096, WO 96/33735, and WO 91/10741; each of which is incorporated herein by reference in its entirety.

Human antibodies can also be produced using transgenic mice which are incapable of expressing functional endogenous immunoglobulins, but which can
5 express human immunoglobulin genes. For example, the human heavy and light chain immunoglobulin gene complexes may be introduced randomly or by homologous recombination into mouse embryonic stem cells. Alternatively, the human variable region, constant region, and diversity region may be introduced into mouse embryonic stem cells in addition to the human heavy and light chain genes.
10 The mouse heavy and light chain immunoglobulin genes may be rendered non-functional separately or simultaneously with the introduction of human immunoglobulin loci by homologous recombination. In particular, homozygous deletion of the JH region prevents endogenous antibody production. The modified embryonic stem cells are expanded and microinjected into blastocysts to produce
15 chimeric mice. The chimeric mice are then bred to produce homozygous offspring which express human antibodies. The transgenic mice are immunized in the normal fashion with a selected antigen, e.g., all or a portion of a polypeptide of the invention. Monoclonal antibodies directed against the antigen can be obtained from the immunized, transgenic mice using conventional hybridoma technology. The human
20 immunoglobulin transgenes harbored by the transgenic mice rearrange during B cell differentiation, and subsequently undergo class switching and somatic mutation. Thus, using such a technique, it is possible to produce therapeutically useful IgG, IgA, IgM and IgE antibodies. For an overview of this technology for producing human antibodies, see Lonberg and Huszar, *Int. Rev. Immunol.* 13:65-93 (1995). For a
25 detailed discussion of this technology for producing human antibodies and human monoclonal antibodies and protocols for producing such antibodies, see, e.g., PCT publications WO 98/24893; WO 92/01047; WO 96/34096; WO 96/33735; European Patent No. 0 598 877; U.S. Patent Nos. 5,413,923; 5,625,126; 5,633,425; 5,569,825; 5,661,016; 5,545,806; 5,814,318; 5,885,793; 5,916,771; and 5,939,598, which are
30 incorporated by reference herein in their entirety. In addition, companies such as Abgenix, Inc. (Freemont, CA) and Genpharm (San Jose, CA) can be engaged to

provide human antibodies directed against a selected antigen using technology similar to that described above.

Completely human antibodies which recognize a selected epitope can be generated using a technique referred to as "guided selection." In this approach a
5 selected non-human monoclonal antibody, e.g., a mouse antibody, is used to guide the selection of a completely human antibody recognizing the same epitope. (Jespers et al., Bio/technology 12:899-903 (1988)).

Further, antibodies to the polypeptides of the invention can, in turn, be utilized to generate anti-idiotypic antibodies that "mimic" polypeptides of the invention using
10 techniques well known to those skilled in the art. (See, e.g., Greenspan & Bona, FASEB J. 7(5):437-444; (1989) and Nissinoff, J. Immunol. 147(8):2429-2438 (1991)). For example, antibodies which bind to and competitively inhibit polypeptide multimerization and/or binding of a polypeptide of the invention to a ligand can be used to generate anti-idiotypes that "mimic" the polypeptide multimerization and/or
15 binding domain and, as a consequence, bind to and neutralize polypeptide and/or its ligand. Such neutralizing anti-idiotypes or Fab fragments of such anti-idiotypes can be used in therapeutic regimens to neutralize polypeptide ligand. For example, such anti-idiotypic antibodies can be used to bind a polypeptide of the invention and/or to bind its ligands/receptors, and thereby block its biological activity.

20

Polynucleotides Encoding Antibodies

The invention further provides polynucleotides comprising a nucleotide sequence encoding an antibody of the invention and fragments thereof. The invention also encompasses polynucleotides that hybridize under stringent or lower
25 stringency hybridization conditions, e.g., as defined supra, to polynucleotides that encode an antibody, preferably, that specifically binds to a polypeptide of the invention, preferably, an antibody that binds to a polypeptide having the amino acid sequence of SEQ ID NO:Y.

The polynucleotides may be obtained, and the nucleotide sequence of the
30 polynucleotides determined, by any method known in the art. For example, if the nucleotide sequence of the antibody is known, a polynucleotide encoding the antibody

may be assembled from chemically synthesized oligonucleotides (e.g., as described in Kutmeier et al., *BioTechniques* 17:242 (1994)), which, briefly, involves the synthesis of overlapping oligonucleotides containing portions of the sequence encoding the antibody, annealing and ligating of those oligonucleotides, and then
5 amplification of the ligated oligonucleotides by PCR.

Alternatively, a polynucleotide encoding an antibody may be generated from nucleic acid from a suitable source. If a clone containing a nucleic acid encoding a particular antibody is not available, but the sequence of the antibody molecule is known, a nucleic acid encoding the immunoglobulin may be chemically synthesized
10 or obtained from a suitable source (e.g., an antibody cDNA library, or a cDNA library generated from, or nucleic acid, preferably poly A+ RNA, isolated from, any tissue or cells expressing the antibody, such as hybridoma cells selected to express an antibody of the invention) by PCR amplification using synthetic primers hybridizable to the 3' and 5' ends of the sequence or by cloning using an oligonucleotide probe
15 specific for the particular gene sequence to identify, e.g., a cDNA clone from a cDNA library that encodes the antibody. Amplified nucleic acids generated by PCR may then be cloned into replicable cloning vectors using any method well known in the art.

Once the nucleotide sequence and corresponding amino acid sequence of the
20 antibody is determined, the nucleotide sequence of the antibody may be manipulated using methods well known in the art for the manipulation of nucleotide sequences, e.g., recombinant DNA techniques, site directed mutagenesis, PCR, etc. (see, for example, the techniques described in Sambrook et al., 1990, *Molecular Cloning, A Laboratory Manual*, 2d Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor,
25 NY and Ausubel et al., eds., 1998, *Current Protocols in Molecular Biology*, John Wiley & Sons, NY, which are both incorporated by reference herein in their entireties), to generate antibodies having a different amino acid sequence, for example to create amino acid substitutions, deletions, and/or insertions.

In a specific embodiment, the amino acid sequence of the heavy and/or light
30 chain variable domains may be inspected to identify the sequences of the complementarity determining regions (CDRs) by methods that are well known in the

art, e.g., by comparison to known amino acid sequences of other heavy and light chain variable regions to determine the regions of sequence hypervariability. Using routine recombinant DNA techniques, one or more of the CDRs may be inserted within framework regions, e.g., into human framework regions to humanize a non-
5 human antibody, as described supra. The framework regions may be naturally occurring or consensus framework regions, and preferably human framework regions (see, e.g., Chothia et al., J. Mol. Biol. 278: 457-479 (1998) for a listing of human framework regions). Preferably, the polynucleotide generated by the combination of the framework regions and CDRs encodes an antibody that specifically binds a
10 polypeptide of the invention. Preferably, as discussed supra, one or more amino acid substitutions may be made within the framework regions, and, preferably, the amino acid substitutions improve binding of the antibody to its antigen. Additionally, such methods may be used to make amino acid substitutions or deletions of one or more variable region cysteine residues participating in an intrachain disulfide bond to
15 generate antibody molecules lacking one or more intrachain disulfide bonds. Other alterations to the polynucleotide are encompassed by the present invention and within the skill of the art.

In addition, techniques developed for the production of "chimeric antibodies" (Morrison et al., Proc. Natl. Acad. Sci. 81:851-855 (1984); Neuberger et al., Nature
20 312:604-608 (1984); Takeda et al., Nature 314:452-454 (1985)) by splicing genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity can be used. As described supra, a chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived
25 from a murine mAb and a human immunoglobulin constant region, e.g., humanized antibodies.

Alternatively, techniques described for the production of single chain antibodies (U.S. Patent No. 4,946,778; Bird, Science 242:423-42 (1988); Huston et al., Proc. Natl. Acad. Sci. USA 85:5879-5883 (1988); and Ward et al., Nature
30 334:544-54 (1989)) can be adapted to produce single chain antibodies. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region

via an amino acid bridge, resulting in a single chain polypeptide. Techniques for the assembly of functional Fv fragments in *E. coli* may also be used (Skerra et al., Science 242:1038- 1041 (1988)).

5 Methods of Producing Antibodies

The antibodies of the invention can be produced by any method known in the art for the synthesis of antibodies, in particular, by chemical synthesis or preferably, by recombinant expression techniques.

Recombinant expression of an antibody of the invention, or fragment,
10 derivative or analog thereof, (e.g., a heavy or light chain of an antibody of the invention or a single chain antibody of the invention), requires construction of an expression vector containing a polynucleotide that encodes the antibody. Once a polynucleotide encoding an antibody molecule or a heavy or light chain of an antibody, or portion thereof (preferably containing the heavy or light chain variable
15 domain), of the invention has been obtained, the vector for the production of the antibody molecule may be produced by recombinant DNA technology using techniques well known in the art. Thus, methods for preparing a protein by expressing a polynucleotide containing an antibody encoding nucleotide sequence are described herein. Methods which are well known to those skilled in the art can be
20 used to construct expression vectors containing antibody coding sequences and appropriate transcriptional and translational control signals. These methods include, for example, in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. The invention, thus, provides replicable vectors comprising a nucleotide sequence encoding an antibody molecule of the invention, or a heavy or
25 light chain thereof, or a heavy or light chain variable domain, operably linked to a promoter. Such vectors may include the nucleotide sequence encoding the constant region of the antibody molecule (see, e.g., PCT Publication WO 86/05807; PCT Publication WO 89/01036; and U.S. Patent No. 5,122,464) and the variable domain of the antibody may be cloned into such a vector for expression of the entire heavy or
30 light chain.

The expression vector is transferred to a host cell by conventional techniques and the transfected cells are then cultured by conventional techniques to produce an antibody of the invention. Thus, the invention includes host cells containing a polynucleotide encoding an antibody of the invention, or a heavy or light chain thereof, or a single chain antibody of the invention, operably linked to a heterologous promoter. In preferred embodiments for the expression of double-chained antibodies, vectors encoding both the heavy and light chains may be co-expressed in the host cell for expression of the entire immunoglobulin molecule, as detailed below.

A variety of host-expression vector systems may be utilized to express the antibody molecules of the invention. Such host-expression systems represent vehicles by which the coding sequences of interest may be produced and subsequently purified, but also represent cells which may, when transformed or transfected with the appropriate nucleotide coding sequences, express an antibody molecule of the invention in situ. These include but are not limited to microorganisms such as bacteria (e.g., *E. coli*, *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing antibody coding sequences; yeast (e.g., *Saccharomyces*, *Pichia*) transformed with recombinant yeast expression vectors containing antibody coding sequences; insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing antibody coding sequences; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing antibody coding sequences; or mammalian cell systems (e.g., COS, CHO, BHK, 293, 3T3 cells) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter). Preferably, bacterial cells such as *Escherichia coli*, and more preferably, eukaryotic cells, especially for the expression of whole recombinant antibody molecule, are used for the expression of a recombinant antibody molecule. For example, mammalian cells such as Chinese hamster ovary cells (CHO), in conjunction with a vector such as the major intermediate early gene

promoter element from human cytomegalovirus is an effective expression system for antibodies (Foecking et al., Gene 45:101 (1986); Cockett et al., Bio/Technology 8:2 (1990)).

In bacterial systems, a number of expression vectors may be advantageously
5 selected depending upon the use intended for the antibody molecule being expressed. For example, when a large quantity of such a protein is to be produced, for the generation of pharmaceutical compositions of an antibody molecule, vectors which direct the expression of high levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not limited, to the *E. coli* expression
10 vector pUR278 (Ruther et al., EMBO J. 2:1791 (1983)), in which the antibody coding sequence may be ligated individually into the vector in frame with the lac Z coding region so that a fusion protein is produced; pIN vectors (Inouye & Inouye, Nucleic Acids Res. 13:3101-3109 (1985); Van Heeke & Schuster, J. Biol. Chem. 24:5503-5509 (1989)); and the like. pGEX vectors may also be used to express foreign
15 polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption and binding to matrix glutathione-agarose beads followed by elution in the presence of free glutathione. The pGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene product can be released from the
20 GST moiety.

In an insect system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in
Spodoptera frugiperda cells. The antibody coding sequence may be cloned
25 individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter).

In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, the antibody coding sequence of interest may be ligated to an adenovirus transcription/translation
30 control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by in vitro or in vivo

recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing the antibody molecule in infected hosts. (e.g., see Logan & Shenk, Proc. Natl. Acad. Sci. USA 81:355-359 (1984)). Specific initiation signals may also be required for efficient translation of inserted antibody coding sequences. These signals include the ATG initiation codon and adjacent sequences. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (see Bittner et al., Methods in Enzymol. 153:51-544 (1987)).

In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and gene products. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end, eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such mammalian host cells include but are not limited to CHO, VERY, BHK, HeLa, COS, MDCK, 293, 3T3, WI38, and in particular, breast cancer cell lines such as, for example, BT483, Hs578T, HTB2, BT20 and T47D, and normal mammary gland cell line such as, for example, CRL7030 and Hs578Bst.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express the antibody molecule may be engineered. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (e.g., promoter, enhancer, sequences,

transcription terminators, polyadenylation sites, etc.), and a selectable marker.

Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media.

The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines which express the antibody molecule. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that interact directly or indirectly with the antibody molecule.

A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler et al., *Cell* 11:223 (1977)), hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, *Proc. Natl. Acad. Sci. USA* 48:202 (1992)), and adenine phosphoribosyltransferase (Lowy et al., *Cell* 22:817 (1980)) genes can be employed in tk-, hgpri- or apri- cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: dhfr, which confers resistance to methotrexate (Wigler et al., *Natl. Acad. Sci. USA* 77:357 (1980); O'Hare et al., *Proc. Natl. Acad. Sci. USA* 78:1527 (1981)); gpt, which confers resistance to mycophenolic acid (Mulligan & Berg, *Proc. Natl. Acad. Sci. USA* 78:2072 (1981)); neo, which confers resistance to the aminoglycoside G-418 (*Clinical Pharmacy* 12:488-505; Wu and Wu, *Biotherapy* 3:87-95 (1991); Tolstoshev, *Ann. Rev. Pharmacol. Toxicol.* 32:573-596 (1993); Mulligan, *Science* 260:926-932 (1993); and Morgan and Anderson, *Ann. Rev. Biochem.* 62:191-217 (1993); May, 1993, *TIB TECH* 11(5):155-215); and hygromycin (Santerre et al., *Gene* 30:147 (1984)). Methods commonly known in the art of recombinant DNA technology may be routinely applied to select the desired recombinant clone, and such methods are described, for example, in Ausubel et al. (eds.), *Current Protocols in Molecular Biology*. John Wiley & Sons, NY (1993); Kriegler, *Gene Transfer and Expression, A Laboratory Manual*, Stockton Press, NY (1990); and in Chapters 12 and 13, Dracopoli et al. (eds), *Current Protocols in Human Genetics*, John Wiley & Sons, NY (1994); Colberre-Garapin et al., *J. Mol. Biol.* 150:1 (1981), which are incorporated by reference herein in their entireties.

The expression levels of an antibody molecule can be increased by vector amplification (for a review, see Bebbington and Hentschel, The use of vectors based on gene amplification for the expression of cloned genes in mammalian cells in DNA cloning, Vol.3. (Academic Press, New York, 1987)). When a marker in the vector
5 system expressing antibody is amplifiable, increase in the level of inhibitor present in culture of host cell will increase the number of copies of the marker gene. Since the amplified region is associated with the antibody gene, production of the antibody will also increase (Crouse et al., Mol. Cell. Biol. 3:257 (1983)).

The host cell may be co-transfected with two expression vectors of the
10 invention, the first vector encoding a heavy chain derived polypeptide and the second vector encoding a light chain derived polypeptide. The two vectors may contain identical selectable markers which enable equal expression of heavy and light chain polypeptides. Alternatively, a single vector may be used which encodes, and is capable of expressing, both heavy and light chain polypeptides. In such situations,
15 the light chain should be placed before the heavy chain to avoid an excess of toxic free heavy chain (Proudfoot, Nature 322:52 (1986); Kohler, Proc. Natl. Acad. Sci. USA 77:2197 (1980)). The coding sequences for the heavy and light chains may comprise cDNA or genomic DNA.

Once an antibody molecule of the invention has been produced by an animal,
20 chemically synthesized, or recombinantly expressed, it may be purified by any method known in the art for purification of an immunoglobulin molecule, for example, by chromatography (e.g., ion exchange, affinity, particularly by affinity for the specific antigen after Protein A, and sizing column chromatography), centrifugation, differential solubility, or by any other standard technique for the
25 purification of proteins. In addition, the antibodies of the present invention or fragments thereof can be fused to heterologous polypeptide sequences described herein or otherwise known in the art, to facilitate purification.

The present invention encompasses antibodies recombinantly fused or chemically conjugated (including both covalently and non-covalently conjugations)
30 to a polypeptide (or portion thereof, preferably at least 10, 20, 30, 40, 50, 60, 70, 80, 90 or 100 amino acids of the polypeptide) of the present invention to generate fusion

proteins. The fusion does not necessarily need to be direct, but may occur through linker sequences. The antibodies may be specific for antigens other than polypeptides (or portion thereof, preferably at least 10, 20, 30, 40, 50, 60, 70, 80, 90 or 100 amino acids of the polypeptide) of the present invention. For example, antibodies may be
5 used to target the polypeptides of the present invention to particular cell types, either in vitro or in vivo, by fusing or conjugating the polypeptides of the present invention to antibodies specific for particular cell surface receptors. Antibodies fused or conjugated to the polypeptides of the present invention may also be used in in vitro immunoassays and purification methods using methods known in the art. See e.g.,
10 Harbor et al., supra, and PCT publication WO 93/21232; EP 439,095; Naramura et al., Immunol. Lett. 39:91-99 (1994); U.S. Patent 5,474,981; Gillies et al., PNAS 89:1428-1432 (1992); Fell et al., J. Immunol. 146:2446-2452(1991), which are incorporated by reference in their entirety.

The present invention further includes compositions comprising the
15 polypeptides of the present invention fused or conjugated to antibody domains other than the variable regions. For example, the polypeptides of the present invention may be fused or conjugated to an antibody Fc region, or portion thereof. The antibody portion fused to a polypeptide of the present invention may comprise the constant region, hinge region, CH1 domain, CH2 domain, and CH3 domain or any
20 combination of whole domains or portions thereof. The polypeptides may also be fused or conjugated to the above antibody portions to form multimers. For example, Fc portions fused to the polypeptides of the present invention can form dimers through disulfide bonding between the Fc portions. Higher multimeric forms can be made by fusing the polypeptides to portions of IgA and IgM. Methods for fusing or
25 conjugating the polypeptides of the present invention to antibody portions are known in the art. See, e.g., U.S. Patent Nos. 5,336,603; 5,622,929; 5,359,046; 5,349,053; 5,447,851; 5,112,946; EP 307,434; EP 367,166; PCT publications WO 96/04388; WO 91/06570; Ashkenazi et al., Proc. Natl. Acad. Sci. USA 88:10535-10539 (1991); Zheng et al., J. Immunol. 154:5590-5600 (1995); and Vil et al., Proc. Natl. Acad. Sci.
30 USA 89:11337-11341(1992) (said references incorporated by reference in their entirety).

As discussed, supra, the polypeptides corresponding to a polypeptide, polypeptide fragment, or a variant of SEQ ID NO:Y may be fused or conjugated to the above antibody portions to increase the in vivo half life of the polypeptides or for use in immunoassays using methods known in the art. Further, the polypeptides
5 corresponding to SEQ ID NO:Y may be fused or conjugated to the above antibody portions to facilitate purification. One reported example describes chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins. (EP 394,827; Traunecker et al., *Nature* 331:84-86 (1988)). The
10 polypeptides of the present invention fused or conjugated to an antibody having disulfide-linked dimeric structures (due to the IgG) may also be more efficient in binding and neutralizing other molecules, than the monomeric secreted protein or protein fragment alone. (Fountoulakis et al., *J. Biochem.* 270:3958-3964 (1995)). In many cases, the Fc part in a fusion protein is beneficial in therapy and diagnosis, and
15 thus can result in, for example, improved pharmacokinetic properties. (EP A 232,262). Alternatively, deleting the Fc part after the fusion protein has been expressed, detected, and purified, would be desired. For example, the Fc portion may hinder therapy and diagnosis if the fusion protein is used as an antigen for immunizations. In drug discovery, for example, human proteins, such as hIL-5, have
20 been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists of hIL-5. (See, Bennett et al., *J. Molecular Recognition* 8:52-58 (1995); Johanson et al., *J. Biol. Chem.* 270:9459-9471 (1995)).

Moreover, the antibodies or fragments thereof of the present invention can be fused to marker sequences, such as a peptide to facilitate purification. In preferred
25 embodiments, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311), among others, many of which are commercially available. As described in Gentz et al., *Proc. Natl. Acad. Sci. USA* 86:821-824 (1989), for instance, hexa-histidine provides for convenient purification of the fusion protein. Other peptide tags
30 useful for purification include, but are not limited to, the "HA" tag, which

corresponds to an epitope derived from the influenza hemagglutinin protein (Wilson et al., Cell 37:767 (1984)) and the "flag" tag.

The present invention further encompasses antibodies or fragments thereof conjugated to a diagnostic or therapeutic agent. The antibodies can be used
5 diagnostically to, for example, monitor the development or progression of a tumor as part of a clinical testing procedure to, e.g., determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent
10 materials, radioactive materials, positron emitting metals using various positron emission tomographies, and nonradioactive paramagnetic metal ions. The detectable substance may be coupled or conjugated either directly to the antibody (or fragment thereof) or indirectly, through an intermediate (such as, for example, a linker known in the art) using techniques known in the art. See, for example, U.S. Patent No.
15 4,741,900 for metal ions which can be conjugated to antibodies for use as diagnostics according to the present invention. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, beta-galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone,
20 fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin; and examples of suitable radioactive material include ¹²⁵I, ¹³¹I, ¹¹¹In or ⁹⁹Tc.
25 Further, an antibody or fragment thereof may be conjugated to a therapeutic moiety such as a cytotoxin, e.g., a cytostatic or cytotoxic agent, a therapeutic agent or a radioactive metal ion, e.g., alpha-emitters such as, for example, ²¹³Bi. A cytotoxin or cytotoxic agent includes any agent that is detrimental to cells. Examples include paclitaxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin,
30 etoposide, teniposide, vincristine, vinblastine, colchicine, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-

dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof. Therapeutic agents include, but are not limited to, antimetabolites (e.g., methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (e.g., mechlorethamine, thioepa chlorambucil, melphalan, carmustine (BSNU) and lomustine (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cis-dichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (e.g., daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (e.g., dactinomycin (formerly actinomycin), bleomycin, mithramycin, and anthramycin (AMC)), and anti-mitotic agents (e.g., vincristine and vinblastine).

The conjugates of the invention can be used for modifying a given biological response, the therapeutic agent or drug moiety is not to be construed as limited to classical chemical therapeutic agents. For example, the drug moiety may be a protein or polypeptide possessing a desired biological activity. Such proteins may include, for example, a toxin such as abrin, ricin A, pseudomonas exotoxin, or diphtheria toxin; a protein such as tumor necrosis factor, α -interferon, β -interferon, nerve growth factor, platelet derived growth factor, tissue plasminogen activator, an apoptotic agent, e.g., TNF- α , TNF- β , AIM I (See, International Publication No. WO 97/33899), AIM II (See, International Publication No. WO 97/34911), Fas Ligand (Takahashi *et al.*, *Int. Immunol.*, 6:1567-1574 (1994)), VEGI (See, International Publication No. WO 99/23105), a thrombotic agent or an anti-angiogenic agent, e.g., angiostatin or endostatin; or, biological response modifiers such as, for example, lymphokines, interleukin-1 ("IL-1"), interleukin-2 ("IL-2"), interleukin-6 ("IL-6"), granulocyte macrophage colony stimulating factor ("GM-CSF"), granulocyte colony stimulating factor ("G-CSF"), or other growth factors.

Antibodies may also be attached to solid supports, which are particularly useful for immunoassays or purification of the target antigen. Such solid supports include, but are not limited to, glass, cellulose, polyacrylamide, nylon, polystyrene, polyvinyl chloride or polypropylene.

Techniques for conjugating such therapeutic moiety to antibodies are well known, see, e.g., Arnon *et al.*, "Monoclonal Antibodies For Immunotargeting Of

Drugs In Cancer Therapy", in *Monoclonal Antibodies And Cancer Therapy*, Reisfeld et al. (eds.), pp. 243-56 (Alan R. Liss, Inc. 1985); Hellstrom et al., "Antibodies For Drug Delivery", in *Controlled Drug Delivery* (2nd Ed.), Robinson et al. (eds.), pp. 623-53 (Marcel Dekker, Inc. 1987); Thorpe, "Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review", in *Monoclonal Antibodies '84: Biological And Clinical Applications*, Pinchera et al. (eds.), pp. 475-506 (1985); "Analysis, Results, And Future Prospective Of The Therapeutic Use Of Radiolabeled Antibody In Cancer Therapy", in *Monoclonal Antibodies For Cancer Detection And Therapy*, Baldwin et al. (eds.), pp. 303-16 (Academic Press 1985), and Thorpe et al., "The Preparation And Cytotoxic Properties Of Antibody-Toxin Conjugates", *Immunol. Rev.* 62:119-58 (1982).

Alternatively, an antibody can be conjugated to a second antibody to form an antibody heteroconjugate as described by Segal in U.S. Patent No. 4,676,980, which is incorporated herein by reference in its entirety.

15 An antibody, with or without a therapeutic moiety conjugated to it, administered alone or in combination with cytotoxic factor(s) and/or cytokine(s) can be used as a therapeutic.

Immunophenotyping

20 The antibodies of the invention may be utilized for immunophenotyping of cell lines and biological samples. The translation product of the gene of the present invention may be useful as a cell specific marker, or more specifically as a cellular marker that is differentially expressed at various stages of differentiation and/or maturation of particular cell types. Monoclonal antibodies directed against a specific epitope, or combination of epitopes, will allow for the screening of cellular populations expressing the marker. Various techniques can be utilized using monoclonal antibodies to screen for cellular populations expressing the marker(s), and include magnetic separation using antibody-coated magnetic beads, "panning" with antibody attached to a solid matrix (i.e., plate), and flow cytometry (See, e.g., U.S. Patent 5,985,660; and Morrison *et al.*, *Cell*, 96:737-49 (1999)).

These techniques allow for the screening of particular populations of cells, such as might be found with hematological malignancies (i.e. minimal residual disease (MRD) in acute leukemic patients) and "non-self" cells in transplantations to prevent Graft-versus-Host Disease (GVHD). Alternatively, these techniques allow for the screening of hematopoietic stem and progenitor cells capable of undergoing proliferation and/or differentiation, as might be found in human umbilical cord blood.

Assays For Antibody Binding

The antibodies of the invention may be assayed for immunospecific binding by any method known in the art. The immunoassays which can be used include but are not limited to competitive and non-competitive assay systems using techniques such as western blots, radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoprecipitation assays, precipitin reactions, gel diffusion precipitin reactions, immunodiffusion assays, agglutination assays, complement-fixation assays, immunoradiometric assays, fluorescent immunoassays, protein A immunoassays, to name but a few. Such assays are routine and well known in the art (see, e.g., Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York, which is incorporated by reference herein in its entirety). Exemplary immunoassays are described briefly below (but are not intended by way of limitation).

Immunoprecipitation protocols generally comprise lysing a population of cells in a lysis buffer such as RIPA buffer (1% NP-40 or Triton X-100, 1% sodium deoxycholate, 0.1% SDS, 0.15 M NaCl, 0.01 M sodium phosphate at pH 7.2, 1% Trasylol) supplemented with protein phosphatase and/or protease inhibitors (e.g., EDTA, PMSF, aprotinin, sodium vanadate), adding the antibody of interest to the cell lysate, incubating for a period of time (e.g., 1-4 hours) at 4° C, adding protein A and/or protein G sepharose beads to the cell lysate, incubating for about an hour or more at 4° C, washing the beads in lysis buffer and resuspending the beads in SDS/sample buffer. The ability of the antibody of interest to immunoprecipitate a particular antigen can be assessed by, e.g., western blot analysis. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the

binding of the antibody to an antigen and decrease the background (e.g., pre-clearing the cell lysate with sepharose beads). For further discussion regarding immunoprecipitation protocols see, e.g., Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York at 10.16.1.

5 Western blot analysis generally comprises preparing protein samples, electrophoresis of the protein samples in a polyacrylamide gel (e.g., 8%- 20% SDS-PAGE depending on the molecular weight of the antigen), transferring the protein sample from the polyacrylamide gel to a membrane such as nitrocellulose, PVDF or nylon, blocking the membrane in blocking solution (e.g., PBS with 3% BSA or non-fat milk), washing the membrane in washing buffer (e.g., PBS-Tween 20), blocking
10 the membrane with primary antibody (the antibody of interest) diluted in blocking buffer, washing the membrane in washing buffer, blocking the membrane with a secondary antibody (which recognizes the primary antibody, e.g., an anti-human antibody) conjugated to an enzymatic substrate (e.g., horseradish peroxidase or
15 alkaline phosphatase) or radioactive molecule (e.g., ^{32}P or ^{125}I) diluted in blocking buffer, washing the membrane in wash buffer, and detecting the presence of the antigen. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the signal detected and to reduce the background noise. For further discussion regarding western blot protocols see, e.g., Ausubel et al, eds, 1994,
20 Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York at 10.8.1.

ELISAs comprise preparing antigen, coating the well of a 96 well microtiter plate with the antigen, adding the antibody of interest conjugated to a detectable compound such as an enzymatic substrate (e.g., horseradish peroxidase or alkaline
25 phosphatase) to the well and incubating for a period of time, and detecting the presence of the antigen. In ELISAs the antibody of interest does not have to be conjugated to a detectable compound; instead, a second antibody (which recognizes the antibody of interest) conjugated to a detectable compound may be added to the well. Further, instead of coating the well with the antigen, the antibody may be
30 coated to the well. In this case, a second antibody conjugated to a detectable compound may be added following the addition of the antigen of interest to the

coated well. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the signal detected as well as other variations of ELISAs known in the art. For further discussion regarding ELISAs see, e.g., Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc.,
5 New York at 11.2.1.

The binding affinity of an antibody to an antigen and the off-rate of an antibody-antigen interaction can be determined by competitive binding assays. One example of a competitive binding assay is a radioimmunoassay comprising the incubation of labeled antigen (e.g., ³H or ¹²⁵I) with the antibody of interest in the
10 presence of increasing amounts of unlabeled antigen, and the detection of the antibody bound to the labeled antigen. The affinity of the antibody of interest for a particular antigen and the binding off-rates can be determined from the data by scatchard plot analysis. Competition with a second antibody can also be determined using radioimmunoassays. In this case, the antigen is incubated with antibody of
15 interest conjugated to a labeled compound (e.g., ³H or ¹²⁵I) in the presence of increasing amounts of an unlabeled second antibody.

Therapeutic Uses

The present invention is further directed to antibody-based therapies which
20 involve administering antibodies of the invention to an animal, preferably a mammal, and most preferably a human, patient for treating one or more of the disclosed diseases, disorders, or conditions. Therapeutic compounds of the invention include, but are not limited to, antibodies of the invention (including fragments, analogs and derivatives thereof as described herein) and nucleic acids encoding antibodies of the
25 invention (including fragments, analogs and derivatives thereof and anti-idiotypic antibodies as described herein). The antibodies of the invention can be used to treat, inhibit or prevent diseases, disorders or conditions associated with aberrant expression and/or activity of a polypeptide of the invention, including, but not limited to, any one or more of the diseases, disorders, or conditions described herein. The treatment
30 and/or prevention of diseases, disorders, or conditions associated with aberrant expression and/or activity of a polypeptide of the invention includes, but is not

limited to, alleviating symptoms associated with those diseases, disorders or conditions. Antibodies of the invention may be provided in pharmaceutically acceptable compositions as known in the art or as described herein.

A summary of the ways in which the antibodies of the present invention may
5 be used therapeutically includes binding polynucleotides or polypeptides of the present invention locally or systemically in the body or by direct cytotoxicity of the antibody, e.g. as mediated by complement (CDC) or by effector cells (ADCC). Some of these approaches are described in more detail below. Armed with the teachings provided herein, one of ordinary skill in the art will know how to use the
10 antibodies of the present invention for diagnostic, monitoring or therapeutic purposes without undue experimentation.

The antibodies of this invention may be advantageously utilized in combination with other monoclonal or chimeric antibodies, or with lymphokines or hematopoietic growth factors (such as, e.g., IL-2, IL-3 and IL-7), for example, which
15 serve to increase the number or activity of effector cells which interact with the antibodies.

The antibodies of the invention may be administered alone or in combination with other types of treatments (e.g., radiation therapy, chemotherapy, hormonal therapy, immunotherapy and anti-tumor agents). Generally, administration of
20 products of a species origin or species reactivity (in the case of antibodies) that is the same species as that of the patient is preferred. Thus, in a preferred embodiment, human antibodies, fragments derivatives, analogs, or nucleic acids, are administered to a human patient for therapy or prophylaxis.

It is preferred to use high affinity and/or potent in vivo inhibiting and/or
25 neutralizing antibodies against polypeptides or polynucleotides of the present invention, fragments or regions thereof, for both immunoassays directed to and therapy of disorders related to polynucleotides or polypeptides, including fragments thereof, of the present invention. Such antibodies, fragments, or regions, will preferably have an affinity for polynucleotides or polypeptides of the invention,
30 including fragments thereof. Preferred binding affinities include those with a dissociation constant or K_d less than 5×10^{-2} M, 10^{-2} M, 5×10^{-3} M, 10^{-3} M, 5×10^{-4}

M, 10^{-4} M, 5×10^{-5} M, 10^{-5} M, 5×10^{-6} M, 10^{-6} M, 5×10^{-7} M, 10^{-7} M, 5×10^{-8} M, 10^{-8} M, 5×10^{-9} M, 10^{-9} M, 5×10^{-10} M, 10^{-10} M, 5×10^{-11} M, 10^{-11} M, 5×10^{-12} M, 10^{-12} M, 5×10^{-13} M, 10^{-13} M, 5×10^{-14} M, 10^{-14} M, 5×10^{-15} M, and 10^{-15} M.

5 Gene Therapy

In a specific embodiment, nucleic acids comprising sequences encoding antibodies or functional derivatives thereof, are administered to treat, inhibit or prevent a disease or disorder associated with aberrant expression and/or activity of a polypeptide of the invention, by way of gene therapy. Gene therapy refers to therapy performed by the administration to a subject of an expressed or expressible nucleic acid. In this embodiment of the invention, the nucleic acids produce their encoded protein that mediates a therapeutic effect.

Any of the methods for gene therapy available in the art can be used according to the present invention. Exemplary methods are described below.

15 For general reviews of the methods of gene therapy, see Goldspiel et al., Clinical Pharmacy 12:488-505 (1993); Wu and Wu, Biotherapy 3:87-95 (1991); Tolstoshev, Ann. Rev. Pharmacol. Toxicol. 32:573-596 (1993); Mulligan, Science 260:926-932 (1993); and Morgan and Anderson, Ann. Rev. Biochem. 62:191-217 (1993); May, TIBTECH 11(5):155-215 (1993). Methods commonly known in the art of recombinant DNA technology which can be used are described in Ausubel et al. (eds.), Current Protocols in Molecular Biology, John Wiley & Sons, NY (1993); and Kriegler, Gene Transfer and Expression, A Laboratory Manual, Stockton Press, NY (1990).

25 In a preferred aspect, the compound comprises nucleic acid sequences encoding an antibody, said nucleic acid sequences being part of expression vectors that express the antibody or fragments or chimeric proteins or heavy or light chains thereof in a suitable host. In particular, such nucleic acid sequences have promoters operably linked to the antibody coding region, said promoter being inducible or constitutive, and, optionally, tissue-specific. In another particular embodiment, 30 nucleic acid molecules are used in which the antibody coding sequences and any other desired sequences are flanked by regions that promote homologous recombination at a

desired site in the genome, thus providing for intrachromosomal expression of the antibody encoding nucleic acids (Koller and Smithies, Proc. Natl. Acad. Sci. USA 86:8932-8935 (1989); Zijlstra et al., Nature 342:435-438 (1989). In specific embodiments, the expressed antibody molecule is a single chain antibody;
5 alternatively, the nucleic acid sequences include sequences encoding both the heavy and light chains, or fragments thereof, of the antibody.

Delivery of the nucleic acids into a patient may be either direct, in which case the patient is directly exposed to the nucleic acid or nucleic acid-carrying vectors, or indirect, in which case, cells are first transformed with the nucleic acids in vitro, then
10 transplanted into the patient. These two approaches are known, respectively, as in vivo or ex vivo gene therapy.

In a specific embodiment, the nucleic acid sequences are directly administered in vivo, where it is expressed to produce the encoded product. This can be accomplished by any of numerous methods known in the art, e.g., by constructing
15 them as part of an appropriate nucleic acid expression vector and administering it so that they become intracellular, e.g., by infection using defective or attenuated retrovirals or other viral vectors (see U.S. Patent No. 4,980,286), or by direct injection of naked DNA, or by use of microparticle bombardment (e.g., a gene gun: Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting
20 agents, encapsulation in liposomes, microparticles, or microcapsules, or by administering them in linkage to a peptide which is known to enter the nucleus, by administering it in linkage to a ligand subject to receptor-mediated endocytosis (see, e.g., Wu and Wu, J. Biol. Chem. 262:4429-4432 (1987)) (which can be used to target cell types specifically expressing the receptors), etc. In another embodiment, nucleic
25 acid-ligand complexes can be formed in which the ligand comprises a fusogenic viral peptide to disrupt endosomes, allowing the nucleic acid to avoid lysosomal degradation. In yet another embodiment, the nucleic acid can be targeted in vivo for cell specific uptake and expression, by targeting a specific receptor (see, e.g., PCT Publications WO 92/06180; WO 92/22635; WO92/20316; WO93/14188, WO
30 93/20221). Alternatively, the nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression, by homologous recombination

(Koller and Smithies, *Proc. Natl. Acad. Sci. USA* 86:8932-8935 (1989); Zijlstra et al., *Nature* 342:435-438 (1989)).

In a specific embodiment, viral vectors that contains nucleic acid sequences encoding an antibody of the invention are used. For example, a retroviral vector can be used (see Miller et al., *Meth. Enzymol.* 217:581-599 (1993)). These retroviral vectors contain the components necessary for the correct packaging of the viral genome and integration into the host cell DNA. The nucleic acid sequences encoding the antibody to be used in gene therapy are cloned into one or more vectors, which facilitates delivery of the gene into a patient. More detail about retroviral vectors can be found in Boesen et al., *Biotherapy* 6:291-302 (1994), which describes the use of a retroviral vector to deliver the *mdr1* gene to hematopoietic stem cells in order to make the stem cells more resistant to chemotherapy. Other references illustrating the use of retroviral vectors in gene therapy are: Clowes et al., *J. Clin. Invest.* 93:644-651 (1994); Kiem et al., *Blood* 83:1467-1473 (1994); Salmons and Gunzberg, *Human Gene Therapy* 4:129-141 (1993); and Grossman and Wilson, *Curr. Opin. in Genetics and Devel.* 3:110-114 (1993).

Adenoviruses are other viral vectors that can be used in gene therapy. Adenoviruses are especially attractive vehicles for delivering genes to respiratory epithelia. Adenoviruses naturally infect respiratory epithelia where they cause a mild disease. Other targets for adenovirus-based delivery systems are liver, the central nervous system, endothelial cells, and muscle. Adenoviruses have the advantage of being capable of infecting non-dividing cells. Kozarsky and Wilson, *Current Opinion in Genetics and Development* 3:499-503 (1993) present a review of adenovirus-based gene therapy. Bout et al., *Human Gene Therapy* 5:3-10 (1994) demonstrated the use of adenovirus vectors to transfer genes to the respiratory epithelia of rhesus monkeys. Other instances of the use of adenoviruses in gene therapy can be found in Rosenfeld et al., *Science* 252:431-434 (1991); Rosenfeld et al., *Cell* 68:143-155 (1992); Mastrangeli et al., *J. Clin. Invest.* 91:225-234 (1993); PCT Publication WO94/12649; and Wang, et al., *Gene Therapy* 2:775-783 (1995). In a preferred embodiment, adenovirus vectors are used.

Adeno-associated virus (AAV) has also been proposed for use in gene therapy (Walsh et al., Proc. Soc. Exp. Biol. Med. 204:289-300 (1993); U.S. Patent No. 5,436,146).

Another approach to gene therapy involves transferring a gene to cells in
5 tissue culture by such methods as electroporation, lipofection, calcium phosphate mediated transfection, or viral infection. Usually, the method of transfer includes the transfer of a selectable marker to the cells. The cells are then placed under selection to isolate those cells that have taken up and are expressing the transferred gene. Those cells are then delivered to a patient.

10 In this embodiment, the nucleic acid is introduced into a cell prior to administration in vivo of the resulting recombinant cell. Such introduction can be carried out by any method known in the art, including but not limited to transfection, electroporation, microinjection, infection with a viral or bacteriophage vector containing the nucleic acid sequences, cell fusion, chromosome-mediated gene
15 transfer, microcell-mediated gene transfer, spheroplast fusion, etc. Numerous techniques are known in the art for the introduction of foreign genes into cells (see, e.g., Loeffler and Behr, Meth. Enzymol. 217:599-618 (1993); Cohen et al., Meth. Enzymol. 217:618-644 (1993); Cline, Pharmac. Ther. 29:69-92m (1985) and may be used in accordance with the present invention, provided that the necessary
20 developmental and physiological functions of the recipient cells are not disrupted. The technique should provide for the stable transfer of the nucleic acid to the cell, so that the nucleic acid is expressible by the cell and preferably heritable and expressible by its cell progeny.

The resulting recombinant cells can be delivered to a patient by various
25 methods known in the art. Recombinant blood cells (e.g., hematopoietic stem or progenitor cells) are preferably administered intravenously. The amount of cells envisioned for use depends on the desired effect, patient state, etc., and can be determined by one skilled in the art.

Cells into which a nucleic acid can be introduced for purposes of gene therapy
30 encompass any desired, available cell type, and include but are not limited to epithelial cells, endothelial cells, keratinocytes, fibroblasts, muscle cells, hepatocytes;

blood cells such as Tlymphocytes, Blymphocytes, monocytes, macrophages, neutrophils, eosinophils, megakaryocytes, granulocytes; various stem or progenitor cells, in particular hematopoietic stem or progenitor cells, e.g., as obtained from bone marrow, umbilical cord blood, peripheral blood, fetal liver, etc.

5 In a preferred embodiment, the cell used for gene therapy is autologous to the patient.

In an embodiment in which recombinant cells are used in gene therapy, nucleic acid sequences encoding an antibody are introduced into the cells such that they are expressible by the cells or their progeny, and the recombinant cells are then
10 administered in vivo for therapeutic effect. In a specific embodiment, stem or progenitor cells are used. Any stem and/or progenitor cells which can be isolated and maintained in vitro can potentially be used in accordance with this embodiment of the present invention (see e.g. PCT Publication WO 94/08598; Stemple and Anderson, Cell 71:973-985 (1992); Rheinwald, Meth. Cell Bio. 21A:229 (1980); and
15 Pittelkow and Scott, Mayo Clinic Proc. 61:771 (1986)).

In a specific embodiment, the nucleic acid to be introduced for purposes of gene therapy comprises an inducible promoter operably linked to the coding region, such that expression of the nucleic acid is controllable by controlling the presence or absence of the appropriate inducer of transcription. Demonstration of Therapeutic or
20 Prophylactic Activity

The compounds or pharmaceutical compositions of the invention are preferably tested in vitro, and then in vivo for the desired therapeutic or prophylactic activity, prior to use in humans. For example, in vitro assays to demonstrate the therapeutic or prophylactic utility of a compound or pharmaceutical composition
25 include, the effect of a compound on a cell line or a patient tissue sample. The effect of the compound or composition on the cell line and/or tissue sample can be determined utilizing techniques known to those of skill in the art including, but not limited to, rosette formation assays and cell lysis assays. In accordance with the invention, in vitro assays which can be used to determine whether administration of a
30 specific compound is indicated, include in vitro cell culture assays in which a patient

tissue sample is grown in culture, and exposed to or otherwise administered a compound, and the effect of such compound upon the tissue sample is observed.

Therapeutic/Prophylactic Administration and Composition

5 The invention provides methods of treatment, inhibition and prophylaxis by administration to a subject of an effective amount of a compound or pharmaceutical composition of the invention, preferably an antibody of the invention. In a preferred aspect, the compound is substantially purified (e.g., substantially free from substances that limit its effect or produce undesired side-effects). The subject is
10 preferably an animal, including but not limited to animals such as cows, pigs, horses, chickens, cats, dogs, etc., and is preferably a mammal, and most preferably human.

 Formulations and methods of administration that can be employed when the compound comprises a nucleic acid or an immunoglobulin are described above; additional appropriate formulations and routes of administration can be selected from
15 among those described herein below.

 Various delivery systems are known and can be used to administer a compound of the invention, e.g., encapsulation in liposomes, microparticles, microcapsules, recombinant cells capable of expressing the compound, receptor-mediated endocytosis (see, e.g., Wu and Wu, J. Biol. Chem. 262:4429-4432 (1987)).
20 construction of a nucleic acid as part of a retroviral or other vector, etc. Methods of introduction include but are not limited to intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, and oral routes. The compounds or compositions may be administered by any convenient route, for example by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (e.g.,
25 oral mucosa, rectal and intestinal mucosa, etc.) and may be administered together with other biologically active agents. Administration can be systemic or local. In addition, it may be desirable to introduce the pharmaceutical compounds or compositions of the invention into the central nervous system by any suitable route, including intraventricular and intrathecal injection; intraventricular injection may be
30 facilitated by an intraventricular catheter, for example, attached to a reservoir, such

as an Ommaya reservoir. Pulmonary administration can also be employed, e.g., by use of an inhaler or nebulizer, and formulation with an aerosolizing agent.

In a specific embodiment, it may be desirable to administer the pharmaceutical compounds or compositions of the invention locally to the area in need of treatment; this may be achieved by, for example, and not by way of limitation, local infusion during surgery, topical application, e.g., in conjunction with a wound dressing after surgery, by injection, by means of a catheter, by means of a suppository, or by means of an implant, said implant being of a porous, non-porous, or gelatinous material, including membranes, such as sialastic membranes, or fibers. Preferably, when administering a protein, including an antibody, of the invention, care must be taken to use materials to which the protein does not absorb.

In another embodiment, the compound or composition can be delivered in a vesicle, in particular a liposome (see Langer, *Science* 249:1527-1533 (1990); Treat et al., in *Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 353- 365 (1989); Lopez-Berestein, *ibid.*, pp. 317-327; see generally *ibid.*)

In yet another embodiment, the compound or composition can be delivered in a controlled release system. In one embodiment, a pump may be used (see Langer, *supra*; Sefton, *CRC Crit. Ref. Biomed. Eng.* 14:201 (1987); Buchwald et al., *Surgery* 88:507 (1980); Saudek et al., *N. Engl. J. Med.* 321:574 (1989)). In another embodiment, polymeric materials can be used (see *Medical Applications of Controlled Release*, Langer and Wise (eds.), CRC Pres., Boca Raton, Florida (1974); *Controlled Drug Bioavailability, Drug Product Design and Performance*, Smolen and Ball (eds.), Wiley, New York (1984); Ranger and Peppas, J., *Macromol. Sci. Rev. Macromol. Chem.* 23:61 (1983); see also Levy et al., *Science* 228:190 (1985); During et al., *Ann. Neurol.* 25:351 (1989); Howard et al., *J. Neurosurg.* 71:105 (1989)). In yet another embodiment, a controlled release system can be placed in proximity of the therapeutic target, i.e., the brain, thus requiring only a fraction of the systemic dose (see, e.g., Goodson, in *Medical Applications of Controlled Release*, *supra*, vol. 2, pp. 115-138 (1984)).

Other controlled release systems are discussed in the review by Langer (Science 249:1527-1533 (1990)).

In a specific embodiment where the compound of the invention is a nucleic acid encoding a protein, the nucleic acid can be administered *in vivo* to promote
5 expression of its encoded protein, by constructing it as part of an appropriate nucleic acid expression vector and administering it so that it becomes intracellular, e.g., by use of a retroviral vector (see U.S. Patent No. 4,980,286), or by direct injection, or by use of microparticle bombardment (e.g., a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, or by administering it in
10 linkage to a homeobox- like peptide which is known to enter the nucleus (see e.g., Joliot et al., Proc. Natl. Acad. Sci. USA 88:1864-1868 (1991)), etc. Alternatively, a nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression, by homologous recombination.

The present invention also provides pharmaceutical compositions. Such
15 compositions comprise a therapeutically effective amount of a compound, and a pharmaceutically acceptable carrier. In a specific embodiment, the term "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly in humans. The term
20 "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the therapeutic is administered. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water is a preferred carrier when the pharmaceutical composition is administered intravenously. Saline
25 solutions and aqueous dextrose and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. The composition, if desired,
30 can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. These compositions can take the form of solutions, suspensions, emulsion,

tablets, pills, capsules, powders, sustained-release formulations and the like. The composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides. Oral formulation can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium
5 saccharine, cellulose, magnesium carbonate, etc. Examples of suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E.W. Martin. Such compositions will contain a therapeutically effective amount of the compound, preferably in purified form, together with a suitable amount of carrier so as to provide the form for proper administration to the patient. The formulation
10 should suit the mode of administration.

In a preferred embodiment, the composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous administration to human beings. Typically, compositions for intravenous administration are solutions in sterile isotonic aqueous buffer. Where necessary, the
15 composition may also include a solubilizing agent and a local anesthetic such as lignocaine to ease pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the
20 composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

The compounds of the invention can be formulated as neutral or salt forms.
25 Pharmaceutically acceptable salts include those formed with anions such as those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, etc., and those formed with cations such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, etc.

30 The amount of the compound of the invention which will be effective in the treatment, inhibition and prevention of a disease or disorder associated with aberrant

expression and/or activity of a polypeptide of the invention can be determined by standard clinical techniques. In addition, in vitro assays may optionally be employed to help identify optimal dosage ranges. The precise dose to be employed in the formulation will also depend on the route of administration, and the seriousness of the disease or disorder, and should be decided according to the judgment of the practitioner and each patient's circumstances. Effective doses may be extrapolated from dose-response curves derived from in vitro or animal model test systems.

For antibodies, the dosage administered to a patient is typically 0.1 mg/kg to 100 mg/kg of the patient's body weight. Preferably, the dosage administered to a patient is between 0.1 mg/kg and 20 mg/kg of the patient's body weight, more preferably 1 mg/kg to 10 mg/kg of the patient's body weight. Generally, human antibodies have a longer half-life within the human body than antibodies from other species due to the immune response to the foreign polypeptides. Thus, lower dosages of human antibodies and less frequent administration is often possible. Further, the dosage and frequency of administration of antibodies of the invention may be reduced by enhancing uptake and tissue penetration (e.g., into the brain) of the antibodies by modifications such as, for example, lipidation.

The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration. **Diagnosis and Imaging**

Labeled antibodies, and derivatives and analogs thereof, which specifically bind to a polypeptide of interest can be used for diagnostic purposes to detect, diagnose, or monitor diseases, disorders, and/or conditions associated with the aberrant expression and/or activity of a polypeptide of the invention. The invention provides for the detection of aberrant expression of a polypeptide of interest, comprising (a) assaying the expression of the polypeptide of interest in cells or body fluid of an individual using one or more antibodies specific to the polypeptide interest

and (b) comparing the level of gene expression with a standard gene expression level, whereby an increase or decrease in the assayed polypeptide gene expression level compared to the standard expression level is indicative of aberrant expression.

The invention provides a diagnostic assay for diagnosing a disorder,
5 comprising (a) assaying the expression of the polypeptide of interest in cells or body fluid of an individual using one or more antibodies specific to the polypeptide interest and (b) comparing the level of gene expression with a standard gene expression level, whereby an increase or decrease in the assayed polypeptide gene expression level compared to the standard expression level is indicative of a particular disorder. With
10 respect to cancer, the presence of a relatively high amount of transcript in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier
15 thereby preventing the development or further progression of the cancer.

Antibodies of the invention can be used to assay protein levels in a biological sample using classical immunohistological methods known to those of skill in the art (e.g., see Jalkanen, et al., J. Cell. Biol. 101:976-985 (1985); Jalkanen, et al., J. Cell Biol. 105:3087-3096 (1987)). Other antibody-based methods useful for detecting
20 protein gene expression include immunoassays, such as the enzyme linked immunosorbent assay (ELISA) and the radioimmunoassay (RIA). Suitable antibody assay labels are known in the art and include enzyme labels, such as, glucose oxidase; radioisotopes, such as iodine (^{125}I , ^{121}I), carbon (^{14}C), sulfur (^{35}S), tritium (^3H), indium (^{112}In), and technetium (^{99}Tc); luminescent labels, such as luminol; and
25 fluorescent labels, such as fluorescein and rhodamine, and biotin.

One aspect of the invention is the detection and diagnosis of a disease or disorder associated with aberrant expression of a polypeptide of interest in an animal, preferably a mammal and most preferably a human. In one embodiment, diagnosis comprises: a) administering (for example, parenterally, subcutaneously, or
30 intraperitoneally) to a subject an effective amount of a labeled molecule which specifically binds to the polypeptide of interest; b) waiting for a time interval

following the administering for permitting the labeled molecule to preferentially concentrate at sites in the subject where the polypeptide is expressed (and for unbound labeled molecule to be cleared to background level); c) determining background level; and d) detecting the labeled molecule in the subject, such that
5 detection of labeled molecule above the background level indicates that the subject has a particular disease or disorder associated with aberrant expression of the polypeptide of interest. Background level can be determined by various methods including, comparing the amount of labeled molecule detected to a standard value previously determined for a particular system.

10 It will be understood in the art that the size of the subject and the imaging system used will determine the quantity of imaging moiety needed to produce diagnostic images. In the case of a radioisotope moiety, for a human subject, the quantity of radioactivity injected will normally range from about 5 to 20 millicuries of ^{99m}Tc. The labeled antibody or antibody fragment will then preferentially
15 accumulate at the location of cells which contain the specific protein. In vivo tumor imaging is described in S.W. Burchiel et al., "Immunopharmacokinetics of Radiolabeled Antibodies and Their Fragments." (Chapter 13 in Tumor Imaging: The Radiochemical Detection of Cancer, S.W. Burchiel and B. A. Rhodes, eds., Masson Publishing Inc. (1982).

20 Depending on several variables, including the type of label used and the mode of administration, the time interval following the administration for permitting the labeled molecule to preferentially concentrate at sites in the subject and for unbound labeled molecule to be cleared to background level is 6 to 48 hours or 6 to 24 hours or 6 to 12 hours. In another embodiment the time interval following administration is 5
25 to 20 days or 5 to 10 days.

In an embodiment, monitoring of the disease or disorder is carried out by repeating the method for diagnosing the disease or disease, for example, one month after initial diagnosis, six months after initial diagnosis, one year after initial diagnosis, etc.

30 Presence of the labeled molecule can be detected in the patient using methods known in the art for in vivo scanning. These methods depend upon the type of label

used. Skilled artisans will be able to determine the appropriate method for detecting a particular label. Methods and devices that may be used in the diagnostic methods of the invention include, but are not limited to, computed tomography (CT), whole body scan such as position emission tomography (PET), magnetic resonance imaging
5 (MRI), and sonography.

In a specific embodiment, the molecule is labeled with a radioisotope and is detected in the patient using a radiation responsive surgical instrument (Thurston et al., U.S. Patent No. 5,441,050). In another embodiment, the molecule is labeled with a fluorescent compound and is detected in the patient using a fluorescence responsive
10 scanning instrument. In another embodiment, the molecule is labeled with a positron emitting metal and is detected in the patient using positron emission-tomography. In yet another embodiment, the molecule is labeled with a paramagnetic label and is detected in a patient using magnetic resonance imaging (MRI). Kits

The present invention provides kits that can be used in the above methods. In
15 one embodiment, a kit comprises an antibody of the invention, preferably a purified antibody, in one or more containers. In a specific embodiment, the kits of the present invention contain a substantially isolated polypeptide comprising an epitope which is specifically immunoreactive with an antibody included in the kit. Preferably, the kits of the present invention further comprise a control antibody which does not react with
20 the polypeptide of interest. In another specific embodiment, the kits of the present invention contain a means for detecting the binding of an antibody to a polypeptide of interest (e.g., the antibody may be conjugated to a detectable substrate such as a fluorescent compound, an enzymatic substrate, a radioactive compound or a luminescent compound, or a second antibody which recognizes the first antibody may
25 be conjugated to a detectable substrate).

In another specific embodiment of the present invention, the kit is a diagnostic kit for use in screening serum containing antibodies specific against proliferative and/or cancerous polynucleotides and polypeptides. Such a kit may include a control antibody that does not react with the polypeptide of interest. Such a kit may include a
30 substantially isolated polypeptide antigen comprising an epitope which is specifically immunoreactive with at least one anti-polypeptide antigen antibody. Further, such a

kit includes means for detecting the binding of said antibody to the antigen (e.g., the antibody may be conjugated to a fluorescent compound such as fluorescein or rhodamine which can be detected by flow cytometry). In specific embodiments, the kit may include a recombinantly produced or chemically synthesized polypeptide antigen. The polypeptide antigen of the kit may also be attached to a solid support.

In a more specific embodiment the detecting means of the above-described kit includes a solid support to which said polypeptide antigen is attached. Such a kit may also include a non-attached reporter-labeled anti-human antibody. In this embodiment, binding of the antibody to the polypeptide antigen can be detected by binding of the said reporter-labeled antibody.

In an additional embodiment, the invention includes a diagnostic kit for use in screening serum containing antigens of the polypeptide of the invention. The diagnostic kit includes a substantially isolated antibody specifically immunoreactive with polypeptide or polynucleotide antigens, and means for detecting the binding of the polynucleotide or polypeptide antigen to the antibody. In one embodiment, the antibody is attached to a solid support. In a specific embodiment, the antibody may be a monoclonal antibody. The detecting means of the kit may include a second, labeled monoclonal antibody. Alternatively, or in addition, the detecting means may include a labeled, competing antigen.

In one diagnostic configuration, test serum is reacted with a solid phase reagent having a surface-bound antigen obtained by the methods of the present invention. After binding with specific antigen antibody to the reagent and removing unbound serum components by washing, the reagent is reacted with reporter-labeled anti-human antibody to bind reporter to the reagent in proportion to the amount of bound anti-antigen antibody on the solid support. The reagent is again washed to remove unbound labeled antibody, and the amount of reporter associated with the reagent is determined. Typically, the reporter is an enzyme which is detected by incubating the solid phase in the presence of a suitable fluorometric, luminescent or colorimetric substrate (Sigma, St. Louis, MO).

The solid surface reagent in the above assay is prepared by known techniques for attaching protein material to solid support material, such as polymeric beads, dip

sticks, 96-well plate or filter material. These attachment methods generally include non-specific adsorption of the protein to the support or covalent attachment of the protein, typically through a free amine group, to a chemically reactive group on the solid support, such as an activated carboxyl, hydroxyl, or aldehyde group.

- 5 Alternatively, streptavidin coated plates can be used in conjunction with biotinylated antigen(s).

Thus, the invention provides an assay system or kit for carrying out this diagnostic method. The kit generally includes a support with surface-bound recombinant antigens, and a reporter-labeled anti-human antibody for detecting
10 surface-bound anti-antigen antibody.

Fusion Proteins

Any polypeptide of the present invention can be used to generate fusion
15 proteins. For example, the polypeptide of the present invention, when fused to a second protein, can be used as an antigenic tag. Antibodies raised against the polypeptide of the present invention can be used to indirectly detect the second protein by binding to the polypeptide. Moreover, because secreted proteins target cellular locations based on trafficking signals, the polypeptides of the present
20 invention can be used as targeting molecules once fused to other proteins.

Examples of domains that can be fused to polypeptides of the present invention include not only heterologous signal sequences, but also other heterologous functional regions. The fusion does not necessarily need to be direct, but may occur through linker sequences.

25 Moreover, fusion proteins may also be engineered to improve characteristics of the polypeptide of the present invention. For instance, a region of additional amino acids, particularly charged amino acids, may be added to the N-terminus of the polypeptide to improve stability and persistence during purification from the host cell or subsequent handling and storage. Also, peptide moieties may be added to the
30 polypeptide to facilitate purification. Such regions may be removed prior to final

preparation of the polypeptide. The addition of peptide moieties to facilitate handling of polypeptides are familiar and routine techniques in the art.

Moreover, polypeptides of the present invention, including fragments, and specifically epitopes, can be combined with parts of the constant domain of immunoglobulins (IgA, IgE, IgG, IgM) or portions thereof (CH1, CH2, CH3, and any combination thereof, including both entire domains and portions thereof), resulting in chimeric polypeptides. These fusion proteins facilitate purification and show an increased half-life in vivo. One reported example describes chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins. (EP A 394,827; Traunecker et al., Nature 331:84-86 (1988).) Fusion proteins having disulfide-linked dimeric structures (due to the IgG) can also be more efficient in binding and neutralizing other molecules, than the monomeric secreted protein or protein fragment alone. (Fountoulakis et al., J. Biochem. 15 270:3958-3964 (1995).)

Similarly, EP-A-O 464 533 (Canadian counterpart 2045869) discloses fusion proteins comprising various portions of constant region of immunoglobulin molecules together with another human protein or part thereof. In many cases, the Fc part in a fusion protein is beneficial in therapy and diagnosis, and thus can result in, for example, improved pharmacokinetic properties. (EP-A 0232 262.) Alternatively, deleting the Fc part after the fusion protein has been expressed, detected, and purified, would be desired. For example, the Fc portion may hinder therapy and diagnosis if the fusion protein is used as an antigen for immunizations. In drug discovery, for example, human proteins, such as hIL-5, have been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists of hIL-5. (See, 20 D. Bennett et al., J. Molecular Recognition 8:52-58 (1995); K. Johanson et al., J. Biol. Chem. 270:9459-9471 (1995).) 25

Moreover, the polypeptides of the present invention can be fused to marker sequences, such as a peptide which facilitates purification of the fused polypeptide. In preferred embodiments, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue, 30

Chatsworth, CA, 91311), among others, many of which are commercially available. As described in Gentz et al., Proc. Natl. Acad. Sci. USA 86:821-824 (1989), for instance, hexa-histidine provides for convenient purification of the fusion protein. Another peptide tag useful for purification, the "HA" tag, corresponds to an epitope
5 derived from the influenza hemagglutinin protein. (Wilson et al., Cell 37:767 (1984).)

Thus, any of these above fusions can be engineered using the polynucleotides or the polypeptides of the present invention.

10 **Vectors, Host Cells, and Protein Production**

The present invention also relates to vectors containing the polynucleotide of the present invention, host cells, and the production of polypeptides by recombinant techniques. The vector may be, for example, a phage, plasmid, viral, or retroviral vector. Retroviral vectors may be replication competent or replication defective. In
15 the latter case, viral propagation generally will occur only in complementing host cells.

The polynucleotides may be joined to a vector containing a selectable marker for propagation in a host. Generally, a plasmid vector is introduced in a precipitate, such as a calcium phosphate precipitate, or in a complex with a charged lipid. If the
20 vector is a virus, it may be packaged in vitro using an appropriate packaging cell line and then transduced into host cells.

The polynucleotide insert should be operatively linked to an appropriate promoter, such as the phage lambda PL promoter, the E. coli lac, trp, phoA and tac promoters, the SV40 early and late promoters and promoters of retroviral LTRs, to
25 name a few. Other suitable promoters will be known to the skilled artisan. The expression constructs will further contain sites for transcription initiation, termination, and, in the transcribed region, a ribosome binding site for translation. The coding portion of the transcripts expressed by the constructs will preferably include a translation initiating codon at the beginning and a termination codon (UAA, UGA or
30 UAG) appropriately positioned at the end of the polypeptide to be translated.

As indicated, the expression vectors will preferably include at least one selectable marker. Such markers include dihydrofolate reductase, G418 or neomycin resistance for eukaryotic cell culture and tetracycline, kanamycin or ampicillin resistance genes for culturing in *E. coli* and other bacteria. Representative examples of appropriate hosts include, but are not limited to, bacterial cells, such as *E. coli*, *Streptomyces* and *Salmonella typhimurium* cells; fungal cells, such as yeast cells (e.g., *Saccharomyces cerevisiae* or *Pichia pastoris* (ATCC Accession No. 201178)); insect cells such as *Drosophila* S2 and *Spodoptera Sf9* cells; animal cells such as CHO, COS, 293, and Bowes melanoma cells; and plant cells. Appropriate culture mediums and conditions for the above-described host cells are known in the art.

Among vectors preferred for use in bacteria include pQE70, pQE60 and pQE-9, available from QIAGEN, Inc.; pBluescript vectors, Phagescript vectors, pNH8A, pNH16a, pNH18A, pNH46A, available from Stratagene Cloning Systems, Inc.; and ptrc99a, pKK223-3, pKK233-3, pDR540, pRIT5 available from Pharmacia Biotech, Inc. Among preferred eukaryotic vectors are pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; and pSVK3, pBPV, pMSG and pSVL available from Pharmacia. Preferred expression vectors for use in yeast systems include, but are not limited to pYES2, pYD1, pTEF1/Zeo, pYES2/GS, pPICZ, pGAPZ, pGAPZalph, pPIC9, pPIC3.5, pHIL-D2, pHIL-S1, pPIC3.5K, pPIC9K, and PAO815 (all available from Invitrogen, Carlsbad, CA). Other suitable vectors will be readily apparent to the skilled artisan.

Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection, or other methods. Such methods are described in many standard laboratory manuals, such as Davis et al., Basic Methods In Molecular Biology (1986). It is specifically contemplated that the polypeptides of the present invention may in fact be expressed by a host cell lacking a recombinant vector.

A polypeptide of this invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography,

phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography ("HPLC") is employed for purification.

5 Polypeptides of the present invention, and preferably the secreted form, can also be recovered from: products purified from natural sources, including bodily fluids, tissues and cells, whether directly isolated or cultured; products of chemical synthetic procedures; and products produced by recombinant techniques from a prokaryotic or eukaryotic host, including, for example, bacterial, yeast, higher plant,
10 insect, and mammalian cells. Depending upon the host employed in a recombinant production procedure, the polypeptides of the present invention may be glycosylated or may be non-glycosylated. In addition, polypeptides of the invention may also include an initial modified methionine residue, in some cases as a result of host-mediated processes. Thus, it is well known in the art that the N-terminal methionine
15 encoded by the translation initiation codon generally is removed with high efficiency from any protein after translation in all eukaryotic cells. While the N-terminal methionine on most proteins also is efficiently removed in most prokaryotes, for some proteins, this prokaryotic removal process is inefficient, depending on the nature of the amino acid to which the N-terminal methionine is covalently linked.

20 In one embodiment, the yeast *Pichia pastoris* is used to express the polypeptide of the present invention in a eukaryotic system. *Pichia pastoris* is a methylotrophic yeast which can metabolize methanol as its sole carbon source. A main step in the methanol metabolization pathway is the oxidation of methanol to formaldehyde using O₂. This reaction is catalyzed by the enzyme alcohol oxidase. In
25 order to metabolize methanol as its sole carbon source, *Pichia pastoris* must generate high levels of alcohol oxidase due, in part, to the relatively low affinity of alcohol oxidase for O₂. Consequently, in a growth medium depending on methanol as a main carbon source, the promoter region of one of the two alcohol oxidase genes (*AOX1*) is highly active. In the presence of methanol, alcohol oxidase produced from the *AOX1*
30 gene comprises up to approximately 30% of the total soluble protein in *Pichia*

pastoris. See, Ellis, S.B., *et al.*, *Mol. Cell. Biol.* 5:1111-21 (1985); Koutz, P.J., *et al.*, *Yeast* 5:167-77 (1989); Tschopp, J.F., *et al.*, *Nucl. Acids Res.* 15:3859-76 (1987).

Thus, a heterologous coding sequence, such as, for example, a polynucleotide of the present invention, under the transcriptional regulation of all or part of the *AOX1* regulatory sequence is expressed at exceptionally high levels in *Pichia* yeast grown in the presence of methanol.

In one example, the plasmid vector pPIC9K is used to express DNA encoding a polypeptide of the invention, as set forth herein, in a *Pichea* yeast system essentially as described in "*Pichia* Protocols: Methods in Molecular Biology," D.R. Higgins and J. Cregg, eds. The Humana Press, Totowa, NJ, 1998. This expression vector allows expression and secretion of a protein of the invention by virtue of the strong *AOX1* promoter linked to the *Pichia pastoris* alkaline phosphatase (PHO) secretory signal peptide (i.e., leader) located upstream of a multiple cloning site.

Many other yeast vectors could be used in place of pPIC9K, such as, pYES2, pYD1, pTEF1/Zco, pYES2/GS, pPICZ, pGAPZ, pGAPZalpha, pPIC9, pPIC3.5, pHIL-D2, pHIL-S1, pPIC3.5K, and PAO815, as one skilled in the art would readily appreciate, as long as the proposed expression construct provides appropriately located signals for transcription, translation, secretion (if desired), and the like, including an in-frame AUG as required.

In another embodiment, high-level expression of a heterologous coding sequence, such as, for example, a polynucleotide of the present invention, may be achieved by cloning the heterologous polynucleotide of the invention into an expression vector such as, for example, pGAPZ or pGAPZalpha, and growing the yeast culture in the absence of methanol.

In addition to encompassing host cells containing the vector constructs discussed herein, the invention also encompasses primary, secondary, and immortalized host cells of vertebrate origin, particularly mammalian origin, that have been engineered to delete or replace endogenous genetic material (e.g., coding sequence), and/or to include genetic material (e.g., heterologous polynucleotide sequences) that is operably associated with the polynucleotides of the invention, and

which activates, alters, and/or amplifies endogenous polynucleotides. For example, techniques known in the art may be used to operably associate heterologous control regions (c.g., promoter and/or enhancer) and endogenous polynucleotide sequences via homologous recombination, resulting in the formation of a new transcription unit
5 (see, e.g., U.S. Patent No. 5,641,670, issued June 24, 1997; U.S. Patent No. 5,733,761, issued March 31, 1998; International Publication No. WO 96/29411, published September 26, 1996; International Publication No. WO 94/12650, published August 4, 1994; Koller et al., Proc. Natl. Acad. Sci. USA 86:8932-8935 (1989); and Zijlstra et al., Nature 342:435-438 (1989), the disclosures of each of
10 which are incorporated by reference in their entireties).

In addition, polypeptides of the invention can be chemically synthesized using techniques known in the art (e.g., see Creighton, 1983, Proteins: Structures and Molecular Principles, W.H. Freeman & Co., N.Y., and Hunkapiller et al., *Nature*, 310:105-111 (1984)). For example, a polypeptide corresponding to a fragment of a
15 polypeptide sequence of the invention can be synthesized by use of a peptide synthesizer. Furthermore, if desired, nonclassical amino acids or chemical amino acid analogs can be introduced as a substitution or addition into the polypeptide sequence. Non-classical amino acids include, but are not limited to, to the D-isomers of the common amino acids, 2,4-diaminobutyric acid, α -amino isobutyric acid, 4-
20 aminobutyric acid, Abu, 2-amino butyric acid, γ -Abu, ϵ -Ahx, 6-amino hexanoic acid, Aib, 2-amino isobutyric acid, 3-amino propionic acid, ornithine, norleucine, norvaline, hydroxyproline, sarcosine, citrulline, homocitrulline, cysteic acid, t-butylglycine, t-butylalanine, phenylglycine, cyclohexylalanine, β -alanine, fluoro-amino acids, designer amino acids such as β -methyl amino acids, Ca-methyl amino-
25 acids, Na-methyl amino acids, and amino acid analogs in general. Furthermore, the amino acid can be D (dextrorotary) or L (levorotary).

The invention encompasses polypeptides which are differentially modified during or after translation, e.g., by glycosylation, acetylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage,
30 linkage to an antibody molecule or other cellular ligand, etc. Any of numerous chemical modifications may be carried out by known techniques, including but not

limited, to specific chemical cleavage by cyanogen bromide, trypsin, chymotrypsin, papain, V8 protease, NaBH_4 ; acetylation, formylation, oxidation, reduction; metabolic synthesis in the presence of tunicamycin; etc.

Additional post-translational modifications encompassed by the invention include, for example, e.g., N-linked or O-linked carbohydrate chains, processing of N-terminal or C-terminal ends), attachment of chemical moieties to the amino acid backbone, chemical modifications of N-linked or O-linked carbohydrate chains, and addition or deletion of an N-terminal methionine residue as a result of procaryotic host cell expression. The polypeptides may also be modified with a detectable label, such as an enzymatic, fluorescent, isotopic or affinity label to allow for detection and isolation of the protein.

Also provided by the invention are chemically modified derivatives of the polypeptides of the invention which may provide additional advantages such as increased solubility, stability and circulating time of the polypeptide, or decreased immunogenicity (see U.S. Patent NO: 4,179,337). The chemical moieties for derivitization may be selected from water soluble polymers such as polyethylene glycol, ethylene glycol/propylene glycol copolymers, carboxymethylcellulose, dextran, polyvinyl alcohol and the like. The polypeptides may be modified at random positions within the molecule, or at predetermined positions within the molecule and may include one, two, three or more attached chemical moieties.

The polymer may be of any molecular weight, and may be branched or unbranched. For polyethylene glycol, the preferred molecular weight is between about 1 kDa and about 100 kDa (the term "about" indicating that in preparations of polyethylene glycol, some molecules will weigh more, some less, than the stated molecular weight) for ease in handling and manufacturing. Other sizes may be used, depending on the desired therapeutic profile (e.g., the duration of sustained release desired, the effects, if any on biological activity, the ease in handling, the degree or lack of antigenicity and other known effects of the polyethylene glycol to a therapeutic protein or analog).

The polyethylene glycol molecules (or other chemical moieties) should be attached to the protein with consideration of effects on functional or antigenic

domains of the protein. There are a number of attachment methods available to those skilled in the art, e.g., EP 0 401 384, herein incorporated by reference (coupling PEG to G-CSF), see also Malik et al., Exp. Hematol. 20:1028-1035 (1992) (reporting pegylation of GM-CSF using tresyl chloride). For example, polyethylene glycol may be covalently bound through amino acid residues via a reactive group, such as, a free amino or carboxyl group. Reactive groups are those to which an activated polyethylene glycol molecule may be bound. The amino acid residues having a free amino group may include lysine residues and the N-terminal amino acid residues; those having a free carboxyl group may include aspartic acid residues glutamic acid residues and the C-terminal amino acid residue. Sulfhydryl groups may also be used as a reactive group for attaching the polyethylene glycol molecules. Preferred for therapeutic purposes is attachment at an amino group, such as attachment at the N-terminus or lysine group.

One may specifically desire proteins chemically modified at the N-terminus. Using polyethylene glycol as an illustration of the present composition, one may select from a variety of polyethylene glycol molecules (by molecular weight, branching, etc.), the proportion of polyethylene glycol molecules to protein (polypeptide) molecules in the reaction mix, the type of pegylation reaction to be performed, and the method of obtaining the selected N-terminally pegylated protein. The method of obtaining the N-terminally pegylated preparation (i.e., separating this moiety from other monopegylated moieties if necessary) may be by purification of the N-terminally pegylated material from a population of pegylated protein molecules. Selective proteins chemically modified at the N-terminus modification may be accomplished by reductive alkylation which exploits differential reactivity of different types of primary amino groups (lysine versus the N-terminal) available for derivatization in a particular protein. Under the appropriate reaction conditions, substantially selective derivatization of the protein at the N-terminus with a carbonyl group containing polymer is achieved.

The polypeptides of the invention may be in monomers or multimers (i.e., dimers, trimers, tetramers and higher multimers). Accordingly, the present invention relates to monomers and multimers of the polypeptides of the invention, their

preparation, and compositions (preferably, *Therapeutics*) containing them. In specific embodiments, the polypeptides of the invention are monomers, dimers, trimers or tetramers. In additional embodiments, the multimers of the invention are at least dimers, at least trimers, or at least tetramers.

5 Multimers encompassed by the invention may be homomers or heteromers. As used herein, the term homomer, refers to a multimer containing only polypeptides corresponding to the amino acid sequence of SEQ ID NO:Y or encoded by the cDNA contained in a deposited clone (including fragments, variants, splice variants, and fusion proteins, corresponding to these polypeptides as described herein). These
10 homomers may contain polypeptides having identical or different amino acid sequences. In a specific embodiment, a homomer of the invention is a multimer containing only polypeptides having an identical amino acid sequence. In another specific embodiment, a homomer of the invention is a multimer containing polypeptides having different amino acid sequences. In specific embodiments, the
15 multimer of the invention is a homodimer (*e.g.*, containing polypeptides having identical or different amino acid sequences) or a homotrimer (*e.g.*, containing polypeptides having identical and/or different amino acid sequences). In additional embodiments, the homomeric multimer of the invention is at least a homodimer, at least a homotrimer, or at least a homotetramer.

20 As used herein, the term heteromer refers to a multimer containing one or more heterologous polypeptides (*i.e.*, polypeptides of different proteins) in addition to the polypeptides of the invention. In a specific embodiment, the multimer of the invention is a heterodimer, a heterotrimer, or a heterotetramer. In additional
25 embodiments, the heteromeric multimer of the invention is at least a heterodimer, at least a heterotrimer, or at least a heterotetramer.

 Multimers of the invention may be the result of hydrophobic, hydrophilic, ionic and/or covalent associations and/or may be indirectly linked, by for example, liposome formation. Thus, in one embodiment, multimers of the invention, such as, for example, homodimers or homotrimers, are formed when polypeptides of the
30 invention contact one another in solution. In another embodiment, heteromultimers of the invention, such as, for example, heterotrimers or heterotetramers, are formed

when polypeptides of the invention contact antibodies to the polypeptides of the invention (including antibodies to the heterologous polypeptide sequence in a fusion protein of the invention) in solution. In other embodiments, multimers of the invention are formed by covalent associations with and/or between the polypeptides
5 of the invention. Such covalent associations may involve one or more amino acid residues contained in the polypeptide sequence (e.g., that recited in the sequence listing, or contained in the polypeptide encoded by a deposited clone). In one instance, the covalent associations are cross-linking between cysteine residues located within the polypeptide sequences which interact in the native (i.e., naturally
10 occurring) polypeptide. In another instance, the covalent associations are the consequence of chemical or recombinant manipulation. Alternatively, such covalent associations may involve one or more amino acid residues contained in the heterologous polypeptide sequence in a fusion protein of the invention.

In one example, covalent associations are between the heterologous sequence
15 contained in a fusion protein of the invention (see, e.g., US Patent Number 5,478,925). In a specific example, the covalent associations are between the heterologous sequence contained in an Fc fusion protein of the invention (as described herein). In another specific example, covalent associations of fusion proteins of the invention are between heterologous polypeptide sequence from
20 another protein that is capable of forming covalently associated multimers, such as for example, osteopontin (see, e.g., International Publication NO: WO 98/49305, the contents of which are herein incorporated by reference in its entirety). In another embodiment, two or more polypeptides of the invention are joined through peptide linkers. Examples include those peptide linkers described in U.S. Pat. No. 5,073,627
25 (hereby incorporated by reference). Proteins comprising multiple polypeptides of the invention separated by peptide linkers may be produced using conventional recombinant DNA technology.

Another method for preparing multimer polypeptides of the invention involves use of polypeptides of the invention fused to a leucine zipper or isoleucine zipper
30 polypeptide sequence. Leucine zipper and isoleucine zipper domains are polypeptides that promote multimerization of the proteins in which they are found. Leucine

zippers were originally identified in several DNA-binding proteins (Landschulz et al., Science 240:1759, (1988)), and have since been found in a variety of different proteins. Among the known leucine zippers are naturally occurring peptides and derivatives thereof that dimerize or trimerize. Examples of leucine zipper domains
5 suitable for producing soluble multimeric proteins of the invention are those described in PCT application WO 94/10308, hereby incorporated by reference. Recombinant fusion proteins comprising a polypeptide of the invention fused to a polypeptide sequence that dimerizes or trimerizes in solution are expressed in suitable host cells, and the resulting soluble multimeric fusion protein is recovered from the culture
10 supernatant using techniques known in the art.

Trimeric polypeptides of the invention may offer the advantage of enhanced biological activity. Preferred leucine zipper moieties and isoleucine moieties are those that preferentially form trimers. One example is a leucine zipper derived from lung surfactant protein D (SPD), as described in Hoppe et al. (FEBS Letters 344:191,
15 (1994)) and in U.S. patent application Ser. No. 08/446,922, hereby incorporated by reference. Other peptides derived from naturally occurring trimeric proteins may be employed in preparing trimeric polypeptides of the invention.

In another example, proteins of the invention are associated by interactions between Flag® polypeptide sequence contained in fusion proteins of the invention
20 containing Flag® polypeptide sequence. In a further embodiment, associations proteins of the invention are associated by interactions between heterologous polypeptide sequence contained in Flag® fusion proteins of the invention and anti-Flag® antibody.

The multimers of the invention may be generated using chemical techniques
25 known in the art. For example, polypeptides desired to be contained in the multimers of the invention may be chemically cross-linked using linker molecules and linker molecule length optimization techniques known in the art (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). Additionally, multimers of the invention may be generated using techniques known in
30 the art to form one or more inter-molecule cross-links between the cysteine residues located within the sequence of the polypeptides desired to be contained in the

multimer (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). Further, polypeptides of the invention may be routinely modified by the addition of cysteine or biotin to the C terminus or N-terminus of the polypeptide and techniques known in the art may be applied to generate multimers
5 containing one or more of these modified polypeptides (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). Additionally, techniques known in the art may be applied to generate liposomes containing the polypeptide components desired to be contained in the multimer of the invention (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its
10 entirety).

Alternatively, multimers of the invention may be generated using genetic engineering techniques known in the art. In one embodiment, polypeptides contained in multimers of the invention are produced recombinantly using fusion protein technology described herein or otherwise known in the art (see, e.g., US Patent
15 Number 5,478,925, which is herein incorporated by reference in its entirety). In a specific embodiment, polynucleotides coding for a homodimer of the invention are generated by ligating a polynucleotide sequence encoding a polypeptide of the invention to a sequence encoding a linker polypeptide and then further to a synthetic polynucleotide encoding the translated product of the polypeptide in the reverse
20 orientation from the original C-terminus to the N-terminus (lacking the leader sequence) (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). In another embodiment, recombinant techniques described herein or otherwise known in the art are applied to generate recombinant polypeptides of the invention which contain a transmembrane domain (or hydrophobic or signal
25 peptide) and which can be incorporated by membrane reconstitution techniques into liposomes (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety).

Uses of the Polynucleotides

Each of the polynucleotides identified herein can be used in numerous ways as reagents. The following description should be considered exemplary and utilizes known techniques.

The polynucleotides of the present invention are useful for chromosome
5 identification. There exists an ongoing need to identify new chromosome markers, since few chromosome marking reagents, based on actual sequence data (repeat polymorphisms), are presently available. Each polynucleotide of the present invention can be used as a chromosome marker.

Briefly, sequences can be mapped to chromosomes by preparing PCR primers
10 (preferably 15-25 bp) from the sequences shown in SEQ ID NO:X. Primers can be selected using computer analysis so that primers do not span more than one predicted exon in the genomic DNA. These primers are then used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to the SEQ ID NO:X will yield an
15 amplified fragment.

Similarly, somatic hybrids provide a rapid method of PCR mapping the polynucleotides to particular chromosomes. Three or more clones can be assigned per day using a single thermal cycler. Moreover, sublocalization of the polynucleotides can be achieved with panels of specific chromosome fragments. Other gene mapping
20 strategies that can be used include in situ hybridization, prescreening with labeled flow-sorted chromosomes, and preselection by hybridization to construct chromosome specific-cDNA libraries.

Precise chromosomal location of the polynucleotides can also be achieved using fluorescence in situ hybridization (FISH) of a metaphase chromosomal spread.
25 This technique uses polynucleotides as short as 500 or 600 bases; however, polynucleotides 2,000-4,000 bp are preferred. For a review of this technique, see Verma et al., "Human Chromosomes: a Manual of Basic Techniques," Pergamon Press, New York (1988).

For chromosome mapping, the polynucleotides can be used individually (to
30 mark a single chromosome or a single site on that chromosome) or in panels (for marking multiple sites and/or multiple chromosomes). Preferred polynucleotides

correspond to the noncoding regions of the cDNAs because the coding sequences are more likely conserved within gene families, thus increasing the chance of cross hybridization during chromosomal mapping.

Once a polynucleotide has been mapped to a precise chromosomal location,
5 the physical position of the polynucleotide can be used in linkage analysis. Linkage analysis establishes coinheritance between a chromosomal location and presentation of a particular disease. (Disease mapping data are found, for example, in V. McKusick, Mendelian Inheritance in Man (available on line through Johns Hopkins University Welch Medical Library) .) Assuming 1 megabase mapping resolution and
10 one gene per 20 kb, a cDNA precisely localized to a chromosomal region associated with the disease could be one of 50-500 potential causative genes.

Thus, once coinheritance is established, differences in the polynucleotide and the corresponding gene between affected and unaffected individuals can be examined. First, visible structural alterations in the chromosomes, such as deletions or
15 translocations, are examined in chromosome spreads or by PCR. If no structural alterations exist, the presence of point mutations are ascertained. Mutations observed in some or all affected individuals, but not in normal individuals, indicates that the mutation may cause the disease. However, complete sequencing of the polypeptide and the corresponding gene from several normal individuals is required to distinguish
20 the mutation from a polymorphism. If a new polymorphism is identified, this polymorphic polypeptide can be used for further linkage analysis.

Furthermore, increased or decreased expression of the gene in affected individuals as compared to unaffected individuals can be assessed using polynucleotides of the present invention. Any of these alterations (altered expression,
25 chromosomal rearrangement, or mutation) can be used as a diagnostic or prognostic marker.

Thus, the invention also provides a diagnostic method useful during diagnosis of a disorder, involving measuring the expression level of polynucleotides of the present invention in cells or body fluid from an individual and comparing the
30 measured gene expression level with a standard level of polynucleotide expression

level, whereby an increase or decrease in the gene expression level compared to the standard is indicative of a disorder.

In still another embodiment, the invention includes a kit for analyzing samples for the presence of proliferative and/or cancerous polynucleotides derived from a test
5 subject. In a general embodiment, the kit includes at least one polynucleotide probe containing a nucleotide sequence that will specifically hybridize with a polynucleotide of the present invention and a suitable container. In a specific embodiment, the kit includes two polynucleotide probes defining an internal region of the polynucleotide of the present invention, where each probe has one strand
10 containing a 31' mer-end internal to the region. In a further embodiment, the probes may be useful as primers for polymerase chain reaction amplification.

Where a diagnosis of a disorder, has already been made according to conventional methods, the present invention is useful as a prognostic indicator, whereby patients exhibiting enhanced or depressed polynucleotide of the present
15 invention expression will experience a worse clinical outcome relative to patients expressing the gene at a level nearer the standard level.

By "measuring the expression level of polynucleotide of the present invention" is intended qualitatively or quantitatively measuring or estimating the level of the polypeptide of the present invention or the level of the mRNA encoding the
20 polypeptide in a first biological sample either directly (e.g., by determining or estimating absolute protein level or mRNA level) or relatively (e.g., by comparing to the polypeptide level or mRNA level in a second biological sample). Preferably, the polypeptide level or mRNA level in the first biological sample is measured or estimated and compared to a standard polypeptide level or mRNA level, the standard
25 being taken from a second biological sample obtained from an individual not having the disorder or being determined by averaging levels from a population of individuals not having a disorder. As will be appreciated in the art, once a standard polypeptide level or mRNA level is known, it can be used repeatedly as a standard for comparison.

30 By "biological sample" is intended any biological sample obtained from an individual, body fluid, cell line, tissue culture, or other source which contains the

polypeptide of the present invention or mRNA. As indicated, biological samples include body fluids (such as semen, lymph, sera, plasma, urine, synovial fluid and spinal fluid) which contain the polypeptide of the present invention, and other tissue sources found to express the polypeptide of the present invention. Methods for
5 obtaining tissue biopsies and body fluids from mammals are well known in the art. Where the biological sample is to include mRNA, a tissue biopsy is the preferred source.

The method(s) provided above may preferably be applied in a diagnostic method and/or kits in which polynucleotides and/or polypeptides are attached to a
10 solid support. In one exemplary method, the support may be a "gene chip" or a "biological chip" as described in US Patents 5,837,832, 5,874,219, and 5,856,174. Further, such a gene chip with polynucleotides of the present invention attached may be used to identify polymorphisms between the polynucleotide sequences, with polynucleotides isolated from a test subject. The knowledge of such polymorphisms
15 (i.e. their location, as well as, their existence) would be beneficial in identifying disease loci for many disorders, including cancerous diseases and conditions. Such a method is described in US Patents 5,858,659 and 5,856,104. The US Patents referenced supra are hereby incorporated by reference in their entirety herein.

The present invention encompasses polynucleotides of the present invention
20 that are chemically synthesized, or reproduced as peptide nucleic acids (PNA), or according to other methods known in the art. The use of PNAs would serve as the preferred form if the polynucleotides are incorporated onto a solid support, or gene chip. For the purposes of the present invention, a peptide nucleic acid (PNA) is a polyamide type of DNA analog and the monomeric units for adenine, guanine,
25 thymine and cytosine are available commercially (Perceptive Biosystems). Certain components of DNA, such as phosphorus, phosphorus oxides, or deoxyribose derivatives, are not present in PNAs. As disclosed by P. E. Nielsen, M. Egholm, R. H. Berg and O. Buchardt, *Science* 254, 1497 (1991); and M. Egholm, O. Buchardt, L. Christensen, C. Behrens, S. M. Freier, D. A. Driver, R. H. Berg, S. K. Kim, B. Norden, and P. E. Nielsen, *Nature* 365, 666 (1993), PNAs bind specifically and
30 tightly to complementary DNA strands and are not degraded by nucleases. In fact,

PNA binds more strongly to DNA than DNA itself does. This is probably because there is no electrostatic repulsion between the two strands, and also the polyamide backbone is more flexible. Because of this, PNA/DNA duplexes bind under a wider range of stringency conditions than DNA/DNA duplexes, making it easier to perform
5 multiplex hybridization. Smaller probes can be used than with DNA due to the strong binding. In addition, it is more likely that single base mismatches can be determined with PNA/DNA hybridization because a single mismatch in a PNA/DNA 15-mer lowers the melting point ($T_{sub.m}$) by 8°-20° C, vs. 4°-16° C for the DNA/DNA 15-mer duplex. Also, the absence of charge groups in PNA means that hybridization can
10 be done at low ionic strengths and reduce possible interference by salt during the analysis.

The present invention is useful for detecting cancer in mammals. In particular the invention is useful during diagnosis of pathological cell proliferative neoplasias which include, but are not limited to: acute myelogenous leukemias including acute
15 monocytic leukemia, acute myeloblastic leukemia, acute promyelocytic leukemia, acute myelomonocytic leukemia, acute erythroleukemia, acute megakaryocytic leukemia, and acute undifferentiated leukemia, etc.; and chronic myelogenous leukemias including chronic myelomonocytic leukemia, chronic granulocytic leukemia, etc. Preferred mammals include monkeys, apes, cats, dogs, cows, pigs,
20 horses, rabbits and humans. Particularly preferred are humans.

Pathological cell proliferative diseases, disorders, and/or conditions are often associated with inappropriate activation of proto-oncogenes. (Germann, E. P. et al., "The Etiology of Acute Leukemia: Molecular Genetics and Viral Oncology," in Neoplastic Diseases of the Blood, Vol 1., Wiernik, P. H. et al. eds., 161-182 (1985)).
25 Neoplasias are now believed to result from the qualitative alteration of a normal cellular gene product, or from the quantitative modification of gene expression by insertion into the chromosome of a viral sequence, by chromosomal translocation of a gene to a more actively transcribed region, or by some other mechanism. (Germann et al., supra) It is likely that mutated or altered expression of specific genes is
30 involved in the pathogenesis of some leukemias, among other tissues and cell types. (Germann et al., supra) Indeed, the human counterparts of the oncogenes involved in

some animal neoplasias have been amplified or translocated in some cases of human leukemia and carcinoma. (Germann et al., supra)

For example, c-myc expression is highly amplified in the non-lymphocytic leukemia cell line HL-60. When HL-60 cells are chemically induced to stop proliferation, the
5 level of c-myc is found to be downregulated. (International Publication Number WO 91/15580) However, it has been shown that exposure of HL-60 cells to a DNA construct that is complementary to the 5' end of c-myc or c-myb blocks translation of the corresponding mRNAs which downregulates expression of the c-myc or c-myb proteins and causes arrest of cell proliferation and differentiation of the treated cells.
10 (International Publication Number WO 91/15580; Wickstrom et al., Proc. Natl. Acad. Sci. 85:1028 (1988); Anfossi et al., Proc. Natl. Acad. Sci. 86:3379 (1989)). However, the skilled artisan would appreciate the present invention's usefulness would not be limited to treatment of proliferative diseases, disorders, and/or conditions of hematopoietic cells and tissues, in light of the numerous cells and cell types of
15 varying origins which are known to exhibit proliferative phenotypes.

In addition to the foregoing, a polynucleotide can be used to control gene expression through triple helix formation or antisense DNA or RNA. Antisense techniques are discussed, for example, in Okano, J. Neurochem. 56: 560 (1991); "Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca
20 Raton, FL (1988). Triple helix formation is discussed in, for instance Lee et al., Nucleic Acids Research 6: 3073 (1979); Cooney et al., Science 241: 456 (1988); and Dervan et al., Science 251: 1360 (1991). Both methods rely on binding of the polynucleotide to a complementary DNA or RNA. For these techniques, preferred polynucleotides are usually oligonucleotides 20 to 40 bases in length and
25 complementary to either the region of the gene involved in transcription (triple helix - see Lee et al., Nucl. Acids Res. 6:3073 (1979); Cooney et al., Science 241:456 (1988); and Dervan et al., Science 251:1360 (1991)) or to the mRNA itself (antisense - Okano, J. Neurochem. 56:560 (1991); Oligodeoxy-nucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988).) Triple helix
30 formation optimally results in a shut-off of RNA transcription from DNA, while antisense RNA hybridization blocks translation of an mRNA molecule into

polypeptide. Both techniques are effective in model systems, and the information disclosed herein can be used to design antisense or triple helix polynucleotides in an effort to treat or prevent disease.

Polynucleotides of the present invention are also useful in gene therapy. One
5 goal of gene therapy is to insert a normal gene into an organism having a defective gene, in an effort to correct the genetic defect. The polynucleotides disclosed in the present invention offer a means of targeting such genetic defects in a highly accurate manner. Another goal is to insert a new gene that was not present in the host genome, thereby producing a new trait in the host cell.

10 The polynucleotides are also useful for identifying individuals from minute biological samples. The United States military, for example, is considering the use of restriction fragment length polymorphism (RFLP) for identification of its personnel. In this technique, an individual's genomic DNA is digested with one or more restriction enzymes, and probed on a Southern blot to yield unique bands for
15 identifying personnel. This method does not suffer from the current limitations of "Dog Tags" which can be lost, switched, or stolen, making positive identification difficult. The polynucleotides of the present invention can be used as additional DNA markers for RFLP.

The polynucleotides of the present invention can also be used as an alternative
20 to RFLP, by determining the actual base-by-base DNA sequence of selected portions of an individual's genome. These sequences can be used to prepare PCR primers for amplifying and isolating such selected DNA, which can then be sequenced. Using this technique, individuals can be identified because each individual will have a unique set of DNA sequences. Once an unique ID database is established for an
25 individual, positive identification of that individual, living or dead, can be made from extremely small tissue samples.

Forensic biology also benefits from using DNA-based identification
techniques as disclosed herein. DNA sequences taken from very small biological
samples such as tissues, e.g., hair or skin, or body fluids, e.g., blood, saliva, semen,
30 synovial fluid, amniotic fluid, breast milk, lymph, pulmonary sputum or surfactant, urine, fecal matter, etc., can be amplified using PCR. In one prior art

technique, gene sequences amplified from polymorphic loci, such as DQa class II HLA gene, are used in forensic biology to identify individuals. (Erlich, H., PCR Technology, Freeman and Co. (1992).) Once these specific polymorphic loci are amplified, they are digested with one or more restriction enzymes, yielding an
5 identifying set of bands on a Southern blot probed with DNA corresponding to the DQa class II HLA gene. Similarly, polynucleotides of the present invention can be used as polymorphic markers for forensic purposes.

There is also a need for reagents capable of identifying the source of a particular tissue. Such need arises, for example, in forensics when presented with
10 tissue of unknown origin. Appropriate reagents can comprise, for example, DNA probes or primers specific to particular tissue prepared from the sequences of the present invention. Panels of such reagents can identify tissue by species and/or by organ type. In a similar fashion, these reagents can be used to screen tissue cultures for contamination.

15 In the very least, the polynucleotides of the present invention can be used as molecular weight markers on Southern gels, as diagnostic probes for the presence of a specific mRNA in a particular cell type, as a probe to "subtract-out" known sequences in the process of discovering novel polynucleotides, for selecting and making oligomers for attachment to a "gene chip" or other support, to raise anti-DNA
20 antibodies using DNA immunization techniques, and as an antigen to elicit an immune response.

Uses of the Polypeptides

Each of the polypeptides identified herein can be used in numerous ways. The
25 following description should be considered exemplary and utilizes known techniques.

A polypeptide of the present invention can be used to assay protein levels in a biological sample using antibody-based techniques. For example, protein expression in tissues can be studied with classical immunohistological methods. (Jalkanen, M., et al., J. Cell. Biol. 101:976-985 (1985); Jalkanen, M., et al., J. Cell . Biol. 105:3087-
30 3096 (1987).) Other antibody-based methods useful for detecting protein gene expression include immunoassays, such as the enzyme linked immunosorbent assay

(ELISA) and the radioimmunoassay (RIA). Suitable antibody assay labels are known in the art and include enzyme labels, such as, glucose oxidase, and radioisotopes, such as iodine (^{125}I , ^{121}I), carbon (^{14}C), sulfur (^{35}S), tritium (^3H), indium (^{112}In), and technetium ($^{99\text{m}}\text{Tc}$), and fluorescent labels, such as fluorescein and rhodamine, and
5 biotin.

In addition to assaying secreted protein levels in a biological sample, proteins can also be detected in vivo by imaging. Antibody labels or markers for in vivo imaging of protein include those detectable by X-radiography, NMR or ESR. For X-radiography, suitable labels include radioisotopes such as barium or cesium, which
10 emit detectable radiation but are not overtly harmful to the subject. Suitable markers for NMR and ESR include those with a detectable characteristic spin, such as deuterium, which may be incorporated into the antibody by labeling of nutrients for the relevant hybridoma.

A protein-specific antibody or antibody fragment which has been labeled with
15 an appropriate detectable imaging moiety, such as a radioisotope (for example, ^{131}I , ^{112}In , $^{99\text{m}}\text{Tc}$), a radio-opaque substance, or a material detectable by nuclear magnetic resonance, is introduced (for example, parenterally, subcutaneously, or intraperitoneally) into the mammal. It will be understood in the art that the size of the subject and the imaging system used will determine the quantity of imaging moiety
20 needed to produce diagnostic images. In the case of a radioisotope moiety, for a human subject, the quantity of radioactivity injected will normally range from about 5 to 20 millicuries of $^{99\text{m}}\text{Tc}$. The labeled antibody or antibody fragment will then preferentially accumulate at the location of cells which contain the specific protein. In vivo tumor imaging is described in S.W. Burchiel et al., "Immunopharmacokinetics
25 of Radiolabeled Antibodies and Their Fragments." (Chapter 13 in Tumor Imaging: The Radiochemical Detection of Cancer, S.W. Burchiel and B. A. Rhodes, eds., Masson Publishing Inc. (1982).)

Thus, the invention provides a diagnostic method of a disorder, which involves (a) assaying the expression of a polypeptide of the present invention in cells
30 or body fluid of an individual; (b) comparing the level of gene expression with a standard gene expression level, whereby an increase or decrease in the assayed

polypeptide gene expression level compared to the standard expression level is indicative of a disorder. With respect to cancer, the presence of a relatively high amount of transcript in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for
5 detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Moreover, polypeptides of the present invention can be used to treat, prevent,
10 and/or diagnose disease. For example, patients can be administered a polypeptide of the present invention in an effort to replace absent or decreased levels of the polypeptide (e.g., insulin), to supplement absent or decreased levels of a different polypeptide (e.g., hemoglobin S for hemoglobin B, SOD, catalase, DNA repair proteins), to inhibit the activity of a polypeptide (e.g., an oncogene or tumor
15 suppressor), to activate the activity of a polypeptide (e.g., by binding to a receptor), to reduce the activity of a membrane bound receptor by competing with it for free ligand (e.g., soluble TNF receptors used in reducing inflammation), or to bring about a desired response (e.g., blood vessel growth inhibition, enhancement of the immune response to proliferative cells or tissues).

20 Similarly, antibodies directed to a polypeptide of the present invention can also be used to treat, prevent, and/or diagnose disease. For example, administration of an antibody directed to a polypeptide of the present invention can bind and reduce overproduction of the polypeptide. Similarly, administration of an antibody can activate the polypeptide, such as by binding to a polypeptide bound to a membrane
25 (receptor).

At the very least, the polypeptides of the present invention can be used as molecular weight markers on SDS-PAGE gels or on molecular sieve gel filtration columns using methods well known to those of skill in the art. Polypeptides can also be used to raise antibodies, which in turn are used to measure protein expression from
30 a recombinant cell, as a way of assessing transformation of the host cell. Moreover,

the polypeptides of the present invention can be used to test the following biological activities.

Gene Therapy Methods

5 Another aspect of the present invention is to gene therapy methods for treating or preventing disorders, diseases and conditions. The gene therapy methods relate to the introduction of nucleic acid (DNA, RNA and antisense DNA or RNA) sequences into an animal to achieve expression of a polypeptide of the present invention. This method requires a polynucleotide which codes for a polypeptide of the invention that
10 operatively linked to a promoter and any other genetic elements necessary for the expression of the polypeptide by the target tissue. Such gene therapy and delivery techniques are known in the art, see, for example, WO90/11092, which is herein incorporated by reference.

Thus, for example, cells from a patient may be engineered with a
15 polynucleotide (DNA or RNA) comprising a promoter operably linked to a polynucleotide of the invention *ex vivo*, with the engineered cells then being provided to a patient to be treated with the polypeptide. Such methods are well-known in the art. For example, see Belldgrun et al., J. Natl. Cancer Inst., 85:207-216 (1993); Ferrantini et al., Cancer Research, 53:107-1112 (1993); Ferrantini et al., J.
20 Immunology 153: 4604-4615 (1994); Kaido, T., et al., Int. J. Cancer 60: 221-229 (1995); Ogura et al., Cancer Research 50: 5102-5106 (1990); Santodonato, et al., Human Gene Therapy 7:1-10 (1996); Santodonato, et al., Gene Therapy 4:1246-1255 (1997); and Zhang, et al., Cancer Gene Therapy 3: 31-38 (1996)), which are herein incorporated by reference. In one embodiment, the cells which are engineered are
25 arterial cells. The arterial cells may be reintroduced into the patient through direct injection to the artery, the tissues surrounding the artery, or through catheter injection.

As discussed in more detail below, the polynucleotide constructs can be delivered by any method that delivers injectable materials to the cells of an animal, such as, injection into the interstitial space of tissues (heart, muscle, skin, lung, liver,
30 and the like). The polynucleotide constructs may be delivered in a pharmaceutically acceptable liquid or aqueous carrier.

In one embodiment, the polynucleotide of the invention is delivered as a naked polynucleotide. The term "naked" polynucleotide, DNA or RNA refers to sequences that are free from any delivery vehicle that acts to assist, promote or facilitate entry into the cell, including viral sequences, viral particles, liposome formulations, lipofectin or precipitating agents and the like. However, the polynucleotides of the invention can also be delivered in liposome formulations and lipofectin formulations and the like can be prepared by methods well known to those skilled in the art. Such methods are described, for example, in U.S. Patent Nos. 5,593,972, 5,589,466, and 5,580,859, which are herein incorporated by reference.

The polynucleotide vector constructs of the invention used in the gene therapy method are preferably constructs that will not integrate into the host genome nor will they contain sequences that allow for replication. Appropriate vectors include pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; pSVK3, pBPV, pMSG and pSVL available from Pharmacia; and pEF1/V5, pcDNA3.1, and pRc/CMV2 available from Invitrogen. Other suitable vectors will be readily apparent to the skilled artisan.

Any strong promoter known to those skilled in the art can be used for driving the expression of polynucleotide sequence of the invention. Suitable promoters include adenoviral promoters, such as the adenoviral major late promoter; or heterologous promoters, such as the cytomegalovirus (CMV) promoter; the respiratory syncytial virus (RSV) promoter; inducible promoters, such as the MMT promoter, the metallothionein promoter; heat shock promoters; the albumin promoter; the ApoAI promoter; human globin promoters; viral thymidine kinase promoters, such as the Herpes Simplex thymidine kinase promoter; retroviral LTRs; the b-actin promoter; and human growth hormone promoters. The promoter also may be the native promoter for the polynucleotides of the invention.

Unlike other gene therapy techniques, one major advantage of introducing naked nucleic acid sequences into target cells is the transitory nature of the polynucleotide synthesis in the cells. Studies have shown that non-replicating DNA sequences can be introduced into cells to provide production of the desired polypeptide for periods of up to six months.

The polynucleotide construct of the invention can be delivered to the interstitial space of tissues within the an animal, including of muscle, skin, brain, lung, liver, spleen, bone marrow, thymus, heart, lymph, blood, bone, cartilage, pancreas, kidney, gall bladder, stomach, intestine, testis, ovary, uterus, rectum, nervous system, eye, gland, and connective tissue. Interstitial space of the tissues comprises the intercellular, fluid, mucopolysaccharide matrix among the reticular fibers of organ tissues, elastic fibers in the walls of vessels or chambers, collagen fibers of fibrous tissues, or that same matrix within connective tissue ensheathing muscle cells or in the lacunae of bone. It is similarly the space occupied by the plasma of the circulation and the lymph fluid of the lymphatic channels. Delivery to the interstitial space of muscle tissue is preferred for the reasons discussed below. They may be conveniently delivered by injection into the tissues comprising these cells. They are preferably delivered to and expressed in persistent, non-dividing cells which are differentiated, although delivery and expression may be achieved in non-differentiated or less completely differentiated cells, such as, for example, stem cells of blood or skin fibroblasts. *In vivo* muscle cells are particularly competent in their ability to take up and express polynucleotides.

For the nakednucleic acid sequence injection, an effective dosage amount of DNA or RNA will be in the range of from about 0.05 mg/kg body weight to about 50 mg/kg body weight. Preferably the dosage will be from about 0.005 mg/kg to about 20 mg/kg and more preferably from about 0.05 mg/kg to about 5 mg/kg. Of course, as the artisan of ordinary skill will appreciate, this dosage will vary according to the tissue site of injection. The appropriate and effective dosage of nucleic acid sequence can readily be determined by those of ordinary skill in the art and may depend on the condition being treated and the route of administration.

The preferred route of administration is by the parenteral route of injection into the interstitial space of tissues. However, other parenteral routes may also be used, such as, inhalation of an aerosol formulation particularly for delivery to lungs or bronchial tissues, throat or mucous membranes of the nose. In addition, naked DNA constructs can be delivered to arteries during angioplasty by the catheter used in the procedure.

The naked polynucleotides are delivered by any method known in the art, including, but not limited to, direct needle injection at the delivery site, intravenous injection, topical administration, catheter infusion, and so-called "gene guns". These delivery methods are known in the art.

5 The constructs may also be delivered with delivery vehicles such as viral sequences, viral particles, liposome formulations, lipofectin, precipitating agents, etc. Such methods of delivery are known in the art.

10 In certain embodiments, the polynucleotide constructs of the invention are complexed in a liposome preparation. Liposomal preparations for use in the instant invention include cationic (positively charged), anionic (negatively charged) and neutral preparations. However, cationic liposomes are particularly preferred because a tight charge complex can be formed between the cationic liposome and the polyanionic nucleic acid. Cationic liposomes have been shown to mediate intracellular delivery of plasmid DNA (Felgner et al., Proc. Natl. Acad. Sci. USA ,
15 84:7413-7416 (1987), which is herein incorporated by reference); mRNA (Malone et al., Proc. Natl. Acad. Sci. USA , 86:6077-6081 (1989), which is herein incorporated by reference); and purified transcription factors (Debs et al., J. Biol. Chem., 265:10189-10192 (1990), which is herein incorporated by reference), in functional form.

20 Cationic liposomes are readily available. For example, N[1-2,3-dioleoyloxy)propyl]-N,N,N-triethylammonium (DOTMA) liposomes are particularly useful and are available under the trademark Lipofectin. from GIBCO BRL, Grand Island, N. Y. (See, also, Felgner et al., Proc. Natl Acad. Sci. USA , 84:7413-7416 (1987), which is herein incorporated by reference). Other commercially
25 available liposomes include transfectace (DDAB/DOPE) and DOTAP/DOPE (Boehringer).

Other cationic liposomes can be prepared from readily available materials using techniques well known in the art. See, e.g. PCT Publication NO: WO 90/11092 (which is herein incorporated by reference) for a description of the synthesis of
30 DOTAP (1,2-bis(oleoyloxy)-3-(trimethylammonio)propane) liposomes. Preparation of DOTMA liposomes is explained in the literature, see, e.g., Felgner et al., Proc.

Natl. Acad. Sci. USA, 84:7413-7417, which is herein incorporated by reference.

Similar methods can be used to prepare liposomes from other cationic lipid materials.

Similarly, anionic and neutral liposomes are readily available, such as from Avanti Polar Lipids (Birmingham, Ala.), or can be easily prepared using readily available materials. Such materials include phosphatidyl, choline, cholesterol, phosphatidyl ethanolamine, dioleoylphosphatidyl choline (DOPC), dioleoylphosphatidyl glycerol (DOPG), dioleoylphosphatidyl ethanolamine (DOPE), among others. These materials can also be mixed with the DOTMA and DOTAP starting materials in appropriate ratios. Methods for making liposomes using these materials are well known in the art.

For example, commercially dioleoylphosphatidyl choline (DOPC), dioleoylphosphatidyl glycerol (DOPG), and dioleoylphosphatidyl ethanolamine (DOPE) can be used in various combinations to make conventional liposomes, with or without the addition of cholesterol. Thus, for example, DOPG/DOPC vesicles can be prepared by drying 50 mg each of DOPG and DOPC under a stream of nitrogen gas into a sonication vial. The sample is placed under a vacuum pump overnight and is hydrated the following day with deionized water. The sample is then sonicated for 2 hours in a capped vial, using a Heat Systems model 350 sonicator equipped with an inverted cup (bath type) probe at the maximum setting while the bath is circulated at 15EC. Alternatively, negatively charged vesicles can be prepared without sonication to produce multilamellar vesicles or by extrusion through nucleopore membranes to produce unilamellar vesicles of discrete size. Other methods are known and available to those of skill in the art.

The liposomes can comprise multilamellar vesicles (MLVs), small unilamellar vesicles (SUVs), or large unilamellar vesicles (LUVs), with SUVs being preferred. The various liposome-nucleic acid complexes are prepared using methods well known in the art. See, e.g., Straubinger et al., *Methods of Immunology*, 101:512-527 (1983), which is herein incorporated by reference. For example, MLVs containing nucleic acid can be prepared by depositing a thin film of phospholipid on the walls of a glass tube and subsequently hydrating with a solution of the material to be encapsulated. SUVs are prepared by extended sonication of MLVs to produce a homogeneous

population of unilamellar liposomes. The material to be entrapped is added to a suspension of preformed MLVs and then sonicated. When using liposomes containing cationic lipids, the dried lipid film is resuspended in an appropriate solution such as sterile water or an isotonic buffer solution such as 10 mM Tris/NaCl, sonicated, and then the preformed liposomes are mixed directly with the DNA. The liposome and DNA form a very stable complex due to binding of the positively charged liposomes to the cationic DNA. SUVs find use with small nucleic acid fragments. LUVs are prepared by a number of methods, well known in the art. Commonly used methods include Ca^{2+} -EDTA chelation (Papahadjopoulos et al., Biochim. Biophys. Acta, 394:483 (1975); Wilson et al., Cell, 17:77 (1979)); ether injection (Deamer et al., Biochim. Biophys. Acta, 443:629 (1976); Ostro et al., Biochem. Biophys. Res. Commun., 76:836 (1977); Fraley et al., Proc. Natl. Acad. Sci. USA, 76:3348 (1979)); detergent dialysis (Enoch et al., Proc. Natl. Acad. Sci. USA, 76:145 (1979)); and reverse-phase evaporation (REV) (Fraley et al., J. Biol. Chem., 255:10431 (1980); Szoka et al., Proc. Natl. Acad. Sci. USA, 75:145 (1978); Schaefer-Ridder et al., Science, 215:166 (1982)), which are herein incorporated by reference.

Generally, the ratio of DNA to liposomes will be from about 10:1 to about 1:10. Preferably, the ration will be from about 5:1 to about 1:5. More preferably, the ration will be about 3:1 to about 1:3. Still more preferably, the ratio will be about 1:1.

U.S. Patent NO: 5,676,954 (which is herein incorporated by reference) reports on the injection of genetic material, complexed with cationic liposomes carriers, into mice. U.S. Patent Nos. 4,897,355, 4,946,787, 5,049,386, 5,459,127, 5,589,466, 5,693,622, 5,580,859, 5,703,055, and international publication NO: WO 94/9469 (which are herein incorporated by reference) provide cationic lipids for use in transfecting DNA into cells and mammals. U.S. Patent Nos. 5,589,466, 5,693,622, 5,580,859, 5,703,055, and international publication NO: WO 94/9469 (which are herein incorporated by reference) provide methods for delivering DNA-cationic lipid complexes to mammals.

In certain embodiments, cells are engineered, *ex vivo* or *in vivo*, using a retroviral particle containing RNA which comprises a sequence encoding polypeptides of the invention. Retroviruses from which the retroviral plasmid vectors

may be derived include, but are not limited to, Moloney Murine Leukemia Virus, spleen necrosis virus, Rous sarcoma Virus, Harvey Sarcoma Virus, avian leukosis virus, gibbon ape leukemia virus, human immunodeficiency virus, Myeloproliferative Sarcoma Virus. and mammary tumor virus.

5 The retroviral plasmid vector is employed to transduce packaging cell lines to form producer cell lines. Examples of packaging cells which may be transfected include, but are not limited to, the PE501, PA317, R-2, R-AM, PA12, T19-14X, VT-19-17-H2, RCRE, RCRIP, GP+E-86, GP+envAm12, and DAN cell lines as described in Miller, Human Gene Therapy , 1:5-14 (1990), which is incorporated herein by
10 reference in its entirety. The vector may transduce the packaging cells through any means known in the art. Such means include, but are not limited to, electroporation, the use of liposomes, and CaPO₄ precipitation. In one alternative, the retroviral plasmid vector may be encapsulated into a liposome, or coupled to a lipid, and then administered to a host.

15 The producer cell line generates infectious retroviral vector particles which include polynucleotide encoding polypeptides of the invention. Such retroviral vector particles then may be employed, to transduce eukaryotic cells, either *in vitro* or *in vivo*. The transduced eukaryotic cells will express polypeptides of the invention.

 In certain other embodiments, cells are engineered, *ex vivo* or *in vivo*, with
20 polynucleotides of the invention contained in an adenovirus vector. Adenovirus can be manipulated such that it encodes and expresses polypeptides of the invention, and at the same time is inactivated in terms of its ability to replicate in a normal lytic viral life cycle. Adenovirus expression is achieved without integration of the viral DNA into the host cell chromosome, thereby alleviating concerns about insertional
25 mutagenesis. Furthermore, adenoviruses have been used as live enteric vaccines for many years with an excellent safety profile (Schwartz et al., Am. Rev. Respir. Dis., 109:233-238 (1974)). Finally, adenovirus mediated gene transfer has been demonstrated in a number of instances including transfer of alpha-1-antitrypsin and CFTR to the lungs of cotton rats (Rosenfeld et al., Science , 252:431-434 (1991);
30 Rosenfeld et al., Cell, 68:143-155 (1992)). Furthermore, extensive studies to attempt

to establish adenovirus as a causative agent in human cancer were uniformly negative (Green et al. Proc. Natl. Acad. Sci. USA , 76:6606 (1979)).

Suitable adenoviral vectors useful in the present invention are described, for example, in Kozarsky and Wilson, Curr. Opin. Genet. Devel., 3:499-503 (1993);
5 Rosenfeld et al., Ccell , 68:143-155 (1992); Engelhardt et al., Human Genet. Ther., 4:759-769 (1993); Yang et al., Nature Genet., 7:362-369 (1994); Wilson et al., Nature , 365:691-692 (1993); and U.S. Patent NO: 5,652,224, which are herein incorporated by reference. For example, the adenovirus vector Ad2 is useful and can be grown in human 293 cells. These cells contain the E1 region of adenovirus and
10 constitutively express E1a and E1b, which complement the defective adenoviruses by providing the products of the genes deleted from the vector. In addition to Ad2, other varieties of adenovirus (e.g., Ad3, Ad5, and Ad7) are also useful in the present invention.

Preferably, the adenoviruses used in the present invention are replication
15 deficient. Replication deficient adenoviruses require the aid of a helper virus and/or packaging cell line to form infectious particles. The resulting virus is capable of infecting cells and can express a polynucleotide of interest which is operably linked to a promoter, but cannot replicate in most cells. Replication deficient adenoviruses may be deleted in one or more of all or a portion of the following genes: E1a, E1b,
20 E3, E4, E2a, or L1 through L5.

In certain other embodiments, the cells are engineered, *ex vivo* or *in vivo*, using an adeno-associated virus (AAV). AAVs are naturally occurring defective viruses that require helper viruses to produce infectious particles (Muzyczka, Curr. Topics in Microbiol. Immunol., 158:97 (1992)). It is also one of the few viruses that
25 may integrate its DNA into non-dividing cells. Vectors containing as little as 300 base pairs of AAV can be packaged and can integrate, but space for exogenous DNA is limited to about 4.5 kb. Methods for producing and using such AAVs are known in the art. See, for example, U.S. Patent Nos. 5,139,941, 5,173,414, 5,354,678, 5,436,146, 5,474,935, 5,478,745, and 5,589,377.

30 For example, an appropriate AAV vector for use in the present invention will include all the sequences necessary for DNA replication, encapsidation, and host-cell

integration. The polynucleotide construct containing polynucleotides of the invention is inserted into the AAV vector using standard cloning methods, such as those found in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press (1989). The recombinant AAV vector is then transfected into packaging cells
5 which are infected with a helper virus, using any standard technique, including lipofection, electroporation, calcium phosphate precipitation, etc. Appropriate helper viruses include adenoviruses, cytomegaloviruses, vaccinia viruses, or herpes viruses. Once the packaging cells are transfected and infected, they will produce infectious AAV viral particles which contain the polynucleotide construct of the invention.
10 These viral particles are then used to transduce eukaryotic cells, either *ex vivo* or *in vivo*. The transduced cells will contain the polynucleotide construct integrated into its genome, and will express the desired gene product.

Another method of gene therapy involves operably associating heterologous control regions and endogenous polynucleotide sequences (e.g. encoding the
15 polypeptide sequence of interest) via homologous recombination (see, e.g., U.S. Patent NO: 5,641,670, issued June 24, 1997; International Publication NO: WO 96/29411, published September 26, 1996; International Publication NO: WO 94/12650, published August 4, 1994; Koller et al., *Proc. Natl. Acad. Sci. USA*, 86:8932-8935 (1989); and Zijlstra et al., *Nature*, 342:435-438 (1989). This method
20 involves the activation of a gene which is present in the target cells, but which is not normally expressed in the cells, or is expressed at a lower level than desired.

Polynucleotide constructs are made, using standard techniques known in the art, which contain the promoter with targeting sequences flanking the promoter. Suitable promoters are described herein. The targeting sequence is sufficiently
25 complementary to an endogenous sequence to permit homologous recombination of the promoter-targeting sequence with the endogenous sequence. The targeting sequence will be sufficiently near the 5' end of the desired endogenous polynucleotide sequence so the promoter will be operably linked to the endogenous sequence upon homologous recombination.

30 The promoter and the targeting sequences can be amplified using PCR. Preferably, the amplified promoter contains distinct restriction enzyme sites on the 5'

and 3' ends. Preferably, the 3' end of the first targeting sequence contains the same restriction enzyme site as the 5' end of the amplified promoter and the 5' end of the second targeting sequence contains the same restriction site as the 3' end of the amplified promoter. The amplified promoter and targeting sequences are digested
5 and ligated together.

The promoter-targeting sequence construct is delivered to the cells, either as naked polynucleotide, or in conjunction with transfection-facilitating agents, such as liposomes, viral sequences, viral particles, whole viruses, lipofection, precipitating agents, etc., described in more detail above. The P promoter-targeting sequence can
10 be delivered by any method, included direct needle injection, intravenous injection, topical administration, catheter infusion, particle accelerators, etc. The methods are described in more detail below.

The promoter-targeting sequence construct is taken up by cells. Homologous recombination between the construct and the endogenous sequence takes place, such
15 that an endogenous sequence is placed under the control of the promoter. The promoter then drives the expression of the endogenous sequence.

The polynucleotides encoding polypeptides of the present invention may be administered along with other polynucleotides encoding other angiogenic proteins. Angiogenic proteins include, but are not limited to, acidic and basic fibroblast growth
20 factors, VEGF-1, VEGF-2 (VEGF-C), VEGF-3 (VEGF-B), epidermal growth factor alpha and beta, platelet-derived endothelial cell growth factor, platelet-derived growth factor, tumor necrosis factor alpha, hepatocyte growth factor, insulin like growth factor, colony stimulating factor, macrophage colony stimulating factor, granulocyte/macrophage colony stimulating factor, and nitric oxide synthase.

25 Preferably, the polynucleotide encoding a polypeptide of the invention contains a secretory signal sequence that facilitates secretion of the protein. Typically, the signal sequence is positioned in the coding region of the polynucleotide to be expressed towards or at the 5' end of the coding region. The signal sequence may be homologous or heterologous to the polynucleotide of interest and may be
30 homologous or heterologous to the cells to be transfected. Additionally, the signal sequence may be chemically synthesized using methods known in the art.

Any mode of administration of any of the above-described polynucleotides constructs can be used so long as the mode results in the expression of one or more molecules in an amount sufficient to provide a therapeutic effect. This includes direct needle injection, systemic injection, catheter infusion, biolistic injectors, particle
5 accelerators (i.e., "gene guns"), gelfoam sponge depots, other commercially available depot materials, osmotic pumps (e.g., Alza minipumps), oral or suppositorial solid (tablet or pill) pharmaceutical formulations, and decanting or topical applications during surgery. For example, direct injection of naked calcium phosphate-precipitated plasmid into rat liver and rat spleen or a protein-coated
10 plasmid into the portal vein has resulted in gene expression of the foreign gene in the rat livers. (Kaneda et al., Science, 243:375 (1989)).

A preferred method of local administration is by direct injection. Preferably, a recombinant molecule of the present invention complexed with a delivery vehicle is administered by direct injection into or locally within the area of arteries.
15 Administration of a composition locally within the area of arteries refers to injecting the composition centimeters and preferably, millimeters within arteries.

Another method of local administration is to contact a polynucleotide construct of the present invention in or around a surgical wound. For example, a patient can undergo surgery and the polynucleotide construct can be coated on the
20 surface of tissue inside the wound or the construct can be injected into areas of tissue inside the wound.

Therapeutic compositions useful in systemic administration, include recombinant molecules of the present invention complexed to a targeted delivery vehicle of the present invention. Suitable delivery vehicles for use with systemic
25 administration comprise liposomes comprising ligands for targeting the vehicle to a particular site.

Preferred methods of systemic administration, include intravenous injection, aerosol, oral and percutaneous (topical) delivery. Intravenous injections can be performed using methods standard in the art. Aerosol delivery can also be performed
30 using methods standard in the art (see, for example, Stribling et al., Proc. Natl. Acad. Sci. USA, 189:11277-11281 (1992), which is incorporated herein by reference). Oral

delivery can be performed by complexing a polynucleotide construct of the present invention to a carrier capable of withstanding degradation by digestive enzymes in the gut of an animal. Examples of such carriers, include plastic capsules or tablets, such as those known in the art. Topical delivery can be performed by mixing a
5 polynucleotide construct of the present invention with a lipophilic reagent (e.g., DMSO) that is capable of passing into the skin.

Determining an effective amount of substance to be delivered can depend upon a number of factors including, for example, the chemical structure and biological activity of the substance, the age and weight of the animal, the precise
10 condition requiring treatment and its severity, and the route of administration. The frequency of treatments depends upon a number of factors, such as the amount of polynucleotide constructs administered per dose, as well as the health and history of the subject. The precise amount, number of doses, and timing of doses will be determined by the attending physician or veterinarian. Therapeutic compositions of
15 the present invention can be administered to any animal, preferably to mammals and birds. Preferred mammals include humans, dogs, cats, mice, rats, rabbits sheep, cattle, horses and pigs, with humans being particularly

Biological Activities

20 The polynucleotides or polypeptides, or agonists or antagonists of the present invention can be used in assays to test for one or more biological activities. If these polynucleotides and polypeptides do exhibit activity in a particular assay, it is likely that these molecules may be involved in the diseases associated with the biological activity. Thus, the polynucleotides or polypeptides, or agonists or antagonists could
25 be used to treat the associated disease.

Immune Activity

The polynucleotides or polypeptides, or agonists or antagonists of the present invention may be useful in treating, preventing, and/or diagnosing diseases, disorders,
30 and/or conditions of the immune system, by activating or inhibiting the proliferation, differentiation, or mobilization (chemotaxis) of immune cells. Immune cells develop

through a process called hematopoiesis, producing myeloid (platelets, red blood cells, neutrophils, and macrophages) and lymphoid (B and T lymphocytes) cells from pluripotent stem cells. The etiology of these immune diseases, disorders, and/or conditions may be genetic, somatic, such as cancer or some autoimmune diseases, disorders, and/or conditions, acquired (e.g., by chemotherapy or toxins), or infectious. Moreover, a polynucleotides or polypeptides, or agonists or antagonists of the present invention can be used as a marker or detector of a particular immune system disease or disorder.

A polynucleotides or polypeptides, or agonists or antagonists of the present invention may be useful in treating, preventing, and/or diagnosing diseases, disorders, and/or conditions of hematopoietic cells. A polynucleotides or polypeptides, or agonists or antagonists of the present invention could be used to increase differentiation and proliferation of hematopoietic cells, including the pluripotent stem cells, in an effort to treat or prevent those diseases, disorders, and/or conditions associated with a decrease in certain (or many) types hematopoietic cells. Examples of immunologic deficiency syndromes include, but are not limited to: blood protein diseases, disorders, and/or conditions (e.g. agammaglobulinemia, dysgammaglobulinemia), ataxia telangiectasia, common variable immunodeficiency, Digeorge Syndrome, HIV infection, HTLV-BLV infection, leukocyte adhesion deficiency syndrome, lymphopenia, phagocyte bactericidal dysfunction, severe combined immunodeficiency (SCIDs), Wiskott-Aldrich Disorder, anemia, thrombocytopenia, or hemoglobinuria.

Moreover, a polynucleotides or polypeptides, or agonists or antagonists of the present invention could also be used to modulate hemostatic (the stopping of bleeding) or thrombolytic activity (clot formation). For example, by increasing hemostatic or thrombolytic activity, a polynucleotides or polypeptides, or agonists or antagonists of the present invention could be used to treat or prevent blood coagulation diseases, disorders, and/or conditions (e.g., afibrinogenemia, factor deficiencies), blood platelet diseases, disorders, and/or conditions (e.g. thrombocytopenia), or wounds resulting from trauma, surgery, or other causes. Alternatively, a polynucleotides or polypeptides, or agonists or antagonists of the

present invention that can decrease hemostatic or thrombolytic activity could be used to inhibit or dissolve clotting. These molecules could be important in the treatment or prevention of heart attacks (infarction), strokes, or scarring.

5 A polynucleotides or polypeptides, or agonists or antagonists of the present invention may also be useful in treating, preventing, and/or diagnosing autoimmune diseases, disorders, and/or conditions. Many autoimmune diseases, disorders, and/or conditions result from inappropriate recognition of self as foreign material by immune cells. This inappropriate recognition results in an immune response leading to the destruction of the host tissue. Therefore, the administration of a polynucleotides or
10 polypeptides, or agonists or antagonists of the present invention that inhibits an immune response, particularly the proliferation, differentiation, or chemotaxis of T-cells, may be an effective therapy in preventing autoimmune diseases, disorders, and/or conditions.

Examples of autoimmune diseases, disorders, and/or conditions that can be
15 treated, prevented, and/or diagnosed or detected by the present invention include, but are not limited to: Addison's Disease, hemolytic anemia, antiphospholipid syndrome, rheumatoid arthritis, dermatitis, allergic encephalomyelitis, glomerulonephritis, Goodpasture's Syndrome, Graves' Disease, Multiple Sclerosis, Myasthenia Gravis, Neuritis, Ophthalmia, Bullous Pemphigoid, Pemphigus, Polyendocrinopathies,
20 Purpura, Reiter's Disease, Stiff-Man Syndrome. Autoimmune Thyroiditis, Systemic Lupus Erythematosus, Autoimmune Pulmonary Inflammation, Guillain-Barre Syndrome, insulin dependent diabetes mellitus, and autoimmune inflammatory eye disease.

Similarly, allergic reactions and conditions, such as asthma (particularly
25 allergic asthma) or other respiratory problems, may also be treated, prevented, and/or diagnosed by polynucleotides or polypeptides, or agonists or antagonists of the present invention. Moreover, these molecules can be used to treat anaphylaxis, hypersensitivity to an antigenic molecule, or blood group incompatibility.

A polynucleotides or polypeptides, or agonists or antagonists of the present
30 invention may also be used to treat, prevent, and/or diagnose organ rejection or graft-versus-host disease (GVHD). Organ rejection occurs by host immune cell destruction

of the transplanted tissue through an immune response. Similarly, an immune response is also involved in GVHD, but, in this case, the foreign transplanted immune cells destroy the host tissues. The administration of a polynucleotides or polypeptides, or agonists or antagonists of the present invention that inhibits an
5 immune response, particularly the proliferation, differentiation, or chemotaxis of T-cells, may be an effective therapy in preventing organ rejection or GVHD.

Similarly, a polynucleotides or polypeptides, or agonists or antagonists of the present invention may also be used to modulate inflammation. For example, the polypeptide or polynucleotide or agonists
10 or antagonist may inhibit the proliferation and differentiation of cells involved in an inflammatory response. These molecules can be used to treat, prevent, and/or diagnose inflammatory conditions, both chronic and acute conditions, including chronic prostatitis, granulomatous prostatitis and malacoplakia, inflammation associated with infection (e.g., septic shock, sepsis, or systemic inflammatory
15 response syndrome (SIRS)), ischemia-reperfusion injury, endotoxin lethality, arthritis, complement-mediated hyperacute rejection, nephritis, cytokine or chemokine induced lung injury, inflammatory bowel disease, Crohn's disease, or resulting from over production of cytokines (e.g., TNF or IL-1.)

20 **Hyperproliferative Disorders**

A polynucleotides or polypeptides, or agonists or antagonists of the invention can be used to treat, prevent, and/or diagnose hyperproliferative diseases, disorders, including neoplasms. A polynucleotides or polypeptides, or agonists or antagonists of the present invention may inhibit the proliferation of the disorder through direct or
25 indirect interactions. Alternatively, a polynucleotides or polypeptides, or agonists or antagonists of the present invention may proliferate other cells which can inhibit the hyperproliferative disorder.

For example, by increasing an immune response, particularly increasing antigenic qualities of the hyperproliferative disorder or by proliferating,
30 differentiating, or mobilizing T-cells, hyperproliferative diseases, disorders, and/or conditions can be treated, prevented, and/or diagnosed. This immune response may

be increased by either enhancing an existing immune response, or by initiating a new immune response. Alternatively, decreasing an immune response may also be a method of treating, preventing, and/or diagnosing hyperproliferative diseases, disorders, and/or conditions, such as a chemotherapeutic agent.

5 Examples of hyperproliferative diseases, disorders, and/or conditions that can be treated, prevented, and/or diagnosed by polynucleotides or polypeptides, or agonists or antagonists of the present invention include, but are not limited to neoplasms located in the: colon, abdomen, bone, breast, digestive system, liver, pancreas, peritoneum, endocrine glands (adrenal, parathyroid, pituitary, testicles,
10 ovary, thymus, thyroid), eye, head and neck, nervous (central and peripheral), lymphatic system, pelvic, skin, soft tissue, spleen, thoracic, and urogenital.

 Similarly, other hyperproliferative diseases, disorders, and/or conditions can also be treated, prevented, and/or diagnosed by a polynucleotides or polypeptides, or agonists or antagonists of the present invention. Examples of such hyperproliferative
15 diseases, disorders, and/or conditions include, but are not limited to: hypergammaglobulinemia, lymphoproliferative diseases, disorders, and/or conditions, paraproteinemias, purpura, sarcoidosis, Sezary Syndrome, Waldenström's Macroglobulinemia, Gaucher's Disease, histiocytosis, and any other hyperproliferative disease, besides neoplasia, located in an organ system listed above.

20 One preferred embodiment utilizes polynucleotides of the present invention to inhibit aberrant cellular division, by gene therapy using the present invention, and/or protein fusions or fragments thereof.

 Thus, the present invention provides a method for treating or preventing cell proliferative diseases, disorders, and/or conditions by inserting into an abnormally
25 proliferating cell a polynucleotide of the present invention, wherein said polynucleotide represses said expression.

 Another embodiment of the present invention provides a method of treating or preventing cell-proliferative diseases, disorders, and/or conditions in individuals comprising administration of one or more active gene copies of the present invention
30 to an abnormally proliferating cell or cells. In a preferred embodiment, polynucleotides of the present invention is a DNA construct comprising a

recombinant expression vector effective in expressing a DNA sequence encoding said polynucleotides. In another preferred embodiment of the present invention, the DNA construct encoding the polynucleotides of the present invention is inserted into cells to be treated utilizing a retrovirus, or more preferably an adenoviral vector (See G J. Nabel, et. al., PNAS 1999 96: 324-326, which is hereby incorporated by reference). In a most preferred embodiment, the viral vector is defective and will not transform non-proliferating cells, only proliferating cells. Moreover, in a preferred embodiment, the polynucleotides of the present invention inserted into proliferating cells either alone, or in combination with or fused to other polynucleotides, can then be modulated via an external stimulus (i.e. magnetic, specific small molecule, chemical, or drug administration, etc.), which acts upon the promoter upstream of said polynucleotides to induce expression of the encoded protein product. As such the beneficial therapeutic affect of the present invention may be expressly modulated (i.e. to increase, decrease, or inhibit expression of the present invention) based upon said external stimulus.

Polynucleotides of the present invention may be useful in repressing expression of oncogenic genes or antigens. By "repressing expression of the oncogenic genes " is intended the suppression of the transcription of the gene, the degradation of the gene transcript (pre-message RNA), the inhibition of splicing, the destruction of the messenger RNA, the prevention of the post-translational modifications of the protein, the destruction of the protein, or the inhibition of the normal function of the protein.

For local administration to abnormally proliferating cells, polynucleotides of the present invention may be administered by any method known to those of skill in the art including, but not limited to transfection, electroporation, microinjection of cells, or in vehicles such as liposomes, lipofectin, or as naked polynucleotides, or any other method described throughout the specification. The polynucleotide of the present invention may be delivered by known gene delivery systems such as, but not limited to, retroviral vectors (Gilboa, J. Virology 44:845 (1982); Hocke, Nature 320:275 (1986); Wilson, et al., Proc. Natl. Acad. Sci. U.S.A. 85:3014), vaccinia virus system (Chakrabarty et al., Mol. Cell Biol. 5:3403 (1985) or other efficient DNA

delivery systems (Yates et al., Nature 313:812 (1985)) known to those skilled in the art. These references are exemplary only and are hereby incorporated by reference. In order to specifically deliver or transfect cells which are abnormally proliferating and spare non-dividing cells, it is preferable to utilize a retrovirus, or adenoviral (as
5 described in the art and elsewhere herein) delivery system known to those of skill in the art. Since host DNA replication is required for retroviral DNA to integrate and the retrovirus will be unable to self replicate due to the lack of the retrovirus genes needed for its life cycle. Utilizing such a retroviral delivery system for polynucleotides of the present invention will target said gene and constructs to
10 abnormally proliferating cells and will spare the non-dividing normal cells.

The polynucleotides of the present invention may be delivered directly to cell proliferative disorder/disease sites in internal organs, body cavities and the like by use of imaging devices used to guide an injecting needle directly to the disease site. The polynucleotides of the present invention may also be administered to disease sites at
15 the time of surgical intervention.

By "cell proliferative disease" is meant any human or animal disease or disorder, affecting any one or any combination of organs, cavities, or body parts, which is characterized by single or multiple local abnormal proliferations of cells, groups of cells, or tissues, whether benign or malignant.

20 Any amount of the polynucleotides of the present invention may be administered as long as it has a biologically inhibiting effect on the proliferation of the treated cells. Moreover, it is possible to administer more than one of the polynucleotide of the present invention simultaneously to the same site. By "biologically inhibiting" is meant partial or total growth inhibition as well as
25 decreases in the rate of proliferation or growth of the cells. The biologically inhibitory dose may be determined by assessing the effects of the polynucleotides of the present invention on target malignant or abnormally proliferating cell growth in tissue culture, tumor growth in animals and cell cultures, or any other method known to one of ordinary skill in the art.

30 The present invention is further directed to antibody-based therapies which involve administering of anti-polypeptides and anti-polynucleotide antibodies to a

mammalian, preferably human, patient for treating, preventing, and/or diagnosing one or more of the described diseases, disorders, and/or conditions. Methods for producing anti-polypeptides and anti-polynucleotide antibodies polyclonal and monoclonal antibodies are described in detail elsewhere herein. Such antibodies may
5 be provided in pharmaceutically acceptable compositions as known in the art or as described herein.

A summary of the ways in which the antibodies of the present invention may be used therapeutically includes binding polynucleotides or polypeptides of the present invention locally or systemically in the body or by direct cytotoxicity of the
10 antibody, e.g. as mediated by complement (CDC) or by effector cells (ADCC). Some of these approaches are described in more detail below. Armed with the teachings provided herein, one of ordinary skill in the art will know how to use the antibodies of the present invention for diagnostic, monitoring or therapeutic purposes without undue experimentation.

15 In particular, the antibodies, fragments and derivatives of the present invention are useful for treating, preventing, and/or diagnosing a subject having or developing cell proliferative and/or differentiation diseases, disorders, and/or conditions as described herein. Such treatment comprises administering a single or multiple doses of the antibody, or a fragment, derivative, or a conjugate thereof.

20 The antibodies of this invention may be advantageously utilized in combination with other monoclonal or chimeric antibodies, or with lymphokines or hematopoietic growth factors, for example, which serve to increase the number or activity of effector cells which interact with the antibodies.

It is preferred to use high affinity and/or potent in vivo inhibiting and/or
25 neutralizing antibodies against polypeptides or polynucleotides of the present invention, fragments or regions thereof, for both immunoassays directed to and therapy of diseases, disorders, and/or conditions related to polynucleotides or polypeptides, including fragments thereof, of the present invention. Such antibodies, fragments, or regions, will preferably have an affinity for polynucleotides or
30 polypeptides, including fragments thereof. Preferred binding affinities include those with a dissociation constant or K_d less than $5 \times 10^{-6}M$, $10^{-6}M$, $5 \times 10^{-7}M$, $10^{-7}M$, $5 \times 10^{-8}M$.

8M , 10^{-8}M , $5 \times 10^{-9}\text{M}$, 10^{-9}M , $5 \times 10^{-10}\text{M}$, 10^{-10}M , $5 \times 10^{-11}\text{M}$, 10^{-11}M , $5 \times 10^{-12}\text{M}$, 10^{-12}M , $5 \times 10^{-13}\text{M}$, 10^{-13}M , $5 \times 10^{-14}\text{M}$, 10^{-14}M , $5 \times 10^{-15}\text{M}$, and 10^{-15}M .

Moreover, polypeptides of the present invention are useful in inhibiting the angiogenesis of proliferative cells or tissues, either alone, as a protein fusion, or in combination with other polypeptides directly or indirectly, as described elsewhere herein. In a most preferred embodiment, said anti-angiogenesis effect may be achieved indirectly, for example, through the inhibition of hematopoietic, tumor-specific cells, such as tumor-associated macrophages (See Joseph IB, et al. J Natl Cancer Inst, 90(21):1648-53 (1998), which is hereby incorporated by reference).

Antibodies directed to polypeptides or polynucleotides of the present invention may also result in inhibition of angiogenesis directly, or indirectly (See Witte L, et al., Cancer Metastasis Rev. 17(2):155-61 (1998), which is hereby incorporated by reference)).

Polypeptides, including protein fusions, of the present invention, or fragments thereof may be useful in inhibiting proliferative cells or tissues through the induction of apoptosis. Said polypeptides may act either directly, or indirectly to induce apoptosis of proliferative cells and tissues, for example in the activation of a death-domain receptor, such as tumor necrosis factor (TNF) receptor-1, CD95 (Fas/APO-1), TNF-receptor-related apoptosis-mediated protein (TRAMP) and TNF-related apoptosis-inducing ligand (TRAIL) receptor-1 and -2 (See Schulze-Osthoff K, et al., Eur J Biochem 254(3):439-59 (1998), which is hereby incorporated by reference).

Moreover, in another preferred embodiment of the present invention, said polypeptides may induce apoptosis through other mechanisms, such as in the activation of other proteins which will activate apoptosis, or through stimulating the expression of said proteins, either alone or in combination with small molecule drugs or adjuvants, such as apoptonin, galectins, thioredoxins, antiinflammatory proteins (See for example, Mutat Res 400(1-2):447-55 (1998), Med Hypotheses.50(5):423-33 (1998), Chem Biol Interact. Apr 24;111-112:23-34 (1998), J Mol Med.76(6):402-12 (1998), Int J Tissue React;20(1):3-15 (1998), which are all hereby incorporated by reference).

Polypeptides, including protein fusions to, or fragments thereof, of the present invention are useful in inhibiting the metastasis of proliferative cells or tissues.

Inhibition may occur as a direct result of administering polypeptides, or antibodies directed to said polypeptides as described elsewhere herein, or indirectly, such as

- 5 activating the expression of proteins known to inhibit metastasis, for example alpha 4 integrins, (See, e.g., Curr Top Microbiol Immunol 1998;231:125-41, which is hereby incorporated by reference). Such therapeutic affects of the present invention may be achieved either alone, or in combination with small molecule drugs or adjuvants.

- 10 In another embodiment, the invention provides a method of delivering compositions containing the polypeptides of the invention (e.g., compositions containing polypeptides or polypeptide antibodies associated with heterologous polypeptides, heterologous nucleic acids, toxins, or prodrugs) to targeted cells expressing the polypeptide of the present invention. Polypeptides or polypeptide antibodies of the invention may be associated with with heterologous polypeptides, 15 heterologous nucleic acids, toxins, or prodrugs via hydrophobic, hydrophilic, ionic and/or covalent interactions.

- Polypeptides, protein fusions to, or fragments thereof, of the present invention are useful in enhancing the immunogenicity and/or antigenicity of proliferating cells or tissues, either directly, such as would occur if the polypeptides of the present 20 invention 'vaccinated' the immune response to respond to proliferative antigens and immunogens, or indirectly, such as in activating the expression of proteins known to enhance the immune response (e.g. chemokines), to said antigens and immunogens.

Cardiovascular Disorders

- 25 Polynucleotides or polypeptides, or agonists or antagonists of the invention may be used to treat, prevent, and/or diagnose cardiovascular diseases, disorders, and/or conditions, including peripheral artery disease, such as limb ischemia.

- Cardiovascular diseases, disorders, and/or conditions include cardiovascular abnormalities, such as arterio-arterial fistula, arteriovenous fistula, cerebral 30 arteriovenous malformations, congenital heart defects, pulmonary atresia, and Scimitar Syndrome. Congenital heart defects include aortic coarctation, cor

triatratrium, coronary vessel anomalies, crisscross heart, dextrocardia, patent ductus arteriosus, Ebstein's anomaly, Eisenmenger complex, hypoplastic left heart syndrome, levocardia, tetralogy of fallot, transposition of great vessels, double outlet right ventricle, tricuspid atresia, persistent truncus arteriosus, and heart septal defects, such as aortopulmonary septal defect, endocardial cushion defects, Lutembacher's Syndrome, trilog of Fallot, ventricular heart septal defects.

Cardiovascular diseases, disorders, and/or conditions also include heart disease, such as arrhythmias, carcinoid heart disease, high cardiac output, low cardiac output, cardiac tamponade, endocarditis (including bacterial), heart aneurysm, cardiac arrest, congestive heart failure, congestive cardiomyopathy, paroxysmal dyspnea, cardiac edema, heart hypertrophy, congestive cardiomyopathy, left ventricular hypertrophy, right ventricular hypertrophy, post-infarction heart rupture, ventricular septal rupture, heart valve diseases, myocardial diseases, myocardial ischemia, pericardial effusion, pericarditis (including constrictive and tuberculous), pneumopericardium, postpericardiotomy syndrome, pulmonary heart disease, rheumatic heart disease, ventricular dysfunction, hyperemia, cardiovascular pregnancy complications, Scimitar Syndrome, cardiovascular syphilis, and cardiovascular tuberculosis.

Arrhythmias include sinus arrhythmia, atrial fibrillation, atrial flutter, bradycardia, extrasystole, Adams-Stokes Syndrome, bundle-branch block, sinoatrial block, long QT syndrome, parasystole, Lown-Ganong-Levine Syndrome, Mahaim-type pre-excitation syndrome, Wolff-Parkinson-White syndrome, sick sinus syndrome, tachycardias, and ventricular fibrillation. Tachycardias include paroxysmal tachycardia, supraventricular tachycardia, accelerated idioventricular rhythm, atrioventricular nodal reentry tachycardia, ectopic atrial tachycardia, ectopic junctional tachycardia, sinoatrial nodal reentry tachycardia, sinus tachycardia, Torsades de Pointes, and ventricular tachycardia.

Heart valve disease include aortic valve insufficiency, aortic valve stenosis, hear murmurs, aortic valve prolapse, mitral valve prolapse, tricuspid valve prolapse, mitral valve insufficiency, mitral valve stenosis, pulmonary atresia, pulmonary valve

insufficiency, pulmonary valve stenosis, tricuspid atresia, tricuspid valve insufficiency, and tricuspid valve stenosis.

Myocardial diseases include alcoholic cardiomyopathy, congestive cardiomyopathy, hypertrophic cardiomyopathy, aortic subvalvular stenosis, pulmonary subvalvular stenosis, restrictive cardiomyopathy, Chagas cardiomyopathy, endocardial fibroelastosis, endomyocardial fibrosis, Kearns Syndrome, myocardial reperfusion injury, and myocarditis.

Myocardial ischemias include coronary disease, such as angina pectoris, coronary aneurysm, coronary arteriosclerosis, coronary thrombosis, coronary vasospasm, myocardial infarction and myocardial stunning.

Cardiovascular diseases also include vascular diseases such as aneurysms, angiodyplasia, angiomatosis, bacillary angiomatosis, Hippel-Lindau Disease, Klippel-Trenaunay-Weber Syndrome, Sturge-Weber Syndrome, angioneurotic edema, aortic diseases, Takayasu's Arteritis, aortitis, Leriche's Syndrome, arterial occlusive diseases, arteritis, enarteritis, polyarteritis nodosa, cerebrovascular diseases, disorders, and/or conditions, diabetic angiopathies, diabetic retinopathy, embolisms, thrombosis, erythromelalgia, hemorrhoids, hepatic veno-occlusive disease, hypertension, hypotension, ischemia, peripheral vascular diseases, phlebitis, pulmonary veno-occlusive disease, Raynaud's disease, CREST syndrome, retinal vein occlusion, Scimitar syndrome, superior vena cava syndrome, telangiectasia, atacia telangiectasia, hereditary hemorrhagic telangiectasia, varicocele, varicose veins, varicose ulcer, vasculitis, and venous insufficiency.

Aneurysms include dissecting aneurysms, false aneurysms, infected aneurysms, ruptured aneurysms, aortic aneurysms, cerebral aneurysms, coronary aneurysms, heart aneurysms, and iliac aneurysms.

Arterial occlusive diseases include arteriosclerosis, intermittent claudication, carotid stenosis, fibromuscular dysplasias, mesenteric vascular occlusion, Moyamoya disease, renal artery obstruction, retinal artery occlusion, and thromboangiitis obliterans.

Cerebrovascular diseases, disorders, and/or conditions include carotid artery diseases, cerebral amyloid angiopathy, cerebral aneurysm, cerebral anoxia, cerebral

arteriosclerosis, cerebral arteriovenous malformation, cerebral artery diseases, cerebral embolism and thrombosis, carotid artery thrombosis, sinus thrombosis, Wallenberg's syndrome, cerebral hemorrhage, epidural hematoma, subdural hematoma, subarachnoid hemorrhage, cerebral infarction, cerebral ischemia
5 (including transient), subclavian steal syndrome, periventricular leukomalacia, vascular headache, cluster headache, migraine, and vertebrobasilar insufficiency.

Embolisms include air embolisms, amniotic fluid embolisms, cholesterol embolisms, blue toe syndrome, fat embolisms, pulmonary embolisms; and thromboembolisms. Thrombosis include coronary thrombosis, hepatic vein
10 thrombosis, retinal vein occlusion, carotid artery thrombosis, sinus thrombosis, Wallenberg's syndrome, and thrombophlebitis.

Ischemia includes cerebral ischemia, ischemic colitis, compartment syndromes, anterior compartment syndrome, myocardial ischemia, reperfusion injuries, and peripheral limb ischemia. Vasculitis includes aortitis, arteritis, Behcet's
15 Syndrome, Churg-Strauss Syndrome, mucocutaneous lymph node syndrome, thromboangiitis obliterans, hypersensitivity vasculitis, Schoenlein-Henoch purpura, allergic cutaneous vasculitis, and Wegener's granulomatosis.

Polynucleotides or polypeptides, or agonists or antagonists of the invention, are especially effective for the treatment of critical limb ischemia and coronary
20 disease.

Polypeptides may be administered using any method known in the art, including, but not limited to, direct needle injection at the delivery site, intravenous injection, topical administration, catheter infusion, biolistic injectors, particle accelerators, gel foam sponge depots, other commercially available depot materials,
25 osmotic pumps, oral or suppositorial solid pharmaceutical formulations, decanting or topical applications during surgery, aerosol delivery. Such methods are known in the art. Polypeptides of the invention may be administered as part of a *Therapeutic*, described in more detail below. Methods of delivering polynucleotides of the invention are described in more detail herein.

30

Anti-Angiogenesis Activity

The naturally occurring balance between endogenous stimulators and inhibitors of angiogenesis is one in which inhibitory influences predominate. Rastinejad *et al.*, *Cell* 56:345-355 (1989). In those rare instances in which neovascularization occurs under normal physiological conditions, such as wound healing, organ regeneration, embryonic development, and female reproductive processes, angiogenesis is stringently regulated and spatially and temporally delimited. Under conditions of pathological angiogenesis such as that characterizing solid tumor growth, these regulatory controls fail. Unregulated angiogenesis becomes pathologic and sustains progression of many neoplastic and non-neoplastic diseases.

5 A number of serious diseases are dominated by abnormal neovascularization including solid tumor growth and metastases, arthritis, some types of eye diseases, disorders, and/or conditions, and psoriasis. See, e.g., reviews by Moses *et al.*, *Biotech.* 9:630-634 (1991); Folkman *et al.*, *N. Engl. J. Med.*, 333:1757-1763 (1995); Auerbach *et al.*, *J. Microvasc. Res.* 29:401-411 (1985); Folkman, *Advances in*

15 *Cancer Research*, eds. Klein and Weinhouse, Academic Press, New York, pp. 175-203 (1985); Patz, *Am. J. Ophthalmol.* 94:715-743 (1982); and Folkman *et al.*, *Science* 221:719-725 (1983). In a number of pathological conditions, the process of angiogenesis contributes to the disease state. For example, significant data have accumulated which suggest that the growth of solid tumors is dependent on

20 angiogenesis. Folkman and Klagsbrun, *Science* 235:442-447 (1987).

The present invention provides for treatment of diseases, disorders, and/or conditions associated with neovascularization by administration of the polynucleotides and/or polypeptides of the invention, as well as agonists or antagonists of the present invention. Malignant and metastatic conditions which can

25 be treated with the polynucleotides and polypeptides, or agonists or antagonists of the invention include, but are not limited to, malignancies, solid tumors, and cancers described herein and otherwise known in the art (for a review of such disorders, see Fishman *et al.*, *Medicine*, 2d Ed., J. B. Lippincott Co., Philadelphia (1985)). Thus, the present invention provides a method of treating, preventing, and/or diagnosing an

30 angiogenesis-related disease and/or disorder, comprising administering to an individual in need thereof a therapeutically effective amount of a polynucleotide,

polypeptide, antagonist and/or agonist of the invention. For example, polynucleotides, polypeptides, antagonists and/or agonists may be utilized in a variety of additional methods in order to therapeutically treat or prevent a cancer or tumor. Cancers which may be treated, prevented, and/or diagnosed with polynucleotides, polypeptides, antagonists and/or agonists include, but are not limited to solid tumors, including prostate, lung, breast, ovarian, stomach, pancreas, larynx, esophagus, testes, liver, parotid, biliary tract, colon, rectum, cervix, uterus, endometrium, kidney, bladder, thyroid cancer; primary tumors and metastases; melanomas; glioblastoma; Kaposi's sarcoma; leiomyosarcoma; non-small cell lung cancer; colorectal cancer; advanced malignancies; and blood born tumors such as leukemias. For example, polynucleotides, polypeptides, antagonists and/or agonists may be delivered topically, in order to treat or prevent cancers such as skin cancer, head and neck tumors, breast tumors, and Kaposi's sarcoma.

Within yet other aspects, polynucleotides, polypeptides, antagonists and/or agonists may be utilized to treat superficial forms of bladder cancer by, for example, intravesical administration. Polynucleotides, polypeptides, antagonists and/or agonists may be delivered directly into the tumor, or near the tumor site, via injection or a catheter. Of course, as the artisan of ordinary skill will appreciate, the appropriate mode of administration will vary according to the cancer to be treated. Other modes of delivery are discussed herein.

Polynucleotides, polypeptides, antagonists and/or agonists may be useful in treating, preventing, and/or diagnosing other diseases, disorders, and/or conditions, besides cancers, which involve angiogenesis. These diseases, disorders, and/or conditions include, but are not limited to: benign tumors, for example hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas; arteriosclerotic plaques; ocular angiogenic diseases, for example, diabetic retinopathy, retinopathy of prematurity, macular degeneration, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, rubeosis, retinoblastoma, uveitis and Pterygia (abnormal blood vessel growth) of the eye; rheumatoid arthritis; psoriasis; delayed wound healing; endometriosis; vasculogenesis; granulations; hypertrophic scars (keloids); nonunion fractures; scleroderma; trachoma; vascular adhesions; myocardial

angiogenesis; coronary collaterals; cerebral collaterals; arteriovenous malformations; ischemic limb angiogenesis; Osler-Webber Syndrome; plaque neovascularization; telangiectasia; hemophiliac joints; angiofibroma; fibromuscular dysplasia; wound granulation; Crohn's disease; and atherosclerosis.

5 For example, within one aspect of the present invention methods are provided for treating, preventing, and/or diagnosing hypertrophic scars and keloids, comprising the step of administering a polynucleotide, polypeptide, antagonist and/or agonist of the invention to a hypertrophic scar or keloid.

 Within one embodiment of the present invention polynucleotides,
10 polypeptides, antagonists and/or agonists are directly injected into a hypertrophic scar or keloid, in order to prevent the progression of these lesions. This therapy is of particular value in the prophylactic treatment of conditions which are known to result in the development of hypertrophic scars and keloids (e.g., burns), and is preferably initiated after the proliferative phase has had time to progress (approximately 14 days
15 after the initial injury), but before hypertrophic scar or keloid development. As noted above, the present invention also provides methods for treating, preventing, and/or diagnosing neovascular diseases of the eye, including for example, corneal neovascularization, neovascular glaucoma, proliferative diabetic retinopathy, retrolental fibroplasia and macular degeneration.

20 Moreover, Ocular diseases, disorders, and/or conditions associated with neovascularization which can be treated, prevented, and/or diagnosed with the polynucleotides and polypeptides of the present invention (including agonists and/or antagonists) include, but are not limited to: neovascular glaucoma, diabetic retinopathy, retinoblastoma, retrolental fibroplasia, uveitis, retinopathy of prematurity
25 macular degeneration, corneal graft neovascularization, as well as other eye inflammatory diseases, ocular tumors and diseases associated with choroidal or iris neovascularization. See, e.g., reviews by Waltman *et al.*, *Am. J. Ophthalm.* 85:704-710 (1978) and Gartner *et al.*, *Surv. Ophthalm.* 22:291-312 (1978).

 Thus, within one aspect of the present invention methods are provided for
30 treating or preventing neovascular diseases of the eye such as corneal neovascularization (including corneal graft neovascularization), comprising the step

of administering to a patient a therapeutically effective amount of a compound (as described above) to the cornea, such that the formation of blood vessels is inhibited. Briefly, the cornea is a tissue which normally lacks blood vessels. In certain pathological conditions however, capillaries may extend into the cornea from the pericorneal vascular plexus of the limbus. When the cornea becomes vascularized, it also becomes clouded, resulting in a decline in the patient's visual acuity. Visual loss may become complete if the cornea completely opacitates. A wide variety of diseases, disorders, and/or conditions can result in corneal neovascularization, including for example, corneal infections (e.g., trachoma, herpes simplex keratitis, leishmaniasis and onchocerciasis), immunological processes (e.g., graft rejection and Stevens-Johnson's syndrome), alkali burns, trauma, inflammation (of any cause), toxic and nutritional deficiency states, and as a complication of wearing contact lenses.

Within particularly preferred embodiments of the invention, may be prepared for topical administration in saline (combined with any of the preservatives and antimicrobial agents commonly used in ocular preparations), and administered in eyedrop form. The solution or suspension may be prepared in its pure form and administered several times daily. Alternatively, anti-angiogenic compositions, prepared as described above, may also be administered directly to the cornea. Within preferred embodiments, the anti-angiogenic composition is prepared with a muco-adhesive polymer which binds to cornea. Within further embodiments, the anti-angiogenic factors or anti-angiogenic compositions may be utilized as an adjunct to conventional steroid therapy. Topical therapy may also be useful prophylactically in corneal lesions which are known to have a high probability of inducing an angiogenic response (such as chemical burns). In these instances the treatment, likely in combination with steroids, may be instituted immediately to help prevent subsequent complications.

Within other embodiments, the compounds described above may be injected directly into the corneal stroma by an ophthalmologist under microscopic guidance. The preferred site of injection may vary with the morphology of the individual lesion, but the goal of the administration would be to place the composition at the advancing

front of the vasculature (i.e., interspersed between the blood vessels and the normal cornea). In most cases this would involve perilimbic corneal injection to "protect" the cornea from the advancing blood vessels. This method may also be utilized shortly after a corneal insult in order to prophylactically prevent corneal neovascularization.

- 5 In this situation the material could be injected in the perilimbic cornea interspersed between the corneal lesion and its undesired potential limbic blood supply. Such methods may also be utilized in a similar fashion to prevent capillary invasion of transplanted corneas. In a sustained-release form injections might only be required 2-3 times per year. A steroid could also be added to the injection solution to reduce
10 inflammation resulting from the injection itself.

- Within another aspect of the present invention, methods are provided for treating or preventing neovascular glaucoma, comprising the step of administering to a patient a therapeutically effective amount of a polynucleotide, polypeptide, antagonist and/or agonist to the eye, such that the formation of blood vessels is
15 inhibited. In one embodiment, the compound may be administered topically to the eye in order to treat or prevent early forms of neovascular glaucoma. Within other embodiments, the compound may be implanted by injection into the region of the anterior chamber angle. Within other embodiments, the compound may also be placed in any location such that the compound is continuously released into the
20 aqueous humor. Within another aspect of the present invention, methods are provided for treating or preventing proliferative diabetic retinopathy, comprising the step of administering to a patient a therapeutically effective amount of a polynucleotide, polypeptide, antagonist and/or agonist to the eyes, such that the formation of blood vessels is inhibited.

- 25 Within particularly preferred embodiments of the invention, proliferative diabetic retinopathy may be treated by injection into the aqueous humor or the vitreous, in order to increase the local concentration of the polynucleotide, polypeptide, antagonist and/or agonist in the retina. Preferably, this treatment should be initiated prior to the acquisition of severe disease requiring photocoagulation.

- 30 Within another aspect of the present invention, methods are provided for treating or preventing retrolental fibroplasia, comprising the step of administering to a

patient a therapeutically effective amount of a polynucleotide, polypeptide, antagonist and/or agonist to the eye, such that the formation of blood vessels is inhibited. The compound may be administered topically, via intravitreal injection and/or via intraocular implants.

5 Additionally, diseases, disorders, and/or conditions which can be treated, prevented, and/or diagnosed with the polynucleotides, polypeptides, agonists and/or agonists include, but are not limited to, hemangioma, arthritis, psoriasis, angiofibroma, atherosclerotic plaques, delayed wound healing, granulations, hemophilic joints, hypertrophic scars, nonunion fractures, Osler-Weber syndrome,
10 pyogenic granuloma, scleroderma, trachoma, and vascular adhesions.

 Moreover, diseases, disorders, and/or conditions and/or states, which can be treated, prevented, and/or diagnosed with the the polynucleotides, polypeptides, agonists and/or agonists include, but are not limited to, solid tumors, blood born tumors such as leukemias, tumor metastasis, Kaposi's sarcoma, benign tumors, for
15 example hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas, rheumatoid arthritis, psoriasis, ocular angiogenic diseases, for example, diabetic retinopathy, retinopathy of prematurity, macular degeneration, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, rubeosis, retinoblastoma, and uveitis, delayed wound healing, endometriosis, vasculogenesis, granulations,
20 hypertrophic scars (keloids), nonunion fractures, scleroderma, trachoma, vascular adhesions, myocardial angiogenesis, coronary collaterals, cerebral collaterals, arteriovenous malformations, ischemic limb angiogenesis, Osler-Webber Syndrome, plaque neovascularization, telangiectasia, hemophilic joints, angiofibroma fibromuscular dysplasia, wound granulation, Crohn's disease, atherosclerosis, birth
25 control agent by preventing vascularization required for embryo implantation controlling menstruation, diseases that have angiogenesis as a pathologic consequence such as cat scratch disease (*Rochelle minalia quintosa*), ulcers (*Helicobacter pylori*), Bartonellosis and bacillary angiomatosis.

 In one aspect of the birth control method, an amount of the compound
30 sufficient to block embryo implantation is administered before or after intercourse and fertilization have occurred, thus providing an effective method of birth control,

possibly a "morning after" method. Polynucleotides, polypeptides, agonists and/or agonists may also be used in controlling menstruation or administered as either a peritoneal lavage fluid or for peritoneal implantation in the treatment of endometriosis.

5 Polynucleotides, polypeptides, agonists and/or agonists of the present invention may be incorporated into surgical sutures in order to prevent stitch granulomas.

Polynucleotides, polypeptides, agonists and/or agonists may be utilized in a wide variety of surgical procedures. For example, within one aspect of the present invention a compositions (in the form of, for example, a spray or film) may be utilized
10 to coat or spray an area prior to removal of a tumor, in order to isolate normal surrounding tissues from malignant tissue, and/or to prevent the spread of disease to surrounding tissues. Within other aspects of the present invention, compositions (e.g., in the form of a spray) may be delivered via endoscopic procedures in order to coat
15 tumors, or inhibit angiogenesis in a desired locale. Within yet other aspects of the present invention, surgical meshes which have been coated with anti- angiogenic compositions of the present invention may be utilized in any procedure wherein a surgical mesh might be utilized. For example, within one embodiment of the invention a surgical mesh laden with an anti-angiogenic composition may be utilized
20 during abdominal cancer resection surgery (e.g., subsequent to colon resection) in order to provide support to the structure, and to release an amount of the anti-angiogenic factor.

Within further aspects of the present invention, methods are provided for treating tumor excision sites, comprising administering a polynucleotide, polypeptide,
25 agonist and/or agonist to the resection margins of a tumor subsequent to excision, such that the local recurrence of cancer and the formation of new blood vessels at the site is inhibited. Within one embodiment of the invention, the anti-angiogenic compound is administered directly to the tumor excision site (e.g., applied by swabbing, brushing or otherwise coating the resection margins of the tumor with the
30 anti-angiogenic compound). Alternatively, the anti-angiogenic compounds may be incorporated into known surgical pastes prior to administration. Within particularly

preferred embodiments of the invention, the anti-angiogenic compounds are applied after hepatic resections for malignancy, and after neurosurgical operations.

Within one aspect of the present invention, polynucleotides, polypeptides, agonists and/or agonists may be administered to the resection margin of a wide
5 variety of tumors, including for example, breast, colon, brain and hepatic tumors. For example, within one embodiment of the invention, anti-angiogenic compounds may be administered to the site of a neurological tumor subsequent to excision, such that the formation of new blood vessels at the site are inhibited.

The polynucleotides, polypeptides, agonists and/or agonists of the present
10 invention may also be administered along with other anti-angiogenic factors. Representative examples of other anti-angiogenic factors include: Anti-Invasive Factor, retinoic acid and derivatives thereof, paclitaxel, Suramin, Tissue Inhibitor of Metalloproteinase-1, Tissue Inhibitor of Metalloproteinase-2, Plasminogen Activator Inhibitor-1, Plasminogen Activator Inhibitor-2, and various forms of the lighter "d
15 group" transition metals.

Lighter "d group" transition metals include, for example, vanadium, molybdenum, tungsten, titanium, niobium, and tantalum species. Such transition metal species may form transition metal complexes. Suitable complexes of the above-mentioned transition metal species include oxo transition metal complexes.

Representative examples of vanadium complexes include oxo vanadium
20 complexes such as vanadate and vanadyl complexes. Suitable vanadate complexes include metavanadate and orthovanadate complexes such as, for example, ammonium metavanadate, sodium metavanadate, and sodium orthovanadate. Suitable vanadyl complexes include, for example, vanadyl acetylacetonate and vanadyl sulfate
25 including vanadyl sulfate hydrates such as vanadyl sulfate mono- and trihydrates.

Representative examples of tungsten and molybdenum complexes also include oxo complexes. Suitable oxo tungsten complexes include tungstate and tungsten oxide complexes. Suitable tungstate complexes include ammonium tungstate, calcium tungstate, sodium tungstate dihydrate, and tungstic acid. Suitable tungsten
30 oxides include tungsten (IV) oxide and tungsten (VI) oxide. Suitable oxo molybdenum complexes include molybdate, molybdenum oxide, and molybdenyl

complexes. Suitable molybdate complexes include ammonium molybdate and its hydrates, sodium molybdate and its hydrates, and potassium molybdate and its hydrates. Suitable molybdenum oxides include molybdenum (VI) oxide, molybdenum (VI) oxide, and molybdic acid. Suitable molybdenyl complexes include, for example,
 5 molybdenyl acetylacetonate. Other suitable tungsten and molybdenum complexes include hydroxo derivatives derived from, for example, glycerol, tartaric acid, and sugars.

A wide variety of other anti-angiogenic factors may also be utilized within the context of the present invention. Representative examples include platelet factor 4;
 10 protamine sulphate; sulphated chitin derivatives (prepared from queen crab shells), (Murata et al., Cancer Res. 51:22-26, 1991); Sulphated Polysaccharide Peptidoglycan Complex (SP- PG) (the function of this compound may be enhanced by the presence of steroids such as estrogen, and tamoxifen citrate); Staurosporine; modulators of matrix metabolism, including for example, proline analogs, cis-hydroxyproline, d,L-
 15 3,4-dehydroproline, Thiaproline, alpha,alpha-dipyridyl, aminopropionitrile fumarate; 4-propyl-5-(4-pyridinyl)-2(3H)-oxazolone; Methotrexate; Mitoxantrone; Heparin; Interferons; 2 Macroglobulin-serum; ChIMP-3 (Pavloff et al., J. Bio. Chem. 267:17321-17326, 1992); Chymostatin (Tomkinson et al., Biochem J. 286:475-480, 1992); Cyclodextrin Tetradesulfate; Eponemycin; Camptothecin; Fumagillin
 20 (Ingber et al., Nature 348:555-557, 1990); Gold Sodium Thiomalate ("GST"; Matsubara and Ziff, J. Clin. Invest. 79:1440-1446, 1987); anticollagenase-serum; alpha2-antiplasmin (Holmes et al., J. Biol. Chem. 262(4):1659-1664, 1987); Bisantrene (National Cancer Institute); Lobenzarit disodium (N-(2)-carboxyphenyl-4-chloroanthronilic acid disodium or "CCA"; Takeuchi et al., Agents Actions 36:312-
 25 316, 1992); Thalidomide; Angostatic steroid; AGM-1470; carboxyaminolimidazole; and metalloproteinase inhibitors such as BB94.

Diseases at the Cellular Level

Diseases associated with increased cell survival or the inhibition of apoptosis
 30 that could be treated, prevented, and/or diagnosed by the polynucleotides or polypeptides and/or antagonists or agonists of the invention, include cancers (such as

follicular lymphomas, carcinomas with p53 mutations, and hormone-dependent tumors, including, but not limited to colon cancer, cardiac tumors, pancreatic cancer, melanoma, retinoblastoma, glioblastoma, lung cancer, intestinal cancer, testicular cancer, stomach cancer, neuroblastoma, myxoma, myoma, lymphoma, endothelioma, osteoblastoma, osteoclastoma, osteosarcoma, chondrosarcoma, adenoma, breast cancer, prostate cancer, Kaposi's sarcoma and ovarian cancer); autoimmune diseases, disorders, and/or conditions (such as, multiple sclerosis, Sjogren's syndrome, Hashimoto's thyroiditis, biliary cirrhosis, Behcet's disease, Crohn's disease, polymyositis, systemic lupus erythematosus and immune-related glomerulonephritis and rheumatoid arthritis) and viral infections (such as herpes viruses, pox viruses and adenoviruses), inflammation, graft v. host disease, acute graft rejection, and chronic graft rejection. In preferred embodiments, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention are used to inhibit growth, progression, and/or metastasis of cancers, in particular those listed above.

Additional diseases or conditions associated with increased cell survival that could be treated, prevented or diagnosed by the polynucleotides or polypeptides, or agonists or antagonists of the invention, include, but are not limited to, progression, and/or metastases of malignancies and related disorders such as leukemia (including acute leukemias (e.g., acute lymphocytic leukemia, acute myelocytic leukemia (including myeloblastic, promyelocytic, myelomonocytic, monocytic, and erythroleukemia)) and chronic leukemias (e.g., chronic myelocytic (granulocytic) leukemia and chronic lymphocytic leukemia)), polycythemia vera, lymphomas (e.g., Hodgkin's disease and non-Hodgkin's disease), multiple myeloma, Waldenstrom's macroglobulinemia, heavy chain disease, and solid tumors including, but not limited to, sarcomas and carcinomas such as fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary

carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilm's tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, 5 craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, melanoma, neuroblastoma, and retinoblastoma.

Diseases associated with increased apoptosis that could be treated, prevented, and/or diagnosed by the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, include AIDS; neurodegenerative diseases, disorders, 10 and/or conditions (such as Alzheimer's disease, Parkinson's disease, Amyotrophic lateral sclerosis, Retinitis pigmentosa, Cerebellar degeneration and brain tumor or prior associated disease); autoimmune diseases, disorders, and/or conditions (such as, multiple sclerosis, Sjogren's syndrome, Hashimoto's thyroiditis, biliary cirrhosis, Behcet's disease, Crohn's disease, polymyositis, systemic lupus erythematosus and 15 immune-related glomerulonephritis and rheumatoid arthritis) myelodysplastic syndromes (such as aplastic anemia), graft v. host disease, ischemic injury (such as that caused by myocardial infarction, stroke and reperfusion injury), liver injury (e.g., hepatitis related liver injury, ischemia/reperfusion injury, cholestasis (bile duct injury) and liver cancer); toxin-induced liver disease (such as that caused by alcohol), septic 20 shock, cachexia and anorexia.

Wound Healing and Epithelial Cell Proliferation

In accordance with yet a further aspect of the present invention, there is provided a process for utilizing the polynucleotides or polypeptides, and/or agonists 25 or antagonists of the invention, for therapeutic purposes, for example, to stimulate epithelial cell proliferation and basal keratinocytes for the purpose of wound healing, and to stimulate hair follicle production and healing of dermal wounds. Polynucleotides or polypeptides, as well as agonists or antagonists of the invention, may be clinically useful in stimulating wound healing including surgical wounds, 30 excisional wounds, deep wounds involving damage of the dermis and epidermis, eye tissue wounds, dental tissue wounds, oral cavity wounds, diabetic ulcers, dermal

ulcers, cubitus ulcers, arterial ulcers, venous stasis ulcers, burns resulting from heat exposure or chemicals, and other abnormal wound healing conditions such as uremia, malnutrition, vitamin deficiencies and complications associated with systemic treatment with steroids, radiation therapy and antineoplastic drugs and

5 antimetabolites. Polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to promote dermal reestablishment subsequent to dermal loss

The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to increase the adherence of skin grafts to a wound bed and

10 to stimulate re-epithelialization from the wound bed. The following are a non-exhaustive list of grafts that polynucleotides or polypeptides, agonists or antagonists of the invention, could be used to increase adherence to a wound bed: autografts, artificial skin, allografts, autodermic graft, autocpdermic grafts, avacular grafts, Blair-Brown grafts, bone graft, brephoplastic grafts, cutis graft, delayed graft, dermic graft,

15 epidermic graft, fascia graft, full thickness graft, heterologous graft, xenograft, homologous graft, hyperplastic graft, lamellar graft, mesh graft, mucosal graft, Ollier-Thiersch graft, omenpal graft, patch graft, pedicle graft, penetrating graft, split skin graft, thick split graft. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, can be used to promote skin strength and to improve the

20 appearance of aged skin.

It is believed that the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, will also produce changes in hepatocyte proliferation, and epithelial cell proliferation in the lung, breast, pancreas, stomach, small intesting, and large intestine. The polynucleotides or polypeptides, and/or agonists or

25 antagonists of the invention, could promote proliferation of epithelial cells such as sebocytes, hair follicles, hepatocytes, type II pneumocytes, mucin-producing goblet cells, and other epithelial cells and their progenitors contained within the skin, lung, liver, and gastrointestinal tract. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, may promote proliferation of endothelial cells,

30 keratinocytes, and basal keratinocytes.

The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could also be used to reduce the side effects of gut toxicity that result from radiation, chemotherapy treatments or viral infections. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, may have a
5 cytoprotective effect on the small intestine mucosa. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, may also stimulate healing of mucositis (mouth ulcers) that result from chemotherapy and viral infections.

The polynucleotides or polypeptides, and/or agonists or antagonists of the
10 invention, could further be used in full regeneration of skin in full and partial thickness skin defects, including burns, (i.e., repopulation of hair follicles, sweat glands, and sebaceous glands), treatment of other skin defects such as psoriasis. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to treat epidermolysis bullosa, a defect in adherence of the epidermis to the
15 underlying dermis which results in frequent, open and painful blisters by accelerating reepithelialization of these lesions. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could also be used to treat gastric and duodenal ulcers and help heal by scar formation of the mucosal lining and regeneration of glandular mucosa and duodenal mucosal lining more rapidly.
20 Inflammatory bowel diseases, such as Crohn's disease and ulcerative colitis, are diseases which result in destruction of the mucosal surface of the small or large intestine, respectively. Thus, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to promote the resurfacing of the mucosal surface to aid more rapid healing and to prevent progression of inflammatory bowel
25 disease. Treatment with the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, is expected to have a significant effect on the production of mucus throughout the gastrointestinal tract and could be used to protect the intestinal mucosa from injurious substances that are ingested or following surgery. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention,
30 could be used to treat diseases associate with the under expression of the polynucleotides of the invention.

Moreover, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to prevent and heal damage to the lungs due to various pathological states. A growth factor such as the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, which could stimulate proliferation and differentiation and promote the repair of alveoli and bronchiolar epithelium to prevent or treat acute or chronic lung damage. For example, emphysema, which results in the progressive loss of aveoli, and inhalation injuries, i.e., resulting from smoke inhalation and burns, that cause necrosis of the bronchiolar epithelium and alveoli could be effectively treated, prevented, and/or diagnosed using the polynucleotides or polypeptides, and/or agonists or antagonists of the invention. Also, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to stimulate the proliferation of and differentiation of type II pneumocytes, which may help treat or prevent disease such as hyaline membrane diseases, such as infant respiratory distress syndrome and bronchopulmonary dysplasia, in premature infants.

The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could stimulate the proliferation and differentiation of hepatocytes and, thus, could be used to alleviate or treat liver diseases and pathologies such as fulminant liver failure caused by cirrhosis, liver damage caused by viral hepatitis and toxic substances (i.e., acetaminophen, carbon tetrachloride and other hepatotoxins known in the art).

In addition, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used treat or prevent the onset of diabetes mellitus. In patients with newly diagnosed Types I and II diabetes, where some islet cell function remains, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to maintain the islet function so as to alleviate, delay or prevent permanent manifestation of the disease. Also, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used as an auxiliary in islet cell transplantation to improve or promote islet cell function.

Neurological Diseases

Nervous system diseases, disorders, and/or conditions, which can be treated, prevented, and/or diagnosed with the compositions of the invention (e.g., polypeptides, polynucleotides, and/or agonists or antagonists), include, but are not limited to, nervous system injuries, and diseases, disorders, and/or conditions which result in either a disconnection of axons, a diminution or degeneration of neurons, or demyelination. Nervous system lesions which may be treated, prevented, and/or diagnosed in a patient (including human and non-human mammalian patients) according to the invention, include but are not limited to, the following lesions of either the central (including spinal cord, brain) or peripheral nervous systems: (1) ischemic lesions, in which a lack of oxygen in a portion of the nervous system results in neuronal injury or death, including cerebral infarction or ischemia, or spinal cord infarction or ischemia; (2) traumatic lesions, including lesions caused by physical injury or associated with surgery, for example, lesions which sever a portion of the nervous system, or compression injuries; (3) malignant lesions, in which a portion of the nervous system is destroyed or injured by malignant tissue which is either a nervous system associated malignancy or a malignancy derived from non-nervous system tissue; (4) infectious lesions, in which a portion of the nervous system is destroyed or injured as a result of infection, for example, by an abscess or associated with infection by human immunodeficiency virus, herpes zoster, or herpes simplex virus or with Lyme disease, tuberculosis, syphilis; (5) degenerative lesions, in which a portion of the nervous system is destroyed or injured as a result of a degenerative process including but not limited to degeneration associated with Parkinson's disease, Alzheimer's disease, Huntington's chorea, or amyotrophic lateral sclerosis (ALS); (6) lesions associated with nutritional diseases, disorders, and/or conditions, in which a portion of the nervous system is destroyed or injured by a nutritional disorder or disorder of metabolism including but not limited to, vitamin B12 deficiency, folic acid deficiency, Wernicke disease, tobacco-alcohol amblyopia, Marchiafava-Bignami disease (primary degeneration of the corpus callosum), and alcoholic cerebellar degeneration; (7) neurological lesions associated with systemic diseases including, but not limited to, diabetes (diabetic neuropathy, Bell's palsy), systemic lupus

erythematosis, carcinoma, or sarcoidosis; (8) lesions caused by toxic substances including alcohol, lead, or particular neurotoxins; and (9) demyelinated lesions in which a portion of the nervous system is destroyed or injured by a demyelinating disease including, but not limited to, multiple sclerosis, human immunodeficiency virus-associated myelopathy, transverse myelopathy or various etiologies, progressive multifocal leukoencephalopathy, and central pontine myelinolysis.

In a preferred embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to protect neural cells from the damaging effects of cerebral hypoxia. According to this embodiment, the compositions of the invention are used to treat, prevent, and/or diagnose neural cell injury associated with cerebral hypoxia. In one aspect of this embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to treat, prevent, and/or diagnose neural cell injury associated with cerebral ischemia. In another aspect of this embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to treat, prevent, and/or diagnose neural cell injury associated with cerebral infarction. In another aspect of this embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to treat, prevent, and/or diagnose or prevent neural cell injury associated with a stroke. In a further aspect of this embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to treat, prevent, and/or diagnose neural cell injury associated with a heart attack.

The compositions of the invention which are useful for treating or preventing a nervous system disorder may be selected by testing for biological activity in promoting the survival or differentiation of neurons. For example, and not by way of limitation, compositions of the invention which elicit any of the following effects may be useful according to the invention: (1) increased survival time of neurons in culture; (2) increased sprouting of neurons in culture or *in vivo*; (3) increased production of a neuron-associated molecule in culture or *in vivo*, *e.g.*, choline acetyltransferase or acetylcholinesterase with respect to motor neurons; or (4) decreased symptoms of neuron dysfunction *in vivo*. Such effects may be measured by any method known in the art. In preferred, non-limiting embodiments, increased survival of neurons may

1 routinely be measured using a method set forth herein or otherwise known in the art,
2 such as, for example, the method set forth in Arakawa et al. (J. Neurosci.
3 10:3507-3515 (1990)); increased sprouting of neurons may be detected by methods
4 known in the art, such as, for example, the methods set forth in Pestronk et al. (Exp.
5 Neurol. 70:65-82 (1980)) or Brown et al. (Ann. Rev. Neurosci. 4:17-42 (1981));
6 increased production of neuron-associated molecules may be measured by bioassay,
7 enzymatic assay, antibody binding, Northern blot assay, etc., using techniques known
8 in the art and depending on the molecule to be measured; and motor neuron
9 dysfunction may be measured by assessing the physical manifestation of motor
10 neuron disorder, e.g., weakness, motor neuron conduction velocity, or functional
11 disability.

12 In specific embodiments, motor neuron diseases, disorders, and/or conditions
13 that may be treated, prevented, and/or diagnosed according to the invention include,
14 but are not limited to, diseases, disorders, and/or conditions such as infarction,
15 infection, exposure to toxin, trauma, surgical damage, degenerative disease or
16 malignancy that may affect motor neurons as well as other components of the nervous
17 system, as well as diseases, disorders, and/or conditions that selectively affect neurons
18 such as amyotrophic lateral sclerosis, and including, but not limited to, progressive
19 spinal muscular atrophy, progressive bulbar palsy, primary lateral sclerosis, infantile
20 and juvenile muscular atrophy, progressive bulbar paralysis of childhood (Fazio-
21 Londe syndrome), poliomyelitis and the post polio syndrome, and Hereditary
22 Motorsensory Neuropathy (Charcot-Marie-Tooth Disease).

Infectious Disease

23 A polypeptide or polynucleotide and/or agonist or antagonist of the present
24 invention can be used to treat, prevent, and/or diagnose infectious agents. For
25 example, by increasing the immune response, particularly increasing the proliferation
26 and differentiation of B and/or T cells, infectious diseases may be treated, prevented,
27 and/or diagnosed. The immune response may be increased by either enhancing an
28 existing immune response, or by initiating a new immune response. Alternatively,
29 polypeptide or polynucleotide and/or agonist or antagonist of the present invention
30

may also directly inhibit the infectious agent, without necessarily eliciting an immune response.

Viruses are one example of an infectious agent that can cause disease or symptoms that can be treated, prevented, and/or diagnosed by a polynucleotide or polypeptide and/or agonist or antagonist of the present invention. Examples of viruses, include, but are not limited to Examples of viruses, include, but are not limited to the following DNA and RNA viruses and viral families: Arbovirus, Adenoviridae, Arenaviridae, Arterivirus, Birnaviridae, Bunyaviridae, Caliciviridae, Circoviridae, Coronaviridae, Dengue, EBV, HIV, Flaviviridae, Hepadnaviridae (Hepatitis), Herpesviridae (such as, Cytomegalovirus, Herpes Simplex, Herpes Zoster), Mononegavirus (e.g., Paramyxoviridae, Morbillivirus, Rhabdoviridae), Orthomyxoviridae (e.g., Influenza A, Influenza B, and parainfluenza), Papilloma virus, Papovaviridae, Parvoviridae, Picornaviridae, Poxviridae (such as Smallpox or Vaccinia), Reoviridae (e.g., Rotavirus), Retroviridae (HTLV-I, HTLV-II, Lentivirus), and Togaviridae (e.g., Rubivirus). Viruses falling within these families can cause a variety of diseases or symptoms, including, but not limited to: arthritis, bronchiolitis, respiratory syncytial virus, encephalitis, eye infections (e.g., conjunctivitis, keratitis), chronic fatigue syndrome, hepatitis (A, B, C, E, Chronic Active, Delta), Japanese B encephalitis, Junin, Chikungunya, Rift Valley fever, yellow fever, meningitis, opportunistic infections (e.g., AIDS), pneumonia, Burkitt's Lymphoma, chickenpox, hemorrhagic fever, Measles, Mumps, Parainfluenza, Rabies, the common cold, Polio, leukemia, Rubella, sexually transmitted diseases, skin diseases (e.g., Kaposi's, warts), and viremia. polynucleotides or polypeptides, or agonists or antagonists of the invention, can be used to treat, prevent, and/or diagnose any of these symptoms or diseases. In specific embodiments, polynucleotides, polypeptides, or agonists or antagonists of the invention are used to treat, prevent, and/or diagnose: meningitis, Dengue, EBV, and/or hepatitis (e.g., hepatitis B). In an additional specific embodiment polynucleotides, polypeptides, or agonists or antagonists of the invention are used to treat patients nonresponsive to one or more other commercially available hepatitis vaccines. In a further specific embodiment polynucleotides, polypeptides, or

agonists or antagonists of the invention are used to treat, prevent, and/or diagnose AIDS.

- Similarly, bacterial or fungal agents that can cause disease or symptoms and that can be treated, prevented, and/or diagnosed by a polynucleotide or polypeptide and/or agonist or antagonist of the present invention include, but not limited to, include, but not limited to, the following Gram-Negative and Gram-positive bacteria and bacterial families and fungi: Actinomycetales (e.g., *Corynebacterium*, *Mycobacterium*, *Nocardia*), *Cryptococcus neoformans*, Aspergillosis, Bacillaceae (e.g., *Anthrax*, *Clostridium*), Bacteroidaceae, Blastomycosis, *Bordetella*, *Borrelia* (e.g., *Borrelia burgdorferi*), Brucellosis, Candidiasis, *Campylobacter*, Coccidioidomycosis, Cryptococcosis, Dermatocycoses, *E. coli* (e.g., Enterotoxigenic *E. coli* and Enterohemorrhagic *E. coli*), Enterobacteriaceae (*Klebsiella*, *Salmonella* (e.g., *Salmonella typhi*, and *Salmonella paratyphi*), *Serratia*, *Yersinia*), *Erysipelothrix*, *Helicobacter*, Legionellosis, Leptospirosis, *Listeria*, Mycoplasmatales, *Mycobacterium leprae*, *Vibrio cholerae*, Neisseriaceae (e.g., *Acinetobacter*, *Gonorrhea*, *Menigococcal*), *Meisseria meningitidis*. Pasteurellacea Infections (e.g., *Actinobacillus*, *Heamophilus* (e.g., *Heamophilus influenza type B*), *Pasteurella*), *Pseudomonas*, Rickettsiaceae, Chlamydiaceae, Syphilis, *Shigella* spp., Staphylococcal, Meningiococcal, Pneumococcal and Streptococcal (e.g., *Streptococcus pneumoniae* and Group B *Streptococcus*). These bacterial or fungal families can cause the following diseases or symptoms, including, but not limited to: bacteremia, endocarditis, eye infections (conjunctivitis, tuberculosis, uveitis), gingivitis, opportunistic infections (e.g., AIDS related infections), paronychia, prosthesis-related infections, Reiter's Disease, respiratory tract infections, such as Whooping Cough or Empyema, sepsis, Lyme Disease, Cat-Scratch Disease, Dysentery, Paratyphoid Fever, food poisoning, Typhoid, pneumonia, *Gonorrhea*, meningitis (e.g., meningitis types A and B), Chlamydia, Syphilis, Diphtheria, Leprosy, Paratuberculosis, Tuberculosis, Lupus, Botulism, gangrene, tetanus, impetigo, Rheumatic Fever, Scarlet Fever, sexually transmitted diseases, skin diseases (e.g., cellulitis, dermatocycoses), toxemia, urinary tract infections, wound infections. Polynucleotides or polypeptides, agonists or antagonists of the invention, can be used

to treat, prevent, and/or diagnose any of these symptoms or diseases. In specific embodiments, polynucleotides, polypeptides, agonists or antagonists of the invention are used to treat, prevent, and/or diagnose: tetanus, Diphtheria, botulism, and/or meningitis type B.

- 5 Moreover, parasitic agents causing disease or symptoms that can be treated, prevented, and/or diagnosed by a polynucleotide or polypeptide and/or agonist or antagonist of the present invention include, but not limited to, the following families or class: Amebiasis, Babesiosis, Coccidiosis, Cryptosporidiosis, Dientamoebiasis, Dourine, Ectoparasitic, Giardiasis, Helminthiasis, Leishmaniasis, Theileriasis,
- 10 Toxoplasmosis, Trypanosomiasis, and Trichomonas and Sporozoans (e.g., Plasmodium virax, Plasmodium falciparum, Plasmodium malariae and Plasmodium ovale). These parasites can cause a variety of diseases or symptoms, including, but not limited to: Scabies, Trombiculiasis, eye infections, intestinal disease (e.g., dysentery, giardiasis), liver disease, lung disease, opportunistic infections (e.g., AIDS
- 15 related), malaria, pregnancy complications, and toxoplasmosis. polynucleotides or polypeptides, or agonists or antagonists of the invention, can be used to treat, prevent, and/or diagnose any of these symptoms or diseases. In specific embodiments, polynucleotides, polypeptides, or agonists or antagonists of the invention are used to treat, prevent, and/or diagnose malaria.
- 20 Preferably, treatment or prevention using a polypeptide or polynucleotide and/or agonist or antagonist of the present invention could either be by administering an effective amount of a polypeptide to the patient, or by removing cells from the patient, supplying the cells with a polynucleotide of the present invention, and returning the engineered cells to the patient (ex vivo therapy). Moreover, the
- 25 polypeptide or polynucleotide of the present invention can be used as an antigen in a vaccine to raise an immune response against infectious disease.

Regeneration

- 30 A polynucleotide or polypeptide and/or agonist or antagonist of the present invention can be used to differentiate, proliferate, and attract cells, leading to the regeneration of tissues. (See, Science 276:59-87 (1997).) The regeneration of tissues

could be used to repair, replace, or protect tissue damaged by congenital defects, trauma (wounds, burns, incisions, or ulcers), age, disease (e.g. osteoporosis, osteoarthritis, periodontal disease, liver failure), surgery, including cosmetic plastic surgery, fibrosis, reperfusion injury, or systemic cytokine damage.

5 Tissues that could be regenerated using the present invention include organs (e.g., pancreas, liver, intestine, kidney, skin, endothelium), muscle (smooth, skeletal or cardiac), vasculature (including vascular and lymphatics), nervous, hematopoietic, and skeletal (bone, cartilage, tendon, and ligament) tissue. Preferably, regeneration occurs without or decreased scarring. Regeneration also may include angiogenesis.

10 Moreover, a polynucleotide or polypeptide and/or agonist or antagonist of the present invention may increase regeneration of tissues difficult to heal. For example, increased tendon/ligament regeneration would quicken recovery time after damage. A polynucleotide or polypeptide and/or agonist or antagonist of the present invention could also be used prophylactically in an effort to avoid damage. Specific diseases
15 that could be treated, prevented, and/or diagnosed include of tendinitis, carpal tunnel syndrome, and other tendon or ligament defects. A further example of tissue regeneration of non-healing wounds includes pressure ulcers, ulcers associated with vascular insufficiency, surgical, and traumatic wounds.

 Similarly, nerve and brain tissue could also be regenerated by using a
20 polynucleotide or polypeptide and/or agonist or antagonist of the present invention to proliferate and differentiate nerve cells. Diseases that could be treated, prevented, and/or diagnosed using this method include central and peripheral nervous system diseases, neuropathies, or mechanical and traumatic diseases, disorders, and/or conditions (e.g., spinal cord disorders, head trauma, cerebrovascular disease, and
25 stroke). Specifically, diseases associated with peripheral nerve injuries, peripheral neuropathy (e.g., resulting from chemotherapy or other medical therapies), localized neuropathies, and central nervous system diseases (e.g., Alzheimer's disease, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome), could all be treated, prevented, and/or diagnosed using the
30 polynucleotide or polypeptide and/or agonist or antagonist of the present invention.

Chemotaxis

A polynucleotide or polypeptide and/or agonist or antagonist of the present invention may have chemotaxis activity. A chemotactic molecule attracts or mobilizes cells (e.g., monocytes, fibroblasts, neutrophils, T-cells, mast cells, eosinophils, epithelial and/or endothelial cells) to a particular site in the body, such as inflammation, infection, or site of hyperproliferation. The mobilized cells can then fight off and/or heal the particular trauma or abnormality.

A polynucleotide or polypeptide and/or agonist or antagonist of the present invention may increase chemotactic activity of particular cells. These chemotactic molecules can then be used to treat, prevent, and/or diagnose inflammation, infection, hyperproliferative diseases, disorders, and/or conditions, or any immune system disorder by increasing the number of cells targeted to a particular location in the body. For example, chemotactic molecules can be used to treat, prevent, and/or diagnose wounds and other trauma to tissues by attracting immune cells to the injured location. Chemotactic molecules of the present invention can also attract fibroblasts, which can be used to treat, prevent, and/or diagnose wounds.

It is also contemplated that a polynucleotide or polypeptide and/or agonist or antagonist of the present invention may inhibit chemotactic activity. These molecules could also be used to treat, prevent, and/or diagnose diseases, disorders, and/or conditions. Thus, a polynucleotide or polypeptide and/or agonist or antagonist of the present invention could be used as an inhibitor of chemotaxis.

Binding Activity

A polypeptide of the present invention may be used to screen for molecules that bind to the polypeptide or for molecules to which the polypeptide binds. The binding of the polypeptide and the molecule may activate (agonist), increase, inhibit (antagonist), or decrease activity of the polypeptide or the molecule bound. Examples of such molecules include antibodies, oligonucleotides, proteins (e.g., receptors), or small molecules.

Preferably, the molecule is closely related to the natural ligand of the polypeptide, e.g., a fragment of the ligand, or a natural substrate, a ligand, a structural or functional mimetic. (See, Coligan et al., Current Protocols in Immunology 1(2):Chapter 5 (1991).) Similarly, the molecule can be closely related to the natural
5 receptor to which the polypeptide binds, or at least, a fragment of the receptor capable of being bound by the polypeptide (e.g., active site). In either case, the molecule can be rationally designed using known techniques.

Preferably, the screening for these molecules involves producing appropriate cells which express the polypeptide, either as a secreted protein or on the cell
10 membrane. Preferred cells include cells from mammals, yeast, *Drosophila*, or *E. coli*. Cells expressing the polypeptide (or cell membrane containing the expressed polypeptide) are then preferably contacted with a test compound potentially containing the molecule to observe binding, stimulation, or inhibition of activity of either the polypeptide or the molecule.

15 The assay may simply test binding of a candidate compound to the polypeptide, wherein binding is detected by a label, or in an assay involving competition with a labeled competitor. Further, the assay may test whether the candidate compound results in a signal generated by binding to the polypeptide.

Alternatively, the assay can be carried out using cell-free preparations,
20 polypeptide/molecule affixed to a solid support, chemical libraries, or natural product mixtures. The assay may also simply comprise the steps of mixing a candidate compound with a solution containing a polypeptide, measuring polypeptide/molecule activity or binding, and comparing the polypeptide/molecule activity or binding to a standard.

25 Preferably, an ELISA assay can measure polypeptide level or activity in a sample (e.g., biological sample) using a monoclonal or polyclonal antibody. The antibody can measure polypeptide level or activity by either binding, directly or indirectly, to the polypeptide or by competing with the polypeptide for a substrate.

Additionally, the receptor to which a polypeptide of the invention binds can be
30 identified by numerous methods known to those of skill in the art, for example, ligand panning and FACS sorting (Coligan, et al., Current Protocols in Immun., 1(2),

Chapter 5, (1991)). For example, expression cloning is employed wherein polyadenylated RNA is prepared from a cell responsive to the polypeptides, for example, NIH3T3 cells which are known to contain multiple receptors for the FGF family proteins, and SC-3 cells, and a cDNA library created from this RNA is divided
5 into pools and used to transfect COS cells or other cells that are not responsive to the polypeptides. Transfected cells which are grown on glass slides are exposed to the polypeptide of the present invention, after they have been labelled. The polypeptides can be labeled by a variety of means including iodination or inclusion of a recognition site for a site-specific protein kinase.

10 Following fixation and incubation, the slides are subjected to autoradiographic analysis. Positive pools are identified and sub-pools are prepared and re-transfected using an iterative sub-pooling and re-screening process, eventually yielding a single clones that encodes the putative receptor.

As an alternative approach for receptor identification, the labeled polypeptides
15 can be photoaffinity linked with cell membrane or extract preparations that express the receptor molecule. Cross-linked material is resolved by PAGE analysis and exposed to X-ray film. The labeled complex containing the receptors of the polypeptides can be excised, resolved into peptide fragments, and subjected to protein microsequencing. The amino acid sequence obtained from microsequencing would
20 be used to design a set of degenerate oligonucleotide probes to screen a cDNA library to identify the genes encoding the putative receptors.

Moreover, the techniques of gene-shuffling, motif-shuffling, exon-shuffling, and/or codon-shuffling (collectively referred to as "DNA shuffling") may be employed to modulate the activities of polypeptides of the invention thereby
25 effectively generating agonists and antagonists of polypeptides of the invention. See generally, U.S. Patent Nos. 5,605,793, 5,811,238, 5,830,721, 5,834,252, and 5,837,458, and Patten, P. A., et al., Curr. Opinion Biotechnol. 8:724-33 (1997); Harayama, S. Trends Biotechnol. 16(2):76-82 (1998); Hansson, L. O., et al., J. Mol. Biol. 287:265-76 (1999); and Lorenzo, M. M. and Blasco, R. Biotechniques
30 24(2):308-13 (1998) (each of these patents and publications are hereby incorporated by reference). In one embodiment, alteration of polynucleotides and corresponding

polypeptides of the invention may be achieved by DNA shuffling. DNA shuffling involves the assembly of two or more DNA segments into a desired polynucleotide sequence of the invention molecule by homologous, or site-specific, recombination. In another embodiment, polynucleotides and corresponding polypeptides of the invention may be altered by being subjected to random mutagenesis by error-prone 5 PCR, random nucleotide insertion or other methods prior to recombination. In another embodiment, one or more components, motifs, sections, parts, domains, fragments, etc., of the polypeptides of the invention may be recombined with one or more components, motifs, sections, parts, domains, fragments, etc. of one or more 10 heterologous molecules. In preferred embodiments, the heterologous molecules are family members. In further preferred embodiments, the heterologous molecule is a growth factor such as, for example, platelet-derived growth factor (PDGF), insulin-like growth factor (IGF-I), transforming growth factor (TGF)-alpha, epidermal growth factor (EGF), fibroblast growth factor (FGF), TGF-beta, bone morphogenetic 15 protein (BMP)-2, BMP-4, BMP-5, BMP-6, BMP-7, activins A and B, decapentaplegic(dpp), 60A, OP-2, dorsalin, growth differentiation factors (GDFs), nodal, MIS, inhibin-alpha, TGF-beta1, TGF-beta2, TGF-beta3, TGF-beta5, and glial-derived neurotrophic factor (GDNF).

Other preferred fragments are biologically active fragments of the 20 polypeptides of the invention. Biologically active fragments are those exhibiting activity similar, but not necessarily identical, to an activity of the polypeptide. The biological activity of the fragments may include an improved desired activity, or a decreased undesirable activity.

Additionally, this invention provides a method of screening compounds to 25 identify those which modulate the action of the polypeptide of the present invention. An example of such an assay comprises combining a mammalian fibroblast cell, a the polypeptide of the present invention, the compound to be screened and 3[H] thymidine under cell culture conditions where the fibroblast cell would normally proliferate. A control assay may be performed in the absence of the compound to be 30 screened and compared to the amount of fibroblast proliferation in the presence of the compound to determine if the compound stimulates proliferation by determining the

uptake of 3[H] thymidine in each case. The amount of fibroblast cell proliferation is measured by liquid scintillation chromatography which measures the incorporation of 3[H] thymidine. Both agonist and antagonist compounds may be identified by this procedure.

5 In another method, a mammalian cell or membrane preparation expressing a receptor for a polypeptide of the present invention is incubated with a labeled polypeptide of the present invention in the presence of the compound. The ability of the compound to enhance or block this interaction could then be measured. Alternatively, the response of a known second messenger system following
10 interaction of a compound to be screened and the receptor is measured and the ability of the compound to bind to the receptor and elicit a second messenger response is measured to determine if the compound is a potential agonist or antagonist. Such second messenger systems include but are not limited to, cAMP guanylate cyclase, ion channels or phosphoinositide hydrolysis.

15 All of these above assays can be used as diagnostic or prognostic markers. The molecules discovered using these assays can be used to treat, prevent, and/or diagnose disease or to bring about a particular result in a patient (e.g., blood vessel growth) by activating or inhibiting the polypeptide/molecule. Moreover, the assays can discover agents which may inhibit or enhance the production of the polypeptides
20 of the invention from suitably manipulated cells or tissues. Therefore, the invention includes a method of identifying compounds which bind to the polypeptides of the invention comprising the steps of: (a) incubating a candidate binding compound with the polypeptide; and (b) determining if binding has occurred. Moreover, the invention includes a method of identifying agonists/antagonists comprising the steps
25 of: (a) incubating a candidate compound with the polypeptide, (b) assaying a biological activity, and (b) determining if a biological activity of the polypeptide has been altered.

Also, one could identify molecules bind a polypeptide of the invention experimentally by using the beta-pleated sheet regions contained in the polypeptide
30 sequence of the protein. Accordingly, specific embodiments of the invention are directed to polynucleotides encoding polypeptides which comprise, or alternatively

consist of, the amino acid sequence of each beta pleated sheet regions in a disclosed polypeptide sequence. Additional embodiments of the invention are directed to polynucleotides encoding polypeptides which comprise, or alternatively consist of, any combination or all of contained in the polypeptide sequences of the invention.

- 5 Additional preferred embodiments of the invention are directed to polypeptides which comprise, or alternatively consist of, the amino acid sequence of each of the beta pleated sheet regions in one of the polypeptide sequences of the invention. Additional embodiments of the invention are directed to polypeptides which comprise, or alternatively consist of, any combination or all of the beta pleated sheet regions in one
- 10 of the polypeptide sequences of the invention.

Targeted Delivery

- In another embodiment, the invention provides a method of delivering compositions to targeted cells expressing a receptor for a polypeptide of the invention,
- 15 or cells expressing a cell bound form of a polypeptide of the invention.

- As discussed herein, polypeptides or antibodies of the invention may be associated with heterologous polypeptides, heterologous nucleic acids, toxins, or prodrugs via hydrophobic, hydrophilic, ionic and/or covalent interactions. In one embodiment, the invention provides a method for the specific delivery of
- 20 compositions of the invention to cells by administering polypeptides of the invention (including antibodies) that are associated with heterologous polypeptides or nucleic acids. In one example, the invention provides a method for delivering a therapeutic protein into the targeted cell. In another example, the invention provides a method for delivering a single stranded nucleic acid (e.g., antisense or ribozymes) or double
- 25 stranded nucleic acid (e.g., DNA that can integrate into the cell's genome or replicate episomally and that can be transcribed) into the targeted cell.

- In another embodiment, the invention provides a method for the specific destruction of cells (e.g., the destruction of tumor cells) by administering polypeptides of the invention (e.g., polypeptides of the invention or antibodies of the invention) in
- 30 association with toxins or cytotoxic prodrugs.

By "toxin" is meant compounds that bind and activate endogenous cytotoxic effector systems, radioisotopes, holotoxins, modified toxins, catalytic subunits of toxins, or any molecules or enzymes not normally present in or on the surface of a cell that under defined conditions cause the cell's death. Toxins that may be used according to the methods of the invention include, but are not limited to, radioisotopes known in the art, compounds such as, for example, antibodies (or complement fixing containing portions thereof) that bind an inherent or induced endogenous cytotoxic effector system, thymidine kinase, endonuclease, RNase, alpha toxin, ricin, abrin, *Pseudomonas* exotoxin A, diphtheria toxin, saporin, momordin, gelonin, pokeweed antiviral protein, alpha-sarcin and cholera toxin. By "cytotoxic prodrug" is meant a non-toxic compound that is converted by an enzyme, normally present in the cell, into a cytotoxic compound. Cytotoxic prodrugs that may be used according to the methods of the invention include, but are not limited to, glutamyl derivatives of benzoic acid mustard alkylating agent, phosphate derivatives of etoposide or mitomycin C, cytosine arabinoside, daunorubisin, and phenoxyacetamide derivatives of doxorubicin.

Drug Screening

Further contemplated is the use of the polypeptides of the present invention, or the polynucleotides encoding these polypeptides, to screen for molecules which modify the activities of the polypeptides of the present invention. Such a method would include contacting the polypeptide of the present invention with a selected compound(s) suspected of having antagonist or agonist activity, and assaying the activity of these polypeptides following binding.

This invention is particularly useful for screening therapeutic compounds by using the polypeptides of the present invention, or binding fragments thereof, in any of a variety of drug screening techniques. The polypeptide or fragment employed in such a test may be affixed to a solid support, expressed on a cell surface, free in solution, or located intracellularly. One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant nucleic acids expressing the polypeptide or fragment. Drugs are screened against such transformed

cells in competitive binding assays. One may measure, for example, the formulation of complexes between the agent being tested and a polypeptide of the present invention.

Thus, the present invention provides methods of screening for drugs or any
5 other agents which affect activities mediated by the polypeptides of the present invention. These methods comprise contacting such an agent with a polypeptide of the present invention or a fragment thereof and assaying for the presence of a complex between the agent and the polypeptide or a fragment thereof, by methods well known
10 in the art. In such a competitive binding assay, the agents to screen are typically labeled. Following incubation, free agent is separated from that present in bound form, and the amount of free or uncomplexed label is a measure of the ability of a particular agent to bind to the polypeptides of the present invention.

Another technique for drug screening provides high throughput screening for compounds having suitable binding affinity to the polypeptides of the present
15 invention, and is described in great detail in European Patent Application 84/03564, published on September 13, 1984, which is incorporated herein by reference herein. Briefly stated, large numbers of different small peptide test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The peptide test compounds are reacted with polypeptides of the present invention and
20 washed. Bound polypeptides are then detected by methods well known in the art. Purified polypeptides are coated directly onto plates for use in the aforementioned drug screening techniques. In addition, non-neutralizing antibodies may be used to capture the peptide and immobilize it on the solid support.

This invention also contemplates the use of competitive drug screening assays
25 in which neutralizing antibodies capable of binding polypeptides of the present invention specifically compete with a test compound for binding to the polypeptides or fragments thereof. In this manner, the antibodies are used to detect the presence of any peptide which shares one or more antigenic epitopes with a polypeptide of the invention.

30

Antisense And Ribozyme (Antagonists)

In specific embodiments, antagonists according to the present invention are nucleic acids corresponding to the sequences contained in SEQ ID NO:X, or the complementary strand thereof, and/or to nucleotide sequences contained a deposited clone. In one embodiment, antisense sequence is generated internally by the
5 organism, in another embodiment, the antisense sequence is separately administered (see, for example, O'Connor, *Neurochem.*, 56:560 (1991). *Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression*, CRC Press, Boca Raton, FL (1988). Antisense technology can be used to control gene expression through antisense DNA or RNA, or through triple-helix formation. Antisense techniques are discussed for
10 example, in Okano, *Neurochem.*, 56:560 (1991); *Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression*, CRC Press, Boca Raton, FL (1988). Triple helix formation is discussed in, for instance, Lee et al., *Nucleic Acids Research*, 6:3073 (1979); Cooney et al., *Science*, 241:456 (1988); and Dervan et al., *Science*, 251:1300 (1991). The methods are based on binding of a polynucleotide to a complementary
15 DNA or RNA.

For example, the use of c-myc and c-myb antisense RNA constructs to inhibit the growth of the non-lymphocytic leukemia cell line HL-60 and other cell lines was previously described. (Wickstrom et al. (1988); Anfossi et al. (1989)). These experiments were performed in vitro by incubating cells with the oligoribonucleotide.
20 A similar procedure for in vivo use is described in WO 91/15580. Briefly, a pair of oligonucleotides for a given antisense RNA is produced as follows: A sequence complimentary to the first 15 bases of the open reading frame is flanked by an EcoRI site on the 5' end and a HindIII site on the 3' end. Next, the pair of oligonucleotides is heated at 90°C for one minute and then annealed in 2X ligation buffer (20mM TRIS
25 HCl pH 7.5, 10mM MgCl₂, 10mM dithiothreitol (DTT) and 0.2 mM ATP) and then ligated to the EcoRI/Hind III site of the retroviral vector PMV7 (WO 91/15580).

For example, the 5' coding portion of a polynucleotide that encodes the mature polypeptide of the present invention may be used to design an antisense RNA oligonucleotide of from about 10 to 40 base pairs in length. A DNA oligonucleotide
30 is designed to be complementary to a region of the gene involved in transcription thereby preventing transcription and the production of the receptor. The antisense

RNA oligonucleotide hybridizes to the mRNA in vivo and blocks translation of the mRNA molecule into receptor polypeptide.

In one embodiment, the antisense nucleic acid of the invention is produced intracellularly by transcription from an exogenous sequence. For example, a vector or
5 a portion thereof, is transcribed, producing an antisense nucleic acid (RNA) of the invention. Such a vector would contain a sequence encoding the antisense nucleic acid of the invention. Such a vector can remain episomal or become chromosomally integrated, as long as it can be transcribed to produce the desired antisense RNA. Such vectors can be constructed by recombinant DNA technology methods standard
10 in the art. Vectors can be plasmid, viral, or others known in the art, used for replication and expression in vertebrate cells. Expression of the sequence encoding a polypeptide of the invention, or fragments thereof, can be by any promoter known in the art to act in vertebrate, preferably human cells. Such promoters can be inducible or constitutive. Such promoters include, but are not limited to, the SV40 early
15 promoter region (Bernoist and Chambon, *Nature*, 29:304-310 (1981)), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto et al., *Cell*, 22:787-797 (1980)), the herpes thymidine promoter (Wagner et al., *Proc. Natl. Acad. Sci. U.S.A.*, 78:1441-1445 (1981)), the regulatory sequences of the metallothionein gene (Brinster et al., *Nature*, 296:39-42 (1982)), etc.

20 The antisense nucleic acids of the invention comprise a sequence complementary to at least a portion of an RNA transcript of a gene of interest. However, absolute complementarity, although preferred, is not required. A sequence "complementary to at least a portion of an RNA," referred to herein, means a sequence having sufficient complementarity to be able to hybridize with the RNA,
25 forming a stable duplex; in the case of double stranded antisense nucleic acids of the invention, a single strand of the duplex DNA may thus be tested, or triplex formation may be assayed. The ability to hybridize will depend on both the degree of complementarity and the length of the antisense nucleic acid. Generally, the larger the hybridizing nucleic acid, the more base mismatches with a RNA sequence of the
30 invention it may contain and still form a stable duplex (or triplex as the case may be).

One skilled in the art can ascertain a tolerable degree of mismatch by use of standard procedures to determine the melting point of the hybridized complex.

Oligonucleotides that are complementary to the 5' end of the message, *e.g.*, the 5' untranslated sequence up to and including the AUG initiation codon, should
5 work most efficiently at inhibiting translation. However, sequences complementary to the 3' untranslated sequences of mRNAs have been shown to be effective at inhibiting translation of mRNAs as well. See generally, Wagner, R., *Nature*, 372:333-335 (1994). Thus, oligonucleotides complementary to either the 5' - or 3' - non- translated, non-coding regions of a polynucleotide sequence of the invention
10 could be used in an antisense approach to inhibit translation of endogenous mRNA. Oligonucleotides complementary to the 5' untranslated region of the mRNA should include the complement of the AUG start codon. Antisense oligonucleotides complementary to mRNA coding regions are less efficient inhibitors of translation but could be used in accordance with the invention. Whether designed to hybridize to the
15 5' -, 3' - or coding region of mRNA, antisense nucleic acids should be at least six nucleotides in length, and are preferably oligonucleotides ranging from 6 to about 50 nucleotides in length. In specific aspects the oligonucleotide is at least 10 nucleotides, at least 17 nucleotides, at least 25 nucleotides or at least 50 nucleotides.

The polynucleotides of the invention can be DNA or RNA or chimeric
20 mixtures or derivatives or modified versions thereof, single-stranded or double-stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone, for example, to improve stability of the molecule, hybridization, etc. The oligonucleotide may include other appended groups such as peptides (*e.g.*, for targeting host cell receptors *in vivo*), or agents facilitating transport across the cell
25 membrane (see, *e.g.*, Letsinger et al., *Proc. Natl. Acad. Sci. U.S.A.* 86:6553-6556 (1989); Lemaitre et al., *Proc. Natl. Acad. Sci.*, 84:648-652 (1987); PCT Publication NO: WO88/09810, published December 15, 1988) or the blood-brain barrier (see, *e.g.*, PCT Publication NO: WO89/10134, published April 25, 1988), hybridization-triggered cleavage agents. (See, *e.g.*, Krol et al., *BioTechniques*, 6:958-976 (1988))
30 or intercalating agents. (See, *e.g.*, Zon, *Pharm. Res.*, 5:539-549 (1988)). To this end, the oligonucleotide may be conjugated to another molecule, *e.g.*, a peptide,

hybridization triggered cross-linking agent, transport agent, hybridization-triggered cleavage agent, etc.

The antisense oligonucleotide may comprise at least one modified base moiety which is selected from the group including, but not limited to, 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xantine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w. and 2,6-diaminopurine.

The antisense oligonucleotide may also comprise at least one modified sugar moiety selected from the group including, but not limited to, arabinose, 2-fluoroarabinose, xylulose, and hexose.

In yet another embodiment, the antisense oligonucleotide comprises at least one modified phosphate backbone selected from the group including, but not limited to, a phosphorothioate, a phosphorodithioate, a phosphoramidothioate, a phosphoramidate, a phosphordiamidate, a methylphosphonate, an alkyl phosphotriester, and a formacetal or analog thereof.

In yet another embodiment, the antisense oligonucleotide is an a-anomeric oligonucleotide. An a-anomeric oligonucleotide forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual b-units, the strands run parallel to each other (Gautier et al., Nucl. Acids Res., 15:6625-6641 (1987)). The oligonucleotide is a 2-O-methylribonucleotide (Inoue et al., Nucl. Acids Res.,

15:6131-6148 (1987)), or a chimeric RNA-DNA analogue (Inoue et al., FEBS Lett. 215:327-330 (1987)).

Polynucleotides of the invention may be synthesized by standard methods known in the art, e.g. by use of an automated DNA synthesizer (such as are
5 commercially available from Biosearch, Applied Biosystems, etc.). As examples, phosphorothioate oligonucleotides may be synthesized by the method of Stein et al. (Nucl. Acids Res., 16:3209 (1988)), methylphosphonate oligonucleotides can be prepared by use of controlled pore glass polymer supports (Sarin et al., Proc. Natl. Acad. Sci. U.S.A., 85:7448-7451 (1988)), etc.

10 While antisense nucleotides complementary to the coding region sequence of the invention could be used, those complementary to the transcribed untranslated region are most preferred.

Potential antagonists according to the invention also include catalytic RNA, or a ribozyme (See, e.g., PCT International Publication WO 90/11364, published
15 October 4, 1990; Sarver et al, Science, 247:1222-1225 (1990). While ribozymes that cleave mRNA at site specific recognition sequences can be used to destroy mRNAs corresponding to the polynucleotides of the invention, the use of hammerhead ribozymes is preferred. Hammerhead ribozymes cleave mRNAs at locations dictated by flanking regions that form complementary base pairs with the target mRNA. The
20 sole requirement is that the target mRNA have the following sequence of two bases: 5' -UG-3'. The construction and production of hammerhead ribozymes is well known in the art and is described more fully in Haseloff and Gerlach, Nature, 334:585-591 (1988). There are numerous potential hammerhead ribozyme cleavage sites within each nucleotide sequence disclosed in the sequence listing. Preferably,
25 the ribozyme is engineered so that the cleavage recognition site is located near the 5' end of the mRNA corresponding to the polynucleotides of the invention; i.e., to increase efficiency and minimize the intracellular accumulation of non-functional mRNA transcripts.

As in the antisense approach, the ribozymes of the invention can be composed
30 of modified oligonucleotides (e.g. for improved stability, targeting, etc.) and should be delivered to cells which express the polynucleotides of the invention in vivo.

DNA constructs encoding the ribozyme may be introduced into the cell in the same manner as described above for the introduction of antisense encoding DNA. A preferred method of delivery involves using a DNA construct "encoding" the ribozyme under the control of a strong constitutive promoter, such as, for example, pol III or pol II promoter, so that transfected cells will produce sufficient quantities of the ribozyme to destroy endogenous messages and inhibit translation. Since ribozymes unlike antisense molecules, are catalytic, a lower intracellular concentration is required for efficiency.

Antagonist/agonist compounds may be employed to inhibit the cell growth and proliferation effects of the polypeptides of the present invention on neoplastic cells and tissues, i.e. stimulation of angiogenesis of tumors, and, therefore, retard or prevent abnormal cellular growth and proliferation, for example, in tumor formation or growth.

The antagonist/agonist may also be employed to prevent hyper-vascular diseases, and prevent the proliferation of epithelial lens cells after extracapsular cataract surgery. Prevention of the mitogenic activity of the polypeptides of the present invention may also be desirable in cases such as restenosis after balloon angioplasty.

The antagonist/agonist may also be employed to prevent the growth of scar tissue during wound healing.

The antagonist/agonist may also be employed to treat, prevent, and/or diagnose the diseases described herein.

Thus, the invention provides a method of treating or preventing diseases, disorders, and/or conditions, including but not limited to the diseases, disorders, and/or conditions listed throughout this application, associated with overexpression of a polynucleotide of the present invention by administering to a patient (a) an antisense molecule directed to the polynucleotide of the present invention, and/or (b) a ribozyme directed to the polynucleotide of the present invention, and/or (b) a ribozyme directed to the polynucleotide of the present invention.

Other Activities

The polypeptide of the present invention, as a result of the ability to stimulate vascular endothelial cell growth, may be employed in treatment for stimulating re-vascularization of ischemic tissues due to various disease conditions such as thrombosis, arteriosclerosis, and other cardiovascular conditions. These polypeptide may also be employed to stimulate angiogenesis and limb regeneration, as discussed above.

The polypeptide may also be employed for treating wounds due to injuries, burns, post-operative tissue repair, and ulcers since they are mitogenic to various cells of different origins, such as fibroblast cells and skeletal muscle cells, and therefore, facilitate the repair or replacement of damaged or diseased tissue.

The polypeptide of the present invention may also be employed stimulate neuronal growth and to treat, prevent, and/or diagnose neuronal damage which occurs in certain neuronal disorders or neuro-degenerative conditions such as Alzheimer's disease, Parkinson's disease, and AIDS-related complex. The polypeptide of the invention may have the ability to stimulate chondrocyte growth, therefore, they may be employed to enhance bone and periodontal regeneration and aid in tissue transplants or bone grafts.

The polypeptide of the present invention may be also be employed to prevent skin aging due to sunburn by stimulating keratinocyte growth.

The polypeptide of the invention may also be employed for preventing hair loss, since FGF family members activate hair-forming cells and promotes melanocyte growth. Along the same lines, the polypeptides of the present invention may be employed to stimulate growth and differentiation of hematopoietic cells and bone marrow cells when used in combination with other cytokines.

The polypeptide of the invention may also be employed to maintain organs before transplantation or for supporting cell culture of primary tissues.

The polypeptide of the present invention may also be employed for inducing tissue of mesodermal origin to differentiate in early embryos.

The polypeptide or polynucleotides and/or agonist or antagonists of the present invention may also increase or decrease the differentiation or proliferation of embryonic stem cells, besides, as discussed above, hematopoietic lineage.

5 The polypeptide or polynucleotides and/or agonist or antagonists of the present invention may also be used to modulate mammalian characteristics, such as body height, weight, hair color, eye color, skin, percentage of adipose tissue, pigmentation, size, and shape (e.g., cosmetic surgery). Similarly, polypeptides or polynucleotides and/or agonist or antagonists of the present invention may be used to modulate mammalian metabolism affecting catabolism, anabolism, processing,
10 utilization, and storage of energy.

Polypeptide or polynucleotides and/or agonist or antagonists of the present invention may be used to change a mammal's mental state or physical state by influencing biorhythms, cardiac rhythms, depression (including depressive diseases, disorders, and/or conditions), tendency for violence, tolerance for pain, reproductive
15 capabilities (preferably by Activin or Inhibin-like activity), hormonal or endocrine levels, appetite, libido, memory, stress, or other cognitive qualities.

Polypeptide or polynucleotides and/or agonist or antagonists of the present invention may also be used as a food additive or preservative, such as to increase or decrease storage capabilities, fat content, lipid, protein, carbohydrate, vitamins,
20 minerals, cofactors or other nutritional components.

Other Preferred Embodiments

Other preferred embodiments of the claimed invention include an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95%
25 identical to a sequence of at least about 50 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1.

Also preferred is a nucleic acid molecule wherein said sequence of contiguous nucleotides is included in the nucleotide sequence of SEQ ID NO:X in the range of positions beginning with the nucleotide at about the position of the 5' Nucleotide of

the Clone Sequence and ending with the nucleotide at about the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X in Table 1.

Also preferred is a nucleic acid molecule wherein said sequence of contiguous nucleotides is included in the nucleotide sequence of SEQ ID NO:X in the range of positions beginning with the nucleotide at about the position of the 5' Nucleotide of the Start Codon and ending with the nucleotide at about the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X in Table 1.

Similarly preferred is a nucleic acid molecule wherein said sequence of contiguous nucleotides is included in the nucleotide sequence of SEQ ID NO:X in the range of positions beginning with the nucleotide at about the position of the 5' Nucleotide of the First Amino Acid of the Signal Peptide and ending with the nucleotide at about the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X in Table 1.

Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least about 150 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X.

Further preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least about 500 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X.

A further preferred embodiment is a nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to the nucleotide sequence of SEQ ID NO:X beginning with the nucleotide at about the position of the 5' Nucleotide of the First Amino Acid of the Signal Peptide and ending with the nucleotide at about the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X in Table 1.

A further preferred embodiment is an isolated nucleic acid molecule comprising a nucleotide sequence, which is at least 95% identical to the complete nucleotide sequence of SEQ ID NO:X.

Also preferred is an isolated nucleic acid molecule which hybridizes under stringent hybridization conditions to a nucleic acid molecule, wherein said nucleic acid molecule which hybridizes does not hybridize under stringent hybridization

conditions to a nucleic acid molecule having a nucleotide sequence consisting of only A residues or of only T residues.

Also preferred is a composition of matter comprising a DNA molecule which comprises a human cDNA clone identified by a cDNA Clone Identifier in Table 1,
5 which DNA molecule is contained in the material deposited with the American Type Culture Collection and given the ATCC Deposit Number shown in Table 1 for said cDNA Clone Identifier.

Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least 50 contiguous
10 nucleotides in the nucleotide sequence of a human cDNA clone identified by a cDNA Clone Identifier in Table 1, which DNA molecule is contained in the deposit given the ATCC Deposit Number shown in Table 1.

Also preferred is an isolated nucleic acid molecule, wherein said sequence of at least 50 contiguous nucleotides is included in the nucleotide sequence of the
15 complete open reading frame sequence encoded by said human cDNA clone.

Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to sequence of at least 150 contiguous nucleotides in the nucleotide sequence encoded by said human cDNA clone.

A further preferred embodiment is an isolated nucleic acid molecule
20 comprising a nucleotide sequence which is at least 95% identical to sequence of at least 500 contiguous nucleotides in the nucleotide sequence encoded by said human cDNA clone.

A further preferred embodiment is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to the complete
25 nucleotide sequence encoded by said human cDNA clone.

A further preferred embodiment is a method for detecting in a biological sample a nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X
30 wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained

in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1; which method comprises a step of comparing a nucleotide sequence of at least one nucleic acid molecule in said sample with a sequence selected from said group and determining whether the sequence of said nucleic acid molecule in said sample is at
5 least 95% identical to said selected sequence.

Also preferred is the above method wherein said step of comparing sequences comprises determining the extent of nucleic acid hybridization between nucleic acid molecules in said sample and a nucleic acid molecule comprising said sequence selected from said group. Similarly, also preferred is the above method wherein said
10 step of comparing sequences is performed by comparing the nucleotide sequence determined from a nucleic acid molecule in said sample with said sequence selected from said group. The nucleic acid molecules can comprise DNA molecules or RNA molecules.

A further preferred embodiment is a method for identifying the species, tissue
15 or cell type of a biological sample which method comprises a step of detecting nucleic acid molecules in said sample, if any, comprising a nucleotide sequence that is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by
20 a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

The method for identifying the species, tissue or cell type of a biological sample can comprise a step of detecting nucleic acid molecules comprising a
25 nucleotide sequence in a panel of at least two nucleotide sequences, wherein at least one sequence in said panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from said group.

Also preferred is a method for diagnosing in a subject a pathological condition associated with abnormal structure or expression of a gene encoding a secreted
30 protein identified in Table 1, which method comprises a step of detecting in a biological sample obtained from said subject nucleic acid molecules, if any,

comprising a nucleotide sequence that is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA
5 Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

The method for diagnosing a pathological condition can comprise a step of detecting nucleic acid molecules comprising a nucleotide sequence in a panel of at least two nucleotide sequences, wherein at least one sequence in said panel is at least
10 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from said group.

Also preferred is a composition of matter comprising isolated nucleic acid molecules wherein the nucleotide sequences of said nucleic acid molecules comprise a panel of at least two nucleotide sequences, wherein at least one sequence in said
15 panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA
20 clone in Table 1. The nucleic acid molecules can comprise DNA molecules or RNA molecules.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 90% identical to a sequence of at least about 10 contiguous amino acids in the amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1.

25 Also preferred is a polypeptide, wherein said sequence of contiguous amino acids is included in the amino acid sequence of SEQ ID NO:Y in the range of positions beginning with the residue at about the position of the First Amino Acid of the Secreted Portion and ending with the residue at about the Last Amino Acid of the Open Reading Frame as set forth for SEQ ID NO:Y in Table 1.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 30 contiguous amino acids in the amino acid sequence of SEQ ID NO:Y.

Further preferred is an isolated polypeptide comprising an amino acid
5 sequence at least 95% identical to a sequence of at least about 100 contiguous amino acids in the amino acid sequence of SEQ ID NO:Y.

Further preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to the complete amino acid sequence of SEQ ID NO:Y.

10 Further preferred is an isolated polypeptide comprising an amino acid sequence at least 90% identical to a sequence of at least about 10 contiguous amino acids in the complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

15 Also preferred is a polypeptide wherein said sequence of contiguous amino acids is included in the amino acid sequence of a secreted portion of the secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

20 Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 30 contiguous amino acids in the amino acid sequence of the secreted portion of the protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

25 Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 100 contiguous amino acids in the amino acid sequence of the secreted portion of the protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

30 Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to the amino acid sequence of the secreted portion of the protein

encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Further preferred is an isolated antibody which binds specifically to a
5 polypeptide comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained
10 in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Further preferred is a method for detecting in a biological sample a polypeptide comprising an amino acid sequence which is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group
15 consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1; which method comprises a step of comparing an amino acid sequence of at least
20 one polypeptide molecule in said sample with a sequence selected from said group and determining whether the sequence of said polypeptide molecule in said sample is at least 90% identical to said sequence of at least 10 contiguous amino acids.

Also preferred is the above method wherein said step of comparing an amino acid sequence of at least one polypeptide molecule in said sample with a sequence
25 selected from said group comprises determining the extent of specific binding of polypeptides in said sample to an antibody which binds specifically to a polypeptide comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in
30 Table 1; and a complete amino acid sequence of a protein encoded by a human cDNA

clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is the above method wherein said step of comparing sequences is performed by comparing the amino acid sequence determined from a polypeptide molecule in said sample with said sequence selected from said group.

Also preferred is a method for identifying the species, tissue or cell type of a biological sample which method comprises a step of detecting polypeptide molecules in said sample, if any, comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is the above method for identifying the species, tissue or cell type of a biological sample, which method comprises a step of detecting polypeptide molecules comprising an amino acid sequence in a panel of at least two amino acid sequences, wherein at least one sequence in said panel is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the above group.

Also preferred is a method for diagnosing in a subject a pathological condition associated with abnormal structure or expression of a gene encoding a secreted protein identified in Table 1, which method comprises a step of detecting in a biological sample obtained from said subject polypeptide molecules comprising an amino acid sequence in a panel of at least two amino acid sequences, wherein at least one sequence in said panel is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

In any of these methods, the step of detecting said polypeptide molecules includes using an antibody.

Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a nucleotide sequence encoding a polypeptide wherein said polypeptide comprises an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is an isolated nucleic acid molecule, wherein said nucleotide sequence encoding a polypeptide has been optimized for expression of said polypeptide in a prokaryotic host.

Also preferred is an isolated nucleic acid molecule, wherein said polypeptide comprises an amino acid sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Further preferred is a method of making a recombinant vector comprising inserting any of the above isolated nucleic acid molecule into a vector. Also preferred is the recombinant vector produced by this method. Also preferred is a method of making a recombinant host cell comprising introducing the vector into a host cell, as well as the recombinant host cell produced by this method.

Also preferred is a method of making an isolated polypeptide comprising culturing this recombinant host cell under conditions such that said polypeptide is expressed and recovering said polypeptide. Also preferred is this method of making an isolated polypeptide, wherein said recombinant host cell is a eukaryotic cell and said polypeptide is a secreted portion of a human secreted protein comprising an amino acid sequence selected from the group consisting of: an amino acid sequence of

SEQ ID NO:Y beginning with the residue at the position of the First Amino Acid of the Secreted Portion of SEQ ID NO:Y wherein Y is an integer set forth in Table 1 and said position of the First Amino Acid of the Secreted Portion of SEQ ID NO:Y is defined in Table 1; and an amino acid sequence of a secreted portion of a protein
5 encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1. The isolated polypeptide produced by this method is also preferred.

Also preferred is a method of treatment of an individual in need of an increased level of a secreted protein activity, which method comprises administering
10 to such an individual a pharmaceutical composition comprising an amount of an isolated polypeptide, polynucleotide, or antibody of the claimed invention effective to increase the level of said protein activity in said individual.

The above-recited applications have uses in a wide variety of hosts. Such hosts include, but are not limited to, human, murine, rabbit, goat, guinea pig, camel,
15 horse, mouse, rat, hamster, pig, micro-pig, chicken, goat, cow, sheep, dog, cat, non-human primate, and human. In specific embodiments, the host is a mouse, rabbit, goat, guinea pig, chicken, rat, hamster, pig, sheep, dog or cat. In preferred embodiments, the host is a mammal. In most preferred embodiments, the host is a human.

20 In specific embodiments of the invention, for each "Contig ID" listed in the fourth column of Table 2, preferably excluded are one or more polynucleotides comprising, or alternatively consisting of, a nucleotide sequence referenced in the fifth column of Table 2 and described by the general formula of a-b, whereas a and b are uniquely determined for the corresponding SEQ ID NO:X referred to in column 3
25 of Table 2. Further specific embodiments are directed to polynucleotide sequences excluding one, two, three, four, or more of the specific polynucleotide sequences referred to in the fifth column of Table 2. In no way is this listing meant to encompass all of the sequences which may be excluded by the general formula, it is just a representative example. All references available through these accessions are hereby
30 incorporated by reference in their entirety.

TABLE 2

Gene No.	cDNA Clone ID	NT SEQ ID NO: X	Contig ID	Public Accession Numbers
2	HMEIJ21	12	767276	R13644, R13953, R40014, R52685, R40014, H07062, H07061, H13318, H43157, H96929, N52992, AA034502, AA054731, AA054738, AA063318, AA074128, AA464554
5	HMEJC96	15	761218	R16862, R55132, R55139, R59547, AA026967, AA026970, AA031692, AA031712, AA055908, AA149827, AA149854
12	HMHBP74	22	765108	R12909, R18967, R36912, R43982, R49254, R49254, R43982, H10040, H78174, AA236377
13	HMIAC52	23	825484	R35395, H20329, H46461, H46982
20	HMIBD93	30	634227	R45361, R45456, R52109, R45456, R45361, R55770, R55849, H17344, R89670, W32915, W35351, W60204
36	HMMBD19	46	607468	T40326, T41189

Having generally described the invention, the same will be more readily understood by reference to the following examples, which are provided by way of illustration and are not intended as limiting.

5

Examples

Example 1: Isolation of a Selected cDNA Clone From the Deposited Sample

Each cDNA clone in a cited ATCC deposit is contained in a plasmid vector. Table 1 identifies the vectors used to construct the cDNA library from which each clone was isolated. In many cases, the vector used to construct the library is a phage vector from which a plasmid has been excised. The table immediately below correlates the related plasmid for each phage vector used in constructing the cDNA library. For example, where a particular clone is identified in Table 1 as being isolated in the vector "Lambda Zap," the corresponding deposited clone is in "pBluescript."

	<u>Vector Used to Construct Library</u>	<u>Corresponding Deposited</u>
	<u>Plasmid</u>	
	Lambda Zap	pBluescript (pBS)
20	Uni-Zap XR	pBluescript (pBS)
	Zap Express	pBK
	lafmid BA	plafmid BA
	pSport1	pSport1
	pCMVSPORT 2.0	pCMVSPORT 2.0
25	pCMVSPORT 3.0	pCMVSPORT 3.0
	pCR [®] 2.1	pCR [®] 2.1

Vectors Lambda Zap (U.S. Patent Nos. 5,128,256 and 5,286,636), Uni-Zap XR (U.S. Patent Nos. 5,128, 256 and 5,286,636), Zap Express (U.S. Patent Nos. 5,128,256 and 5,286,636), pBluescript (pBS) (Short, J. M. et al., Nucleic Acids Res. 16:7583-7600 (1988); Altung-Mees, M. A. and Short, J. M., Nucleic Acids Res. 17:9494 (1989)) and pBK (Altung-Mees, M. A. et al., Strategies 5:58-61 (1992)) are

30

commercially available from Stratagene Cloning Systems, Inc., 11011 N. Torrey Pines Road, La Jolla, CA, 92037. pBS contains an ampicillin resistance gene and pBK contains a neomycin resistance gene. Both can be transformed into E. coli strain XL-1 Blue, also available from Stratagene. pBS comes in 4 forms SK+, SK-, KS+ and KS. The S and K refers to the orientation of the polylinker to the T7 and T3 primer sequences which flank the polylinker region ("S" is for SacI and "K" is for KpnI which are the first sites on each respective end of the linker). "+" or "-" refer to the orientation of the fl origin of replication ("ori"), such that in one orientation, single stranded rescue initiated from the fl ori generates sense strand DNA and in the other, antisense.

Vectors pSport1, pCMVSPORT 2.0 and pCMVSPORT 3.0. were obtained from Life Technologies, Inc., P. O. Box 6009, Gaithersburg, MD 20897. All Sport vectors contain an ampicillin resistance gene and may be transformed into E. coli strain DH10B, also available from Life Technologies. (See, for instance, Gruber, C. E., et al., Focus 15:59 (1993).) Vector lacmid BA (Bento Soares, Columbia University, NY) contains an ampicillin resistance gene and can be transformed into E. coli strain XL-1 Blue. Vector pCR[®]2.1, which is available from Invitrogen, 1600 Faraday Avenue, Carlsbad, CA 92008, contains an ampicillin resistance gene and may be transformed into E. coli strain DH10B, available from Life Technologies. (See, for instance, Clark, J. M., Nuc. Acids Res. 16:9677-9686 (1988) and Mead, D. et al., Bio/Technology 9: (1991).) Preferably, a polynucleotide of the present invention does not comprise the phage vector sequences identified for the particular clone in Table 1, as well as the corresponding plasmid vector sequences designated above.

The deposited material in the sample assigned the ATCC Deposit Number cited in Table 1 for any given cDNA clone also may contain one or more additional plasmids, each comprising a cDNA clone different from that given clone. Thus, deposits sharing the same ATCC Deposit Number contain at least a plasmid for each cDNA clone identified in Table 1. Typically, each ATCC deposit sample cited in Table 1 comprises a mixture of approximately equal amounts (by weight) of about 50 plasmid DNAs, each containing a different cDNA clone; but such a deposit sample

may include plasmids for more or less than 50 cDNA clones, up to about 500 cDNA clones.

Two approaches can be used to isolate a particular clone from the deposited sample of plasmid DNAs cited for that clone in Table 1. First, a plasmid is directly
5 isolated by screening the clones using a polynucleotide probe corresponding to SEQ ID NO:X.

Particularly, a specific polynucleotide with 30-40 nucleotides is synthesized using an Applied Biosystems DNA synthesizer according to the sequence reported. The oligonucleotide is labeled, for instance, with ^{32}P - γ -ATP using T4 polynucleotide
10 kinase and purified according to routine methods. (E.g., Maniatis et al., Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Press, Cold Spring, NY (1982).) The plasmid mixture is transformed into a suitable host, as indicated above (such as XL-1 Blue (Stratagene)) using techniques known to those of skill in the art, such as those provided by the vector supplier or in related publications or patents cited above.
15 The transformants are plated on 1.5% agar plates (containing the appropriate selection agent, e.g., ampicillin) to a density of about 150 transformants (colonies) per plate. These plates are screened using Nylon membranes according to routine methods for bacterial colony screening (e.g., Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd Edit., (1989), Cold Spring Harbor Laboratory Press, pages 1.93 to
20 1.104), or other techniques known to those of skill in the art.

Alternatively, two primers of 17-20 nucleotides derived from both ends of the SEQ ID NO:X (i.e., within the region of SEQ ID NO:X bounded by the 5' NT and the 3' NT of the clone defined in Table 1) are synthesized and used to amplify the desired cDNA using the deposited cDNA plasmid as a template. The polymerase
25 chain reaction is carried out under routine conditions, for instance, in 25 μl of reaction mixture with 0.5 μg of the above cDNA template. A convenient reaction mixture is 1.5-5 mM MgCl_2 , 0.01% (w/v) gelatin, 20 μM each of dATP, dCTP, dGTP, dTTP, 25 pmol of each primer and 0.25 Unit of Taq polymerase. Thirty five cycles of PCR (denaturation at 94 degree C for 1 min; annealing at 55 degree C for 1 min; elongation
30 at 72 degree C for 1 min) are performed with a Perkin-Elmer Cetus automated thermal cycler. The amplified product is analyzed by agarose gel electrophoresis and

Example 2: Isolation of Genomic Clones Corresponding to a Polynucleotide

A human genomic P1 library (Genomic Systems, Inc.) is screened by PCR using primers selected for the cDNA sequence corresponding to SEQ ID NO:X., according to the method described in Example 1. (See also, Sambrook.)

5

Example 3: Tissue Distribution of Polypeptide

Tissue distribution of mRNA expression of polynucleotides of the present invention is determined using protocols for Northern blot analysis, described by, among others, Sambrook et al. For example, a cDNA probe produced by the method described in Example 1 is labeled with P³² using the rediprime™ DNA labeling system (Amersham Life Science), according to manufacturer's instructions. After labeling, the probe is purified using CHROMA SPIN-100™ column (Clontech Laboratories, Inc.), according to manufacturer's protocol number PT1200-1. The purified labeled probe is then used to examine various human tissues for mRNA expression.

10
15

Multiple Tissue Northern (MTN) blots containing various human tissues (H) or human immune system tissues (IM) (Clontech) are examined with the labeled probe using ExpressHyb™ hybridization solution (Clontech) according to manufacturer's protocol number PT1190-1. Following hybridization and washing, the blots are mounted and exposed to film at -70 degree C overnight, and the films developed according to standard procedures.

20

Example 4: Chromosomal Mapping of the Polynucleotides

An oligonucleotide primer set is designed according to the sequence at the 5' end of SEQ ID NO:X. This primer preferably spans about 100 nucleotides. This primer set is then used in a polymerase chain reaction under the following set of conditions : 30 seconds, 95 degree C; 1 minute, 56 degree C; 1 minute, 70 degree C. This cycle is repeated 32 times followed by one 5 minute cycle at 70 degree C. Human, mouse, and hamster DNA is used as template in addition to a somatic cell hybrid panel containing individual chromosomes or chromosome fragments (Bios, Inc). The reactions is analyzed on either 8% polyacrylamide gels or 3.5 % agarose

25
30

gels. Chromosome mapping is determined by the presence of an approximately 100 bp PCR fragment in the particular somatic cell hybrid.

Example 5: Bacterial Expression of a Polypeptide

5 A polynucleotide encoding a polypeptide of the present invention is amplified using PCR oligonucleotide primers corresponding to the 5' and 3' ends of the DNA sequence, as outlined in Example 1, to synthesize insertion fragments. The primers used to amplify the cDNA insert should preferably contain restriction sites, such as BamHI and XbaI, at the 5' end of the primers in order to clone the amplified product
10 into the expression vector. For example, BamHI and XbaI correspond to the restriction enzyme sites on the bacterial expression vector pQE-9. (Qiagen, Inc., Chatsworth, CA). This plasmid vector encodes antibiotic resistance (Amp^r), a bacterial origin of replication (ori), an IPTG-regulatable promoter/operator (P/O), a ribosome binding site (RBS), a 6-histidine tag (6-His), and restriction enzyme cloning
15 sites.

The pQE-9 vector is digested with BamHI and XbaI and the amplified fragment is ligated into the pQE-9 vector maintaining the reading frame initiated at the bacterial RBS. The ligation mixture is then used to transform the E. coli strain M15/rep4 (Qiagen, Inc.) which contains multiple copies of the plasmid pREP4, which
20 expresses the lacI repressor and also confers kanamycin resistance (Kan^r). Transformants are identified by their ability to grow on LB plates and ampicillin/kanamycin resistant colonies are selected. Plasmid DNA is isolated and confirmed by restriction analysis.

Clones containing the desired constructs are grown overnight (O/N) in liquid
25 culture in LB media supplemented with both Amp (100 ug/ml) and Kan (25 ug/ml). The O/N culture is used to inoculate a large culture at a ratio of 1:100 to 1:250. The cells are grown to an optical density 600 (O.D.⁶⁰⁰) of between 0.4 and 0.6. IPTG (Isopropyl-B-D-thiogalacto pyranoside) is then added to a final concentration of 1 mM. IPTG induces by inactivating the lacI repressor, clearing the P/O leading to
30 increased gene expression.

Cells are grown for an extra 3 to 4 hours. Cells are then harvested by centrifugation (20 mins at 6000Xg). The cell pellet is solubilized in the chaotropic agent 6 Molar Guanidine HCl by stirring for 3-4 hours at 4 degree C. The cell debris is removed by centrifugation, and the supernatant containing the polypeptide is loaded
5 onto a nickel-nitrilo-tri-acetic acid ("Ni-NTA") affinity resin column (available from QIAGEN, Inc., *supra*). Proteins with a 6 x His tag bind to the Ni-NTA resin with high affinity and can be purified in a simple one-step procedure (for details see: The QIAexpressionist (1995) QIAGEN, Inc., *supra*).

Briefly, the supernatant is loaded onto the column in 6 M guanidine-HCl, pH
10 8, the column is first washed with 10 volumes of 6 M guanidine-HCl, pH 8, then washed with 10 volumes of 6 M guanidine-HCl pH 6, and finally the polypeptide is eluted with 6 M guanidine-HCl, pH 5.

The purified protein is then renatured by dialyzing it against phosphate-buffered saline (PBS) or 50 mM Na-acetate, pH 6 buffer plus 200 mM NaCl.
15 Alternatively, the protein can be successfully refolded while immobilized on the Ni-NTA column. The recommended conditions are as follows: renature using a linear 6M-1M urea gradient in 500 mM NaCl, 20% glycerol, 20 mM Tris/HCl pH 7.4, containing protease inhibitors. The renaturation should be performed over a period of 1.5 hours or more. After renaturation the proteins are eluted by the addition of 250
20 mM imidazole. Imidazole is removed by a final dialyzing step against PBS or 50 mM sodium acetate pH 6 buffer plus 200 mM NaCl. The purified protein is stored at 4 degree C or frozen at -80 degree C.

In addition to the above expression vector, the present invention further includes an expression vector comprising phage operator and promoter elements
25 operatively linked to a polynucleotide of the present invention, called pHE4a. (ATCC Accession Number 209645, deposited on February 25, 1998.) This vector contains:
1) a neomycinphosphotransferase gene as a selection marker, 2) an E. coli origin of replication, 3) a T5 phage promoter sequence, 4) two lac operator sequences, 5) a Shine-Delgamo sequence, and 6) the lactose operon repressor gene (*lacIq*). The
30 origin of replication (*oriC*) is derived from pUC19 (LTI, Gaithersburg, MD). The promoter sequence and operator sequences are made synthetically.

DNA can be inserted into the pHEa by restricting the vector with NdeI and XbaI, BamHI, XhoI, or Asp718, running the restricted product on a gel, and isolating the larger fragment (the stuffer fragment should be about 310 base pairs). The DNA insert is generated according to the PCR protocol described in Example 1, using PCR
5 primers having restriction sites for NdeI (5' primer) and XbaI, BamHI, XhoI, or Asp718 (3' primer). The PCR insert is gel purified and restricted with compatible enzymes. The insert and vector are ligated according to standard protocols.

The engineered vector could easily be substituted in the above protocol to express protein in a bacterial system.

10

Example 6: Purification of a Polypeptide from an Inclusion Body

The following alternative method can be used to purify a polypeptide expressed in *E. coli* when it is present in the form of inclusion bodies. Unless otherwise specified, all of the following steps are conducted at 4-10 degree C.

15 Upon completion of the production phase of the *E. coli* fermentation, the cell culture is cooled to 4-10 degree C and the cells harvested by continuous centrifugation at 15,000 rpm (Heraeus Sepatech). On the basis of the expected yield of protein per unit weight of cell paste and the amount of purified protein required, an appropriate amount of cell paste, by weight, is suspended in a buffer solution
20 containing 100 mM Tris, 50 mM EDTA, pH 7.4. The cells are dispersed to a homogeneous suspension using a high shear mixer.

The cells are then lysed by passing the solution through a microfluidizer (Microfluidics, Corp. or APV Gaulin, Inc.) twice at 4000-6000 psi. The homogenate is then mixed with NaCl solution to a final concentration of 0.5 M NaCl, followed by
25 centrifugation at 7000 xg for 15 min. The resultant pellet is washed again using 0.5M NaCl, 100 mM Tris, 50 mM EDTA, pH 7.4.

The resulting washed inclusion bodies are solubilized with 1.5 M guanidine hydrochloride (GuHCl) for 2-4 hours. After 7000 xg centrifugation for 15 min., the pellet is discarded and the polypeptide containing supernatant is incubated at 4 degree
30 C overnight to allow further GuHCl extraction.

Following high speed centrifugation (30,000 xg) to remove insoluble particles, the GuHCl solubilized protein is refolded by quickly mixing the GuHCl extract with 20 volumes of buffer containing 50 mM sodium, pH 4.5, 150 mM NaCl, 2 mM EDTA by vigorous stirring. The refolded diluted protein solution is kept at 4 degree C
5 without mixing for 12 hours prior to further purification steps.

To clarify the refolded polypeptide solution, a previously prepared tangential filtration unit equipped with 0.16 um membrane filter with appropriate surface area (e.g., Filtron), equilibrated with 40 mM sodium acetate, pH 6.0 is employed. The filtered sample is loaded onto a cation exchange resin (e.g., Poros HS-50, Perseptive
10 Biosystems). The column is washed with 40 mM sodium acetate, pH 6.0 and eluted with 250 mM, 500 mM, 1000 mM, and 1500 mM NaCl in the same buffer, in a stepwise manner. The absorbance at 280 nm of the effluent is continuously monitored. Fractions are collected and further analyzed by SDS-PAGE.

Fractions containing the polypeptide are then pooled and mixed with 4
15 volumes of water. The diluted sample is then loaded onto a previously prepared set of tandem columns of strong anion (Poros HQ-50, Perseptive Biosystems) and weak anion (Poros CM-20, Perseptive Biosystems) exchange resins. The columns are equilibrated with 40 mM sodium acetate, pH 6.0. Both columns are washed with 40 mM sodium acetate, pH 6.0, 200 mM NaCl. The CM-20 column is then eluted using
20 a 10 column volume linear gradient ranging from 0.2 M NaCl. 50 mM sodium acetate, pH 6.0 to 1.0 M NaCl, 50 mM sodium acetate, pH 6.5. Fractions are collected under constant A_{280} monitoring of the effluent. Fractions containing the polypeptide (determined, for instance, by 16% SDS-PAGE) are then pooled.

The resultant polypeptide should exhibit greater than 95% purity after the
25 above refolding and purification steps. No major contaminant bands should be observed from Commassie blue stained 16% SDS-PAGE gel when 5 ug of purified protein is loaded. The purified protein can also be tested for endotoxin/LPS contamination, and typically the LPS content is less than 0.1 ng/ml according to LAL assays.

Example 7: Cloning and Expression of a Polypeptide in a Baculovirus Expression System

In this example, the plasmid shuttle vector pA2 is used to insert a polynucleotide into a baculovirus to express a polypeptide. This expression vector
5 contains the strong polyhedrin promoter of the *Autographa californica* nuclear polyhedrosis virus (AcMNPV) followed by convenient restriction sites such as BamHI, Xba I and Asp718. The polyadenylation site of the simian virus 40 ("SV40") is used for efficient polyadenylation. For easy selection of recombinant virus, the plasmid contains the beta-galactosidase gene from *E. coli* under control of a weak
10 *Drosophila* promoter in the same orientation, followed by the polyadenylation signal of the polyhedrin gene. The inserted genes are flanked on both sides by viral sequences for cell-mediated homologous recombination with wild-type viral DNA to generate a viable virus that express the cloned polynucleotide.

Many other baculovirus vectors can be used in place of the vector above, such
15 as pAc373, pVL941, and pAcIM1, as one skilled in the art would readily appreciate, as long as the construct provides appropriately located signals for transcription, translation, secretion and the like, including a signal peptide and an in-frame AUG as required. Such vectors are described, for instance, in Luckow et al., *Virology* 170:31-39 (1989).

20 Specifically, the cDNA sequence contained in the deposited clone, including the AUG initiation codon and the naturally associated leader sequence identified in Table 1, is amplified using the PCR protocol described in Example 1. If the naturally occurring signal sequence is used to produce the secreted protein, the pA2 vector does not need a second signal peptide. Alternatively, the vector can be modified (pA2 GP)
25 to include a baculovirus leader sequence, using the standard methods described in Summers et al., "A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures," Texas Agricultural Experimental Station Bulletin No. 1555 (1987).

The amplified fragment is isolated from a 1% agarose gel using a
30 commercially available kit ("GeneClean," BIO 101 Inc., La Jolla, Ca.). The fragment

then is digested with appropriate restriction enzymes and again purified on a 1% agarose gel.

The plasmid is digested with the corresponding restriction enzymes and optionally, can be dephosphorylated using calf intestinal phosphatase, using routine
5 procedures known in the art. The DNA is then isolated from a 1% agarose gel using a commercially available kit ("GeneClean" BIO 101 Inc., La Jolla, Ca.).

The fragment and the dephosphorylated plasmid are ligated together with T4 DNA ligase. *E. coli* HB101 or other suitable *E. coli* hosts such as XL-1 Blue (Stratagene Cloning Systems, La Jolla, CA) cells are transformed with the ligation
10 mixture and spread on culture plates. Bacteria containing the plasmid are identified by digesting DNA from individual colonies and analyzing the digestion product by gel electrophoresis. The sequence of the cloned fragment is confirmed by DNA sequencing.

Five ug of a plasmid containing the polynucleotide is co-transfected with 1.0
15 ug of a commercially available linearized baculovirus DNA ("BaculoGold™ baculovirus DNA", Pharmingen, San Diego, CA), using the lipofection method described by Felgner et al., Proc. Natl. Acad. Sci. USA 84:7413-7417 (1987). One ug of BaculoGold™ virus DNA and 5 ug of the plasmid are mixed in a sterile well of a microtiter plate containing 50 ul of serum-free Grace's medium (Life Technologies
20 Inc., Gaithersburg, MD). Afterwards, 10 ul Lipofectin plus 90 ul Grace's medium are added, mixed and incubated for 15 minutes at room temperature. Then the transfection mixture is added drop-wise to Sf9 insect cells (ATCC CRL 1711) seeded in a 35 mm tissue culture plate with 1 ml Grace's medium without serum. The plate is then incubated for 5 hours at 27 degrees C. The transfection solution is then removed
25 from the plate and 1 ml of Grace's insect medium supplemented with 10% fetal calf serum is added. Cultivation is then continued at 27 degrees C for four days.

After four days the supernatant is collected and a plaque assay is performed, as described by Summers and Smith, *supra*. An agarose gel with "Blue Gal" (Life Technologies Inc., Gaithersburg) is used to allow easy identification and isolation of
30 gal-expressing clones, which produce blue-stained plaques. (A detailed description of a "plaque assay" of this type can also be found in the user's guide for insect cell

culture and baculovirology distributed by Life Technologies Inc., Gaithersburg, page 9-10.) After appropriate incubation, blue stained plaques are picked with the tip of a micropipettor (e.g., Eppendorf). The agar containing the recombinant viruses is then resuspended in a microcentrifuge tube containing 200 ul of Grace's medium and the suspension containing the recombinant baculovirus is used to infect Sf9 cells seeded in 35 mm dishes. Four days later the supernatants of these culture dishes are harvested and then they are stored at 4 degree C.

To verify the expression of the polypeptide, Sf9 cells are grown in Grace's medium supplemented with 10% heat-inactivated FBS. The cells are infected with the recombinant baculovirus containing the polynucleotide at a multiplicity of infection ("MOI") of about 2. If radiolabeled proteins are desired, 6 hours later the medium is removed and is replaced with SF900 II medium minus methionine and cysteine (available from Life Technologies Inc., Rockville, MD). After 42 hours, 5 uCi of ^{35}S -methionine and 5 uCi ^{35}S -cysteine (available from Amersham) are added. The cells are further incubated for 16 hours and then are harvested by centrifugation. The proteins in the supernatant as well as the intracellular proteins are analyzed by SDS-PAGE followed by autoradiography (if radiolabeled).

Microsequencing of the amino acid sequence of the amino terminus of purified protein may be used to determine the amino terminal sequence of the produced protein.

Example 8: Expression of a Polypeptide in Mammalian Cells

The polypeptide of the present invention can be expressed in a mammalian cell. A typical mammalian expression vector contains a promoter element, which mediates the initiation of transcription of mRNA, a protein coding sequence, and signals required for the termination of transcription and polyadenylation of the transcript. Additional elements include enhancers, Kozak sequences and intervening sequences flanked by donor and acceptor sites for RNA splicing. Highly efficient transcription is achieved with the early and late promoters from SV40, the long terminal repeats (LTRs) from Retroviruses, e.g., RSV, HTLV I, HIV I and the early

promoter of the cytomegalovirus (CMV). However, cellular elements can also be used (e.g., the human actin promoter).

Suitable expression vectors for use in practicing the present invention include, for example, vectors such as pSVL and pMSG (Pharmacia, Uppsala, Sweden),
5 pRSVcat (ATCC 37152), pSV2dhfr (ATCC 37146), pBC12MI (ATCC 67109), pCMVSPORT 2.0, and pCMVSPORT 3.0. Mammalian host cells that could be used include, human HeLa, 293, H9 and Jurkat cells, mouse NIH3T3 and C127 cells, Cos 1, Cos 7 and CV1, quail QC1-3 cells, mouse L cells and Chinese hamster ovary (CHO) cells.

10 Alternatively, the polypeptide can be expressed in stable cell lines containing the polynucleotide integrated into a chromosome. The co-transfection with a selectable marker such as dhfr, gpt, neomycin, hygromycin allows the identification and isolation of the transfected cells.

The transfected gene can also be amplified to express large amounts of the
15 encoded protein. The DHFR (dihydrofolate reductase) marker is useful in developing cell lines that carry several hundred or even several thousand copies of the gene of interest. (See, e.g., Alt, F. W., et al., J. Biol. Chem. 253:1357-1370 (1978); Hamlin, J. L. and Ma, C., Biochem. et Biophys. Acta, 1097:107-143 (1990); Page, M. J. and Sydenham, M. A., Biotechnology 9:64-68 (1991).) Another useful selection marker
20 is the enzyme glutamine synthase (GS) (Murphy et al., Biochem J. 227:277-279 (1991); Bebbington et al., Bio/Technology 10:169-175 (1992). Using these markers, the mammalian cells are grown in selective medium and the cells with the highest resistance are selected. These cell lines contain the amplified gene(s) integrated into a chromosome. Chinese hamster ovary (CHO) and NSO cells are often used for the
25 production of proteins.

Derivatives of the plasmid pSV2-dhfr (ATCC Accession No. 37146), the expression vectors pC4 (ATCC Accession No. 209646) and pC6 (ATCC Accession No. 209647) contain the strong promoter (LTR) of the Rous Sarcoma Virus (Cullen et al., Molecular and Cellular Biology, 438-447 (March, 1985)) plus a fragment of the
30 CMV-enhancer (Boshart et al., Cell 41:521-530 (1985).) Multiple cloning sites, e.g., with the restriction enzyme cleavage sites BamHI, XbaI and Asp718, facilitate the

cloning of the gene of interest. The vectors also contain the 3' intron, the polyadenylation and termination signal of the rat preproinsulin gene, and the mouse DHFR gene under control of the SV40 early promoter.

Specifically, the plasmid pC6, for example, is digested with appropriate
5 restriction enzymes and then dephosphorylated using calf intestinal phosphates by procedures known in the art. The vector is then isolated from a 1% agarose gel.

A polynucleotide of the present invention is amplified according to the protocol outlined in Example 1. If the naturally occurring signal sequence is used to produce the secreted protein, the vector does not need a second signal peptide.
10 Alternatively, if the naturally occurring signal sequence is not used, the vector can be modified to include a heterologous signal sequence. (See, e.g., WO 96/34891.)

The amplified fragment is isolated from a 1% agarose gel using a commercially available kit ("GeneClean," BIO 101 Inc., La Jolla, Ca.). The fragment then is digested with appropriate restriction enzymes and again purified on a 1%
15 agarose gel.

The amplified fragment is then digested with the same restriction enzyme and purified on a 1% agarose gel. The isolated fragment and the dephosphorylated vector are then ligated with T4 DNA ligase. *E. coli* HB101 or XL-1 Blue cells are then transformed and bacteria are identified that contain the fragment inserted into plasmid
20 pC6 using, for instance, restriction enzyme analysis.

Chinese hamster ovary cells lacking an active DHFR gene is used for transfection. Five µg of the expression plasmid pC6 a pC4 is cotransfected with 0.5 µg of the plasmid pSVneo using lipofectin (Felgner et al., *supra*). The plasmid pSV2-neo contains a dominant selectable marker, the *neo* gene from Tn5 encoding an
25 enzyme that confers resistance to a group of antibiotics including G418. The cells are seeded in alpha minus MEM supplemented with 1 mg/ml G418. After 2 days, the cells are trypsinized and seeded in hybridoma cloning plates (Greiner, Germany) in alpha minus MEM supplemented with 10, 25, or 50 ng/ml of methotrexate plus 1 mg/ml G418. After about 10-14 days single clones are trypsinized and then seeded in
30 6-well petri dishes or 10 ml flasks using different concentrations of methotrexate (50 nM, 100 nM, 200 nM, 400 nM, 800 nM). Clones growing at the highest

concentrations of methotrexate are then transferred to new 6-well plates containing even higher concentrations of methotrexate (1 μ M, 2 μ M, 5 μ M, 10 mM, 20 mM). The same procedure is repeated until clones are obtained which grow at a concentration of 100 - 200 μ M. Expression of the desired gene product is analyzed, for instance, by SDS-PAGE and Western blot or by reversed phase HPLC analysis.

Example 9: Protein Fusions

The polypeptides of the present invention are preferably fused to other proteins. These fusion proteins can be used for a variety of applications. For example, fusion of the present polypeptides to His-tag, HA-tag, protein A, IgG domains, and maltose binding protein facilitates purification. (See Example 5; see also EP A 394,827; Traunecker. et al., Nature 331:84-86 (1988).) Similarly, fusion to IgG-1, IgG-3, and albumin increases the halflife time in vivo. Nuclear localization signals fused to the polypeptides of the present invention can target the protein to a specific subcellular localization, while covalent heterodimer or homodimers can increase or decrease the activity of a fusion protein. Fusion proteins can also create chimeric molecules having more than one function. Finally, fusion proteins can increase solubility and/or stability of the fused protein compared to the non-fused protein. All of the types of fusion proteins described above can be made by modifying the following protocol, which outlines the fusion of a polypeptide to an IgG molecule, or the protocol described in Example 5.

Briefly, the human Fc portion of the IgG molecule can be PCR amplified, using primers that span the 5' and 3' ends of the sequence described below. These primers also should have convenient restriction enzyme sites that will facilitate cloning into an expression vector, preferably a mammalian expression vector.

For example, if pC4 (Accession No. 209646) is used, the human Fc portion can be ligated into the BamHI cloning site. Note that the 3' BamHI site should be destroyed. Next, the vector containing the human Fc portion is re-restricted with BamHI, linearizing the vector, and a polynucleotide of the present invention, isolated by the PCR protocol described in Example 1, is ligated into this BamHI site. Note

that the polynucleotide is cloned without a stop codon, otherwise a fusion protein will not be produced.

If the naturally occurring signal sequence is used to produce the secreted protein, pC4 does not need a second signal peptide. Alternatively, if the naturally occurring signal sequence is not used, the vector can be modified to include a
 5 heterologous signal sequence. (See, e.g., WO 96/34891.)

Human IgG Fc region:

GGGATCCGGAGCCCAAATCTTCTGACAAACTCACACATGCCACCGTGC
 10 CCAGCACCTGAATTCGAGGGTGCACCGTCAGTCTTCCTCTTCCCCCAAAA
 CCCAAGGACACCCTCATGATCTCCCGGACTCCTGAGGTACATGCGTGGT
 GGTGGACGTAAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGG
 ACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTA
 CAACAGCACGTACCGTGTGGTCAGCGTCCTACCGTCCTGCACCAGGACT
 15 GGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCCA
 ACCCCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAAC
 CACAGGTGTACACCCTGCCCCATCCCGGGATGAGCTGACCAAGAACCAG
 GTCAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCAAGCGACATCGCCGT
 GGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACACTACAAGACCACGCCT
 20 CCCGTGCTGGACTCCGACGGCTCCTTCTTCCTCTACAGCAAGCTCACCGTG
 GACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCA
 TGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGG
 GTAAATGAGTGCGACGGCCGCGACTCTAGAGGAT (SEQ ID NO:1)

25 **Example 10: Production of an Antibody from a Polypeptide**

The antibodies of the present invention can be prepared by a variety of methods. (See, Current Protocols, Chapter 2.) As one example of such methods, cells expressing a polypeptide of the present invention is administered to an animal to induce the production of sera containing polyclonal antibodies. In a preferred
 30 method, a preparation of the secreted protein is prepared and purified to render it

substantially free of natural contaminants. Such a preparation is then introduced into an animal in order to produce polyclonal antisera of greater specific activity.

In the most preferred method, the antibodies of the present invention are monoclonal antibodies (or protein binding fragments thereof). Such monoclonal
5 antibodies can be prepared using hybridoma technology. (Köhler et al., *Nature* 256:495 (1975); Köhler et al., *Eur. J. Immunol.* 6:511 (1976); Köhler et al., *Eur. J. Immunol.* 6:292 (1976); Hammerling et al., in: *Monoclonal Antibodies and T-Cell Hybridomas*, Elsevier, N.Y., pp. 563-681 (1981).) In general, such procedures involve immunizing an animal (preferably a mouse) with polypeptide or, more
10 preferably, with a secreted polypeptide-expressing cell. Such cells may be cultured in any suitable tissue culture medium; however, it is preferable to culture cells in Earle's modified Eagle's medium supplemented with 10% fetal bovine serum (inactivated at about 56 degrees C), and supplemented with about 10 g/l of nonessential amino acids, about 1.000 U/ml of penicillin, and about 100 ug/ml of streptomycin.

15 The splenocytes of such mice are extracted and fused with a suitable myeloma cell line. Any suitable myeloma cell line may be employed in accordance with the present invention; however, it is preferable to employ the parent myeloma cell line (SP2O), available from the ATCC. After fusion, the resulting hybridoma cells are selectively maintained in HAT medium, and then cloned by limiting dilution as
20 described by Wands et al. (*Gastroenterology* 80:225-232 (1981).) The hybridoma cells obtained through such a selection are then assayed to identify clones which secrete antibodies capable of binding the polypeptide.

Alternatively, additional antibodies capable of binding to the polypeptide can be produced in a two-step procedure using anti-idiotypic antibodies. Such a method
25 makes use of the fact that antibodies are themselves antigens, and therefore, it is possible to obtain an antibody which binds to a second antibody. In accordance with this method, protein specific antibodies are used to immunize an animal, preferably a mouse. The splenocytes of such an animal are then used to produce hybridoma cells, and the hybridoma cells are screened to identify clones which produce an antibody
30 whose ability to bind to the protein-specific antibody can be blocked by the polypeptide. Such antibodies comprise anti-idiotypic antibodies to the protein-

specific antibody and can be used to immunize an animal to induce formation of further protein-specific antibodies.

It will be appreciated that Fab and F(ab')₂ and other fragments of the antibodies of the present invention may be used according to the methods disclosed
5 herein. Such fragments are typically produced by proteolytic cleavage, using enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab')₂ fragments). Alternatively, secreted protein-binding fragments can be produced through the application of recombinant DNA technology or through synthetic chemistry.

10 For in vivo use of antibodies in humans, it may be preferable to use "humanized" chimeric monoclonal antibodies. Such antibodies can be produced using genetic constructs derived from hybridoma cells producing the monoclonal antibodies described above. Methods for producing chimeric antibodies are known in the art. (See, for review, Morrison, Science 229:1202 (1985); Oi et al.,
15 BioTechniques 4:214 (1986); Cabilly et al., U.S. Patent No. 4,816,567; Taniguchi et al., EP 171496; Morrison et al., EP 173494; Neuberger et al., WO 8601533; Robinson et al., WO 8702671; Boulianne et al., Nature 312:643 (1984); Neuberger et al., Nature 314:268 (1985).)

20 **Example 11: Production Of Secreted Protein For High-Throughput Screening Assays**

The following protocol produces a supernatant containing a polypeptide to be tested. This supernatant can then be used in the Screening Assays described in Examples 13-20.

25 First, dilute Poly-D-Lysine (644 587 Boehringer-Mannheim) stock solution (1mg/ml in PBS) 1:20 in PBS (w/o calcium or magnesium 17-516F Biowhittaker) for a working solution of 50ug/ml. Add 200 ul of this solution to each well (24 well plates) and incubate at RT for 20 minutes. Be sure to distribute the solution over each well (note: a 12-channel pipetter may be used with tips on every other channel).
30 Aspirate off the Poly-D-Lysine solution and rinse with 1ml PBS (Phosphate Buffered

Saline). The PBS should remain in the well until just prior to plating the cells and plates may be poly-lysine coated in advance for up to two weeks.

Plate 293T cells (do not carry cells past P+20) at 2×10^5 cells/well in .5ml DMEM(Dulbecco's Modified Eagle Medium)(with 4.5 G/L glucose and L-glutamine
 5 (12-604F Biowhittaker))/10% heat inactivated FBS(14-503F Biowhittaker)/1x Penstrep(17-602E Biowhittaker). Let the cells grow overnight.

The next day, mix together in a sterile solution basin: 300 ul Lipofectamine (18324-012 Gibco/BRL) and 5ml Optimem I (31985070 Gibco/BRL)/96-well plate. With a small volume multi-channel pipetter, aliquot approximately 2ug of an
 10 expression vector containing a polynucleotide insert, produced by the methods described in Examples 8 or 9, into an appropriately labeled 96-well round bottom plate. With a multi-channel pipetter, add 50ul of the Lipofectamine/Optimem I mixture to each well. Pipette up and down gently to mix. Incubate at RT 15-45 minutes. After about 20 minutes, use a multi-channel pipetter to add 150ul Optimem
 15 I to each well. As a control, one plate of vector DNA lacking an insert should be transfected with each set of transfections.

Preferably, the transfection should be performed by tag-teaming the following tasks. By tag-teaming, hands on time is cut in half, and the cells do not spend too much time on PBS. First, person A aspirates off the media from four 24-well plates
 20 of cells, and then person B rinses each well with .5-1ml PBS. Person A then aspirates off PBS rinse, and person B, using a 12-channel pipetter with tips on every other channel, adds the 200ul of DNA/Lipofectamine/Optimem I complex to the odd wells first, then to the even wells, to each row on the 24-well plates. Incubate at 37 degrees C for 6 hours.

25 While cells are incubating, prepare appropriate media, either 1%BSA in DMEM with 1x penstrep, or CHO-5 media (116.6 mg/L of CaCl_2 (anhyd); 0.00130 mg/L $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$; 0.050 mg/L of $\text{Fe}(\text{NO}_3)_3 \cdot 9\text{H}_2\text{O}$; 0.417 mg/L of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$; 311.80 mg/L of KCl; 28.64 mg/L of MgCl_2 ; 48.84 mg/L of MgSO_4 ; 6995.50 mg/L of NaCl; 2400.0 mg/L of NaHCO_3 ; 62.50 mg/L of $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$; 71.02 mg/L of
 30 Na_2HPO_4 ; .4320 mg/L of $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$; .002 mg/L of Arachidonic Acid ; 1.022 mg/L of Cholesterol; .070 mg/L of DL-alpha-Tocopherol-Acetate; 0.0520 mg/L of Linoleic

Acid; 0.010 mg/L of Linolenic Acid; 0.010 mg/L of Myristic Acid; 0.010 mg/L of
 Oleic Acid; 0.010 mg/L of Palmitric Acid; 0.010 mg/L of Palmitic Acid; 100 mg/L of
 Pluronic F-68; 0.010 mg/L of Stearic Acid; 2.20 mg/L of Tween 80; 4551 mg/L of D-
 Glucose; 130.85 mg/ml of L- Alanine; 147.50 mg/ml of L-Arginine-HCL; 7.50 mg/ml
 5 of L-Asparagine-H₂O; 6.65 mg/ml of L-Aspartic Acid; 29.56 mg/ml of L-Cystine-
 2HCL-H₂O; 31.29 mg/ml of L-Cystine-2HCL; 7.35 mg/ml of L-Glutamic Acid; 365.0
 mg/ml of L-Glutamine; 18.75 mg/ml of Glycine; 52.48 mg/ml of L-Histidine-HCL-
 H₂O; 106.97 mg/ml of L-Isoleucine; 111.45 mg/ml of L-Leucine; 163.75 mg/ml of L-
 Lysine HCL; 32.34 mg/ml of L-Methionine; 68.48 mg/ml of L-Phenylalanine; 40.0
 10 mg/ml of L-Proline; 26.25 mg/ml of L-Serine; 101.05 mg/ml of L-Threonine; 19.22
 mg/ml of L-Tryptophan; 91.79 mg/ml of L-Tyrosine-2Na-2H₂O; 99.65 mg/ml of L-
 Valine; 0.0035 mg/L of Biotin; 3.24 mg/L of D-Ca Pantothenate; 11.78 mg/L of
 Choline Chloride; 4.65 mg/L of Folic Acid; 15.60 mg/L of i-Inositol; 3.02 mg/L of
 Niacinamide; 3.00 mg/L of Pyridoxal HCL; 0.031 mg/L of Pyridoxine HCL; 0.319
 15 mg/L of Riboflavin; 3.17 mg/L of Thiamine HCL; 0.365 mg/L of Thymidine; and
 0.680 mg/L of Vitamin B₁₂; 25 mM of HEPES Buffer; 2.39 mg/L of Na
 Hypoxanthine; 0.105 mg/L of Lipoic Acid; 0.081 mg/L of Sodium Putrescine-2HCL;
 55.0 mg/L of Sodium Pyruvate; 0.0067 mg/L of Sodium Selenite; 20uM of
 Ethanolamine; 0.122 mg/L of Ferric Citrate; 41.70 mg/L of Methyl-B-Cyclodextrin
 20 complexed with Linoleic Acid; 33.33 mg/L of Methyl-B-Cyclodextrin complexed
 with Oleic Acid; and 10 mg/L of Methyl-B-Cyclodextrin complexed with Retinal)
 with 2mm glutamine and 1x penstrep. (BSA (81-068-3 Bayer) 100gm dissolved in 1L
 DMEM for a 10% BSA stock solution). Filter the media and collect 50 ul for
 endotoxin assay in 15ml polystyrene conical.

25 The transfection reaction is terminated, preferably by tag-teaming, at the end
 of the incubation period. Person A aspirates off the transfection media, while person
 B adds 1.5ml appropriate media to each well. Incubate at 37 degrees C for 45 or 72
 hours depending on the media used: 1%BSA for 45 hours or CHO-5 for 72 hours.

On day four, using a 300ul multichannel pipetter, aliquot 600ul in one 1ml
 30 deep well plate and the remaining supernatant into a 2ml deep well. The supernatants
 from each well can then be used in the assays described in Examples 13-20.

It is specifically understood that when activity is obtained in any of the assays described below using a supernatant, the activity originates from either the polypeptide directly (e.g., as a secreted protein) or by the polypeptide inducing expression of other proteins, which are then secreted into the supernatant. Thus, the invention further provides a method of identifying the protein in the supernatant characterized by an activity in a particular assay.

Example 12: Construction of GAS Reporter Construct

One signal transduction pathway involved in the differentiation and proliferation of cells is called the Jaks-STATs pathway. Activated proteins in the Jaks-STATs pathway bind to gamma activation site "GAS" elements or interferon-sensitive responsive element ("ISRE"), located in the promoter of many genes. The binding of a protein to these elements alter the expression of the associated gene.

GAS and ISRE elements are recognized by a class of transcription factors called Signal Transducers and Activators of Transcription, or "STATs." There are six members of the STATs family. Stat1 and Stat3 are present in many cell types, as is Stat2 (as response to IFN-alpha is widespread). Stat4 is more restricted and is not in many cell types though it has been found in T helper class I, cells after treatment with IL-12. Stat5 was originally called mammary growth factor, but has been found at higher concentrations in other cells including myeloid cells. It can be activated in tissue culture cells by many cytokines.

The STATs are activated to translocate from the cytoplasm to the nucleus upon tyrosine phosphorylation by a set of kinases known as the Janus Kinase ("Jaks") family. Jaks represent a distinct family of soluble tyrosine kinases and include Tyk2, Jak1, Jak2, and Jak3. These kinases display significant sequence similarity and are generally catalytically inactive in resting cells.

The Jaks are activated by a wide range of receptors summarized in the Table below. (Adapted from review by Schidler and Darnell, Ann. Rev. Biochem. 64:621-51 (1995).) A cytokine receptor family, capable of activating Jaks, is divided into two groups: (a) Class 1 includes receptors for IL-2, IL-3, IL-4, IL-6, IL-7, IL-9, IL-11, IL-12, IL-15, Epo, PRL, GH, G-CSF, GM-CSF, LIF, CNTF, and thrombopoietin; and (b)

Class 2 includes IFN- α , IFN- γ , and IL-10. The Class 1 receptors share a conserved cysteine motif (a set of four conserved cysteines and one tryptophan) and a WSXWS motif (a membrane proximal region encoding Trp-Ser-Xxx-Trp-Ser (SEQ ID NO:2)).

Thus, on binding of a ligand to a receptor, Jaks are activated, which in turn
5 activate STATs, which then translocate and bind to GAS elements. This entire process is encompassed in the Jaks-STATs signal transduction pathway.

Therefore, activation of the Jaks-STATs pathway, reflected by the binding of the GAS or the ISRE element, can be used to indicate proteins involved in the proliferation and differentiation of cells. For example, growth factors and cytokines
10 are known to activate the Jaks-STATs pathway. (See Table below.) Thus, by using GAS elements linked to reporter molecules, activators of the Jaks-STATs pathway can be identified.

	<u>Ligand</u>	<u>tyk2</u>	<u>JAKs</u> <u>Jak1</u>	<u>Jak2</u>	<u>Jak3</u>	<u>STATs</u>	<u>GAS(elements) or ISRE</u>
	<u>IFN family</u>						
5	IFN- α /B	+	+	-	-	1,2,3	ISRE
	IFN-g		+	+	-	1	GAS (IRF1>Lys6>IFP)
	Il-10	+	?	?	-	1,3	
	<u>gp130 family</u>						
10	IL-6 (Pleiotrophic)	+	+	+	?	1,3	GAS (IRF1>Lys6>IFP)
	Il-11(Pleiotrophic)	?	+	?	?	1,3	
	OnM(Pleiotrophic)	?	+	+	?	1,3	
	LIF(Pleiotrophic)	?	+	+	?	1,3	
	CNTF(Pleiotrophic)	-/+	+	+	?	1,3	
15	G-CSF(Pleiotrophic)	?	+	?	?	1,3	
	IL-12(Pleiotrophic)	+	-	+	+	1,3	
	<u>g-C family</u>						
	IL-2 (lymphocytes)	-	+	-	+	1,3,5	GAS
20	IL-4 (lymph/myeloid)	-	+	-	+	6	GAS (IRF1 = IFP >>Ly6)(IgH)
	IL-7 (lymphocytes)	-	+	-	+	5	GAS
	IL-9 (lymphocytes)	-	+	-	+	5	GAS
	IL-13 (lymphocyte)	-	+	?	?	6	GAS
	IL-15	?	+	?	+	5	GAS
25	<u>gp140 family</u>						
	IL-3 (myeloid)	-	-	+	-	5	GAS (IRF1>IFP>>Ly6)
	IL-5 (myeloid)	-	-	+	-	5	GAS
	GM-CSF (myeloid)	-	-	+	-	5	GAS
30	<u>Growth hormone family</u>						
	GH	?	-	+	-	5	
	PRL	?	+/-	+	-	1,3,5	
	EPO	?	-	+	-	5	GAS(B-CAS>IRF1=IFP>>Ly6)
35	<u>Receptor Tyrosine Kinases</u>						
	EGF	?	+	+	-	1,3	GAS (IRF1)
	PDGF	?	+	+	-	1,3	
	CSF-1	?	+	+	-	1,3	GAS (not IRF1)
40							

To construct a synthetic GAS containing promoter element, which is used in the Biological Assays described in Examples 13-14, a PCR based strategy is employed to generate a GAS-SV40 promoter sequence. The 5' primer contains four tandem copies of the GAS binding site found in the IRF1 promoter and previously demonstrated to bind STATs upon induction with a range of cytokines (Rothman et al., Immunity 1:457-468 (1994).), although other GAS or ISRE elements can be used instead. The 5' primer also contains 18bp of sequence complementary to the SV40 early promoter sequence and is flanked with an XhoI site. The sequence of the 5' primer is:

5':GCGCCTCGAGATTTCCTCCCGAAATCTAGATTTCCTCCCGAAATGATTTCCTCCGAAATGATTTCCTCCCGAAATATCTGCCATCTCAATTAG:3' (SEQ ID NO:3)

The downstream primer is complementary to the SV40 promoter and is flanked with a Hind III site: 5':GCGGCAAGCTTTTTTGCAAAGCCTAGGC:3' (SEQ ID NO:4)

PCR amplification is performed using the SV40 promoter template present in the B-gal:promoter plasmid obtained from Clontech. The resulting PCR fragment is digested with XhoI/Hind III and subcloned into BLSK2-. (Stratagene.) Sequencing with forward and reverse primers confirms that the insert contains the following sequence:

5':CTCGAGATTTCCTCCCGAAATCTAGATTTCCTCCCGAAATGATTTCCTCCCGAAATGATTTCCTCCCGAAATATCTGCCATCTCAATTAGTCAGCAACCATAGTCCCGCCCTAACTCCGCCCATCCGCCCCTAACTCCGCCCAGTTCCGCCCATTCTCCGCCCCATGGCTGACTAATTTTTTTTATTTATGCAGAGGCCGAGGCCGCCTCGGCCTCTGAGCTATTCCAGAAGTAGTGAGGAGGCTTTTTTGGAGGCCTAGGCTTTTGCAAAAAAGCTT:3' (SEQ ID NO:5)

With this GAS promoter element linked to the SV40 promoter, a GAS:SEAP2 reporter construct is next engineered. Here, the reporter molecule is a secreted alkaline phosphatase, or "SEAP." Clearly, however, any reporter molecule can be instead of SEAP, in this or in any of the other Examples. Well known reporter molecules that can be used instead of SEAP include chloramphenicol

acetyltransferase (CAT), luciferase, alkaline phosphatase, B-galactosidase, green fluorescent protein (GFP), or any protein detectable by an antibody.

The above sequence confirmed synthetic GAS-SV40 promoter element is subcloned into the pSEAP-Promoter vector obtained from Clontech using HindIII and XhoI, effectively replacing the SV40 promoter with the amplified GAS:SV40 promoter element, to create the GAS-SEAP vector. However, this vector does not contain a neomycin resistance gene, and therefore, is not preferred for mammalian expression systems.

Thus, in order to generate mammalian stable cell lines expressing the GAS-SEAP reporter, the GAS-SEAP cassette is removed from the GAS-SEAP vector using SalI and NotI, and inserted into a backbone vector containing the neomycin resistance gene, such as pGFP-1 (Clontech), using these restriction sites in the multiple cloning site, to create the GAS-SEAP/Neo vector. Once this vector is transfected into mammalian cells, this vector can then be used as a reporter molecule for GAS binding as described in Examples 13-14.

Other constructs can be made using the above description and replacing GAS with a different promoter sequence. For example, construction of reporter molecules containing NFK-B and EGR promoter sequences are described in Examples 15 and 16. However, many other promoters can be substituted using the protocols described in these Examples. For instance, SRE, IL-2, NFAT, or Osteocalcin promoters can be substituted, alone or in combination (e.g., GAS/NF-KB/EGR, GAS/NF-KB, IL-2/NFAT, or NF-KB/GAS). Similarly, other cell lines can be used to test reporter construct activity, such as HELA (epithelial), HUVEC (endothelial), Reh (B-cell), Saos-2 (osteoblast), HUVAC (aortic), or Cardiomyocyte.

Example 13: High-Throughput Screening Assay for T-cell Activity.

The following protocol is used to assess T-cell activity by identifying factors, and determining whether supernate containing a polypeptide of the invention proliferates and/or differentiates T-cells. T-cell activity is assessed using the GAS/SEAP/Neo construct produced in Example 12. Thus, factors that increase SEAP activity indicate the ability to activate the Jaks-STATS signal transduction pathway.

The T-cell used in this assay is Jurkat T-cells (ATCC Accession No. TIB-152), although Molt-3 cells (ATCC Accession No. CRL-1552) and Molt-4 cells (ATCC Accession No. CRL-1582) cells can also be used.

Jurkat T-cells are lymphoblastic CD4+ Th1 helper cells. In order to generate
5 stable cell lines, approximately 2 million Jurkat cells are transfected with the GAS-SEAP/neo vector using DMRIE-C (Life Technologies)(transfection procedure described below). The transfected cells are seeded to a density of approximately 20,000 cells per well and transfectants resistant to 1 mg/ml gentamicin selected. Resistant colonies are expanded and then tested for their response to increasing
10 concentrations of interferon gamma. The dose response of a selected clone is demonstrated.

Specifically, the following protocol will yield sufficient cells for 75 wells containing 200 ul of cells. Thus, it is either scaled up, or performed in multiple to generate sufficient cells for multiple 96 well plates. Jurkat cells are maintained in
15 RPMI + 10% serum with 1%Pen-Strep. Combine 2.5 mls of OPTI-MEM (Life Technologies) with 10 ug of plasmid DNA in a T25 flask. Add 2.5 ml OPTI-MEM containing 50 ul of DMRIE-C and incubate at room temperature for 15-45 mins.

During the incubation period, count cell concentration, spin down the required number of cells (10^7 per transfection), and resuspend in OPTI-MEM to a final
20 concentration of 10^7 cells/ml. Then add 1ml of 1×10^7 cells in OPTI-MEM to T25 flask and incubate at 37 degrees C for 6 hrs. After the incubation, add 10 ml of RPMI + 15% serum.

The Jurkat:GAS-SEAP stable reporter lines are maintained in RPMI + 10% serum, 1 mg/ml Gentamicin, and 1% Pen-Strep. These cells are treated with
25 supernatants containing polypeptides of the invention and/or induced polypeptides of the invention as produced by the protocol described in Example 11.

On the day of treatment with the supernatant, the cells should be washed and resuspended in fresh RPMI + 10% serum to a density of 500,000 cells per ml. The exact number of cells required will depend on the number of supernatants being
30 screened. For one 96 well plate, approximately 10 million cells (for 10 plates, 100 million cells) are required.

Transfer the cells to a triangular reservoir boat, in order to dispense the cells into a 96 well dish, using a 12 channel pipette. Using a 12 channel pipette, transfer 200 ul of cells into each well (therefore adding 100,000 cells per well).

After all the plates have been seeded, 50 ul of the supernatants are transferred
5 directly from the 96 well plate containing the supernatants into each well using a 12 channel pipette. In addition, a dose of exogenous interferon gamma (0.1, 1.0, 10 ng) is added to wells H9, H10, and H11 to serve as additional positive controls for the assay.

The 96 well dishes containing Jurkat cells treated with supernatants are placed
10 in an incubator for 48 hrs (note: this time is variable between 48-72 hrs). 35 ul samples from each well are then transferred to an opaque 96 well plate using a 12 channel pipette. The opaque plates should be covered (using sellophane covers) and stored at -20 degrees C until SEAP assays are performed according to Example 17. The plates containing the remaining treated cells are placed at 4 degrees C and serve
15 as a source of material for repeating the assay on a specific well if desired.

As a positive control, 100 Unit/ml interferon gamma can be used which is known to activate Jurkat T cells. Over 30 fold induction is typically observed in the positive control wells.

The above protocol may be used in the generation of both transient, as well as,
20 stable transfected cells, which would be apparent to those of skill in the art.

Example 14: High-Throughput Screening Assay Identifying Myeloid Activity

The following protocol is used to assess myeloid activity by determining whether polypeptides of the invention proliferates and/or differentiates myeloid cells.
25 Myeloid cell activity is assessed using the GAS/SEAP/Neo construct produced in Example 12. Thus, factors that increase SEAP activity indicate the ability to activate the Jaks-STATS signal transduction pathway. The myeloid cell used in this assay is U937, a pre-monocyte cell line, although TF-1, HL60, or KG1 can be used.

To transiently transfect U937 cells with the GAS/SEAP/Neo construct
30 produced in Example 12, a DEAE-Dextran method (Kharbanda et. al., 1994, Cell Growth & Differentiation, 5:259-265) is used. First, harvest 2×10^7 U937 cells and

wash with PBS. The U937 cells are usually grown in RPMI 1640 medium containing 10% heat-inactivated fetal bovine serum (FBS) supplemented with 100 units/ml penicillin and 100 mg/ml streptomycin.

Next, suspend the cells in 1 ml of 20 mM Tris-HCl (pH 7.4) buffer containing
5 0.5 mg/ml DEAE-Dextran, 8 ug GAS-SEAP2 plasmid DNA, 140 mM NaCl, 5 mM KCl, 375 uM Na₂HPO₄·7H₂O, 1 mM MgCl₂, and 675 uM CaCl₂. Incubate at 37 degrees C for 45 min.

Wash the cells with RPMI 1640 medium containing 10% FBS and then resuspend in 10 ml complete medium and incubate at 37 degrees C for 36 hr.

10 The GAS-SEAP/U937 stable cells are obtained by growing the cells in 400 ug/ml G418. The G418-free medium is used for routine growth but every one to two months, the cells should be re-grown in 400 ug/ml G418 for couple of passages.

These cells are tested by harvesting 1×10^8 cells (this is enough for ten 96-well plates assay) and wash with PBS. Suspend the cells in 200 ml above described
15 growth medium, with a final density of 5×10^5 cells/ml. Plate 200 ul cells per well in the 96-well plate (or 1×10^5 cells/well).

Add 50 ul of the supernatant prepared by the protocol described in Example 11. Incubate at 37 degrees C for 48 to 72 hr. As a positive control, 100 Unit/ml interferon gamma can be used which is known to activate U937 cells. Over 30 fold
20 induction is typically observed in the positive control wells. SEAP assay the supernatant according to the protocol described in Example 17.

Example 15: High-Throughput Screening Assay Identifying Neuronal Activity.

When cells undergo differentiation and proliferation, a group of genes are
25 activated through many different signal transduction pathways. One of these genes, EGR1 (early growth response gene 1), is induced in various tissues and cell types upon activation. The promoter of EGR1 is responsible for such induction. Using the EGR1 promoter linked to reporter molecules, activation of cells can be assessed.

Particularly, the following protocol is used to assess neuronal activity in PC12
30 cell lines. PC12 cells (rat pheochromocytoma cells) are known to proliferate and/or differentiate by activation with a number of mitogens, such as TPA (tetradecanoyl

phorbol acetate), NGF (nerve growth factor), and EGF (epidermal growth factor). The EGR1 gene expression is activated during this treatment. Thus, by stably transfecting PC12 cells with a construct containing an EGR promoter linked to SEAP reporter, activation of PC12 cells can be assessed.

5 The EGR/SEAP reporter construct can be assembled by the following protocol. The EGR-1 promoter sequence (-633 to +1)(Sakamoto K et al., Oncogene 6:867-871 (1991)) can be PCR amplified from human genomic DNA using the following primers:

5' GCGCTCGAGGGATGACAGCGATAGAACCCCGG -3' (SEQ ID NO:6)

10 5' GCGAAGCTTCGCGACTCCCCGGATCCGCCTC-3' (SEQ ID NO:7)

Using the GAS:SEAP/Neo vector produced in Example 12, EGR1 amplified product can then be inserted into this vector. Linearize the GAS:SEAP/Neo vector using restriction enzymes XhoI/HindIII, removing the GAS/SV40 stuffer. Restrict the EGR1 amplified product with these same enzymes. Ligate the vector and the EGR1
15 promoter.

To prepare 96 well-plates for cell culture, two mls of a coating solution (1:30 dilution of collagen type I (Upstate Biotech Inc. Cat#08-115) in 30% ethanol (filter sterilized)) is added per one 10 cm plate or 50 ml per well of the 96-well plate, and allowed to air dry for 2 hr.

20 PC12 cells are routinely grown in RPMI-1640 medium (Bio Whittaker) containing 10% horse serum (JRH BIOSCIENCES. Cat. # 12449-78P), 5% heat-inactivated fetal bovine serum (FBS) supplemented with 100 units/ml penicillin and 100 ug/ml streptomycin on a precoated 10 cm tissue culture dish. One to four split is done every three to four days. Cells are removed from the plates by scraping and
25 resuspended with pipetting up and down for more than 15 times.

Transfect the EGR/SEAP/Neo construct into PC12 using the Lipofectamine protocol described in Example 11. EGR-SEAP/PC12 stable cells are obtained by growing the cells in 300 ug/ml G418. The G418-free medium is used for routine growth but every one to two months, the cells should be re-grown in 300 ug/ml G418
30 for couple of passages.

To assay for neuronal activity, a 10 cm plate with cells around 70 to 80% confluent is screened by removing the old medium. Wash the cells once with PBS (Phosphate buffered saline). Then starve the cells in low serum medium (RPMI-1640 containing 1% horse serum and 0.5% FBS with antibiotics) overnight.

- 5 The next morning, remove the medium and wash the cells with PBS. Scrape off the cells from the plate, suspend the cells well in 2 ml low serum medium. Count the cell number and add more low serum medium to reach final cell density as 5×10^5 cells/ml.

- 10 Add 200 ul of the cell suspension to each well of 96-well plate (equivalent to 1×10^5 cells/well). Add 50 ul supernatant produced by Example 11, 37°C for 48 to 72 hr. As a positive control, a growth factor known to activate PC12 cells through EGR can be used, such as 50 ng/ul of Neuronal Growth Factor (NGF). Over fifty-fold induction of SEAP is typically seen in the positive control wells. SEAP assay the supernatant according to Example 17.

15

Example 16: High-Throughput Screening Assay for T-cell Activity

- NF-KB (Nuclear Factor KB) is a transcription factor activated by a wide variety of agents including the inflammatory cytokines IL-1 and TNF, CD30 and CD40, lymphotoxin-alpha and lymphotoxin-beta, by exposure to LPS or thrombin, and by expression of certain viral gene products. As a transcription factor, NF-KB regulates the expression of genes involved in immune cell activation, control of apoptosis (NF- KB appears to shield cells from apoptosis), B and T-cell development, anti-viral and antimicrobial responses, and multiple stress responses.
- 20

- In non-stimulated conditions, NF- KB is retained in the cytoplasm with I-KB (Inhibitor KB). However, upon stimulation, I- KB is phosphorylated and degraded, causing NF- KB to shuttle to the nucleus, thereby activating transcription of target genes. Target genes activated by NF- KB include IL-2, IL-6, GM-CSF, ICAM-1 and class I MHC.
- 25

- Due to its central role and ability to respond to a range of stimuli, reporter constructs utilizing the NF-KB promoter element are used to screen the supernatants produced in Example 11. Activators or inhibitors of NF-KB would be useful in
- 30

treating diseases. For example, inhibitors of NF-KB could be used to treat those diseases related to the acute or chronic activation of NF-KB, such as rheumatoid arthritis.

To construct a vector containing the NF-KB promoter element, a PCR based strategy is employed. The upstream primer contains four tandem copies of the NF-KB binding site (GGGGACTTTCCC) (SEQ ID NO:8), 18 bp of sequence complementary to the 5' end of the SV40 early promoter sequence, and is flanked with an XhoI site:

5':GCGGCCTCGAGGGGACTTTCCCGGGACTTTCCGGGGACTTTCCGGGAC
TTTCCATCCTGCCATCTCAATTAG:3' (SEQ ID NO:9)

The downstream primer is complementary to the 3' end of the SV40 promoter and is flanked with a Hind III site:

5':GCGGCAAGCTTTTTGCAAAGCCTAGGC:3' (SEQ ID NO:4)

PCR amplification is performed using the SV40 promoter template present in the pB-gal:promoter plasmid obtained from Clontech. The resulting PCR fragment is digested with XhoI and Hind III and subcloned into BLSK2-. (Stratagene) Sequencing with the T7 and T3 primers confirms the insert contains the following sequence:

5' : CTCGAGGGGACTTTCCCGGGGACTTTCCGGGGACTTTCCGGGACTTTCCATCTGCCATCTCAATTA
GTCAGCAACCATAGTCCCGCCCCCTAACTCCGCCCATCCCGCCCCCTAACTCCGCCCAGTCCGCCCATTC
TCCGCCCATGGCTGACTAATTTTTTTATTTATGTCAGAGGCCGAGGCCCTCGGCCTCTGAGCTATT
CCAGAAGTAGTGAGGAGGCTTTTTTGAGGCCCTAGGCTTTTGCAAAAAGCTT:3' (SEQ ID
NO:10)

Next, replace the SV40 minimal promoter element present in the pSEAP2-promoter plasmid (Clontech) with this NF-KB/SV40 fragment using XhoI and HindIII. However, this vector does not contain a neomycin resistance gene, and therefore, is not preferred for mammalian expression systems.

In order to generate stable mammalian cell lines, the NF-KB/SV40/SEAP cassette is removed from the above NF-KB/SEAP vector using restriction enzymes SalI and NotI, and inserted into a vector containing neomycin resistance. Particularly,

the NF-KB/SV40/SEAP cassette was inserted into pGFP-1 (Clontech), replacing the GFP gene, after restricting pGFP-1 with SalI and NotI.

Once NF-KB/SV40/SEAP/Neo vector is created, stable Jurkat T-cells are created and maintained according to the protocol described in Example 13. Similarly, the method for assaying supernatants with these stable Jurkat T-cells is also described in Example 13. As a positive control, exogenous TNF alpha (0.1, 1, 10 ng) is added to wells H9, H10, and H11, with a 5-10 fold activation typically observed.

Example 17: Assay for SEAP Activity

As a reporter molecule for the assays described in Examples 13-16, SEAP activity is assayed using the Tropix Phospho-light Kit (Cat. BP-400) according to the following general procedure. The Tropix Phospho-light Kit supplies the Dilution, Assay, and Reaction Buffers used below.

Prime a dispenser with the 2.5x Dilution Buffer and dispense 15 ul of 2.5x dilution buffer into Optiplates containing 35 ul of a supernatant. Seal the plates with a plastic sealer and incubate at 65 degree C for 30 min. Separate the Optiplates to avoid uneven heating.

Cool the samples to room temperature for 15 minutes. Empty the dispenser and prime with the Assay Buffer. Add 50 ml Assay Buffer and incubate at room temperature 5 min. Empty the dispenser and prime with the Reaction Buffer (see the table below). Add 50 ul Reaction Buffer and incubate at room temperature for 20 minutes. Since the intensity of the chemiluminescent signal is time dependent, and it takes about 10 minutes to read 5 plates on luminometer, one should treat 5 plates at each time and start the second set 10 minutes later.

Read the relative light unit in the luminometer. Set H12 as blank, and print the results. An increase in chemiluminescence indicates reporter activity.

Reaction Buffer Formulation:

# of plates	Rxn buffer diluent (ml)	CSPD (ml)
10	60	3
11	65	3.25
12	70	3.5
13	75	3.75

14	80	4
15	85	4.25
16	90	4.5
17	95	4.75
18	100	5
19	105	5.25
20	110	5.5
21	115	5.75
22	120	6
23	125	6.25
24	130	6.5
25	135	6.75
26	140	7
27	145	7.25
28	150	7.5
29	155	7.75
30	160	8
31	165	8.25
32	170	8.5
33	175	8.75
34	180	9
35	185	9.25
36	190	9.5
37	195	9.75
38	200	10
39	205	10.25
40	210	10.5
41	215	10.75
42	220	11
43	225	11.25
44	230	11.5
45	235	11.75
46	240	12
47	245	12.25
48	250	12.5
49	255	12.75
50	260	13

Example 18: High-Throughput Screening Assay Identifying Changes in Small Molecule Concentration and Membrane Permeability

Binding of a ligand to a receptor is known to alter intracellular levels of small molecules, such as calcium, potassium, sodium, and pH, as well as alter membrane potential. These alterations can be measured in an assay to identify supernatants which bind to receptors of a particular cell. Although the following protocol describes an assay for calcium, this protocol can easily be modified to detect changes in potassium, sodium, pH, membrane potential, or any other small molecule which is detectable by a fluorescent probe.

The following assay uses Fluorometric Imaging Plate Reader ("FLIPR") to measure changes in fluorescent molecules (Molecular Probes) that bind small molecules. Clearly, any fluorescent molecule detecting a small molecule can be used instead of the calcium fluorescent molecule, fluo-4 (Molecular Probes, Inc.; catalog no. F-14202), used here.

For adherent cells, seed the cells at 10,000 -20,000 cells/well in a Co-star black 96-well plate with clear bottom. The plate is incubated in a CO₂ incubator for 20 hours. The adherent cells are washed two times in Biotek washer with 200 ul of HBSS (Hank's Balanced Salt Solution) leaving 100 ul of buffer after the final wash.

A stock solution of 1 mg/ml fluo-4 is made in 10% pluronic acid DMSO. To load the cells with fluo-4, 50 ul of 12 ug/ml fluo-4 is added to each well. The plate is incubated at 37 degrees C in a CO₂ incubator for 60 min. The plate is washed four times in the Biotek washer with HBSS leaving 100 ul of buffer.

For non-adherent cells, the cells are spun down from culture media. Cells are re-suspended to $2-5 \times 10^6$ cells/ml with HBSS in a 50-ml conical tube. 4 ul of 1 mg/ml fluo-4 solution in 10% pluronic acid DMSO is added to each ml of cell suspension. The tube is then placed in a 37 degrees C water bath for 30-60 min. The cells are washed twice with HBSS, resuspended to 1×10^6 cells/ml, and dispensed into a microplate, 100 ul/well. The plate is centrifuged at 1000 rpm for 5 min. The plate is then washed once in Denley CellWash with 200 ul, followed by an aspiration step to 100 ul final volume.

For a non-cell based assay, each well contains a fluorescent molecule, such as fluo-4. The supernatant is added to the well, and a change in fluorescence is detected.

To measure the fluorescence of intracellular calcium, the FLIPR is set for the following parameters: (1) System gain is 300-800 mW; (2) Exposure time is 0.4 second; (3) Camera F/stop is F/2; (4) Excitation is 488 nm; (5) Emission is 530 nm; and (6) Sample addition is 50 ul. Increased emission at 530 nm indicates an extracellular signaling event which has resulted in an increase in the intracellular Ca⁺⁺ concentration.

Example 19: High-Throughput Screening Assay Identifying Tyrosine Kinase Activity

The Protein Tyrosine Kinases (PTK) represent a diverse group of transmembrane and cytoplasmic kinases. Within the Receptor Protein Tyrosine Kinase (RPTK) group are receptors for a range of mitogenic and metabolic growth factors including the PDGF, FGF, EGF, NGF, HGF and Insulin receptor subfamilies. In addition there are a large family of RPTKs for which the corresponding ligand is unknown. Ligands for RPTKs include mainly secreted small proteins, but also membrane-bound and extracellular matrix proteins.

Activation of RPTK by ligands involves ligand-mediated receptor dimerization, resulting in transphosphorylation of the receptor subunits and activation of the cytoplasmic tyrosine kinases. The cytoplasmic tyrosine kinases include receptor associated tyrosine kinases of the src-family (e.g., src, yes, lck, lyn, fyn) and non-receptor linked and cytosolic protein tyrosine kinases, such as the Jak family, members of which mediate signal transduction triggered by the cytokine superfamily of receptors (e.g., the Interleukins, Interferons, GM-CSF, and Leptin).

Because of the wide range of known factors capable of stimulating tyrosine kinase activity, the identification of novel human secreted proteins capable of activating tyrosine kinase signal transduction pathways are of interest. Therefore, the following protocol is designed to identify those novel human secreted proteins capable of activating the tyrosine kinase signal transduction pathways.

Seed target cells (e.g., primary keratinocytes) at a density of approximately 25,000 cells per well in a 96 well Loprodyne Silent Screen Plates purchased from Nalge Nunc (Naperville, IL). The plates are sterilized with two 30 minute rinses with 100% ethanol, rinsed with water and dried overnight. Some plates are coated for 2 hr with 100 ml of cell culture grade type I collagen (50 mg/ml), gelatin (2%) or polylysine (50 mg/ml), all of which can be purchased from Sigma Chemicals (St. Louis, MO) or 10% Matrigel purchased from Becton Dickinson (Bedford, MA), or calf serum, rinsed with PBS and stored at 4 degree C. Cell growth on these plates is assayed by seeding 5,000 cells/well in growth medium and indirect quantitation of cell number through use of alamarBlue as described by the manufacturer Alamar

Biosciences, Inc. (Sacramento, CA) after 48 hr. Falcon plate covers #3071 from Becton Dickinson (Bedford, MA) are used to cover the Loprodyne Silent Screen Plates. Falcon Microtest III cell culture plates can also be used in some proliferation experiments.

- 5 To prepare extracts, A431 cells are seeded onto the nylon membranes of Loprodyne plates (20,000/200ml/well) and cultured overnight in complete medium. Cells are quiesced by incubation in serum-free basal medium for 24 hr. After 5-20 minutes treatment with EGF (60ng/ml) or 50 ul of the supernatant produced in Example 11, the medium was removed and 100 ml of extraction buffer ((20 mM
- 10 HEPES pH 7.5, 0.15 M NaCl, 1% Triton X-100, 0.1% SDS, 2 mM Na₃VO₄, 2 mM Na₄P₂O₇ and a cocktail of protease inhibitors (# 1836170) obtained from Boehringer Mannheim (Indianapolis, IN) is added to each well and the plate is shaken on a rotating shaker for 5 minutes at 4 degrees C. The plate is then placed in a vacuum transfer manifold and the extract filtered through the 0.45 mm membrane
- 15 bottoms of each well using house vacuum. Extracts are collected in a 96-well catch/assay plate in the bottom of the vacuum manifold and immediately placed on ice. To obtain extracts clarified by centrifugation, the content of each well, after detergent solubilization for 5 minutes, is removed and centrifuged for 15 minutes at 4 degrees C at 16,000 x g.
- 20 Test the filtered extracts for levels of tyrosine kinase activity. Although many methods of detecting tyrosine kinase activity are known, one method is described here.

Generally, the tyrosine kinase activity of a supernatant is evaluated by determining its ability to phosphorylate a tyrosine residue on a specific substrate (a

25 biotinylated peptide). Biotinylated peptides that can be used for this purpose include PSK1 (corresponding to amino acids 6-20 of the cell division kinase cdc2-p34) and PSK2 (corresponding to amino acids 1-17 of gastrin). Both peptides are substrates for a range of tyrosine kinases and are available from Boehringer Mannheim.

The tyrosine kinase reaction is set up by adding the following components in

30 order. First, add 10ul of 5uM Biotinylated Peptide, then 10ul ATP/Mg₂⁺ (5mM ATP/50mM MgCl₂), then 10ul of 5x Assay Buffer (40mM imidazole hydrochloride,

pH7.3, 40 mM beta-glycerophosphate, 1mM EGTA, 100mM MgCl₂, 5 mM MnCl₂, 0.5 mg/ml BSA), then 5ul of Sodium Vanadate(1mM), and then 5ul of water. Mix the components gently and preincubate the reaction mix at 30 degrees C for 2 min. Initial the reaction by adding 10ul of the control enzyme or the filtered supernatant.

- 5 The tyrosine kinase assay reaction is then terminated by adding 10 ul of 120mM EDTA and place the reactions on ice.

- Tyrosine kinase activity is determined by transferring 50 ul aliquot of reaction mixture to a microtiter plate (MTP) module and incubating at 37 degrees C for 20 min. This allows the streptavidin coated 96 well plate to associate with the
- 10 biotinylated peptide. Wash the MTP module with 300ul/well of PBS four times. Next add 75 ul of anti-phosphotyrosine antibody conjugated to horse radish peroxidase(anti-P-Tyr-POD(0.5u/ml)) to each well and incubate at 37 degrees C for one hour. Wash the well as above.

- Next add 100ul of peroxidase substrate solution (Boehringer Mannheim) and
- 15 incubate at room temperature for at least 5 mins (up to 30 min). Measure the absorbance of the sample at 405 nm by using ELISA reader. The level of bound peroxidase activity is quantitated using an ELISA reader and reflects the level of tyrosine kinase activity.

20 **Example 20: High-Throughput Screening Assay Identifying Phosphorylation Activity**

- As a potential alternative and/or compliment to the assay of protein tyrosine kinase activity described in Example 19, an assay which detects activation (phosphorylation) of major intracellular signal transduction intermediates can also be
- 25 used. For example, as described below one particular assay can detect tyrosine phosphorylation of the Erk-1 and Erk-2 kinases. However, phosphorylation of other molecules, such as Raf, JNK, p38 MAP, Map kinase kinase (MEK), MEK kinase, Src, Muscle specific kinase (MuSK), IRAK, Tec, and Janus, as well as any other phosphoserine, phosphotyrosine, or phosphothreonine molecule, can be detected by
- 30 substituting these molecules for Erk-1 or Erk-2 in the following assay.

Specifically, assay plates are made by coating the wells of a 96-well ELISA plate with 0.1ml of protein G (1ug/ml) for 2 hr at room temp, (RT). The plates are then rinsed with PBS and blocked with 3% BSA/PBS for 1 hr at RT. The protein G plates are then treated with 2 commercial monoclonal antibodies (100ng/well) against
5 Erk-1 and Erk-2 (1 hr at RT) (Santa Cruz Biotechnology). (To detect other molecules, this step can easily be modified by substituting a monoclonal antibody detecting any of the above described molecules.) After 3-5 rinses with PBS, the plates are stored at 4 degrees C until use.

A431 cells are seeded at 20,000/well in a 96-well Loprodyne filterplate and
10 cultured overnight in growth medium. The cells are then starved for 48 hr in basal medium (DMEM) and then treated with EGF (6ng/well) or 50 ul of the supernatants obtained in Example 11 for 5-20 minutes. The cells are then solubilized and extracts filtered directly into the assay plate.

After incubation with the extract for 1 hr at RT, the wells are again rinsed. As
15 a positive control, a commercial preparation of MAP kinase (10ng/well) is used in place of A431 extract. Plates are then treated with a commercial polyclonal (rabbit) antibody (1ug/ml) which specifically recognizes the phosphorylated epitope of the Erk-1 and Erk-2 kinases (1 hr at RT). This antibody is biotinylated by standard procedures. The bound polyclonal antibody is then quantitated by successive
20 incubations with Europium-streptavidin and Europium fluorescence enhancing reagent in the Wallac DELFIA instrument (time-resolved fluorescence). An increased fluorescent signal over background indicates a phosphorylation.

Example 21: Method of Determining Alterations in a Gene Corresponding to a
25 **Polynucleotide**

RNA isolated from entire families or individual patients presenting with a phenotype of interest (such as a disease) is be isolated. cDNA is then generated from these RNA samples using protocols known in the art. (See, Sambrook.) The cDNA is then used as a template for PCR, employing primers surrounding regions of interest
30 in SEQ ID NO:X. Suggested PCR conditions consist of 35 cycles at 95 degrees C for

30 seconds; 60-120 seconds at 52-58 degrees C; and 60-120 seconds at 70 degrees C, using buffer solutions described in Sidransky et al., Science 252:706 (1991).

PCR products are then sequenced using primers labeled at their 5' end with T4 polynucleotide kinase, employing SequiTherm Polymerase. (Epicentre
5 Technologies). The intron-exon borders of selected exons is also determined and genomic PCR products analyzed to confirm the results. PCR products harboring suspected mutations is then cloned and sequenced to validate the results of the direct sequencing.

PCR products is cloned into T-tailed vectors as described in Holton et al.,
10 Nucleic Acids Research, 19:1156 (1991) and sequenced with T7 polymerase (United States Biochemical). Affected individuals are identified by mutations not present in unaffected individuals.

Genomic rearrangements are also observed as a method of determining alterations in a gene corresponding to a polynucleotide. Genomic clones isolated
15 according to Example 2 are nick-translated with digoxigenindeoxy-uridine 5'-triphosphate (Boehringer Mannheim), and FISH performed as described in Johnson et al., Methods Cell Biol. 35:73-99 (1991). Hybridization with the labeled probe is carried out using a vast excess of human cot-1 DNA for specific hybridization to the corresponding genomic locus.

20 Chromosomes are counterstained with 4,6-diamino-2-phenylidole and propidium iodide, producing a combination of C- and R-bands. Aligned images for precise mapping are obtained using a triple-band filter set (Chroma Technology, Brattleboro, VT) in combination with a cooled charge-coupled device camera (Photometrics, Tucson, AZ) and variable excitation wavelength filters. (Johnson et
25 al., Genet. Anal. Tech. Appl., 8:75 (1991).) Image collection, analysis and chromosomal fractional length measurements are performed using the ISee Graphical Program System. (Inovision Corporation, Durham, NC.) Chromosome alterations of the genomic region hybridized by the probe are identified as insertions, deletions, and translocations. These alterations are used as a diagnostic marker for an associated
30 disease.

Example 22: Method of Detecting Abnormal Levels of a Polypeptide in a Biological Sample

A polypeptide of the present invention can be detected in a biological sample, and if an increased or decreased level of the polypeptide is detected, this polypeptide
5 is a marker for a particular phenotype. Methods of detection are numerous, and thus, it is understood that one skilled in the art can modify the following assay to fit their particular needs.

For example, antibody-sandwich ELISAs are used to detect polypeptides in a sample, preferably a biological sample. Wells of a microtiter plate are coated with
10 specific antibodies, at a final concentration of 0.2 to 10 ug/ml. The antibodies are either monoclonal or polyclonal and are produced by the method described in Example 10. The wells are blocked so that non-specific binding of the polypeptide to the well is reduced.

The coated wells are then incubated for > 2 hours at RT with a sample
15 containing the polypeptide. Preferably, serial dilutions of the sample should be used to validate results. The plates are then washed three times with deionized or distilled water to remove unbounded polypeptide.

Next, 50 ul of specific antibody-alkaline phosphatase conjugate, at a concentration of 25-400 ng, is added and incubated for 2 hours at room temperature.
20 The plates are again washed three times with deionized or distilled water to remove unbounded conjugate.

Add 75 ul of 4-methylumbelliferyl phosphate (MUP) or p-nitrophenyl phosphate (NPP) substrate solution to each well and incubate 1 hour at room temperature. Measure the reaction by a microtiter plate reader. Prepare a standard
25 curve, using serial dilutions of a control sample, and plot polypeptide concentration on the X-axis (log scale) and fluorescence or absorbance of the Y-axis (linear scale). Interpolate the concentration of the polypeptide in the sample using the standard curve.

Example 23: Formulation

The invention also provides methods of treatment and/or prevention diseases, disorders, and/or conditions (such as, for example, any one or more of the diseases or disorders disclosed herein) by administration to a subject of an effective amount of a
5 Therapeutic. By therapeutic is meant a polynucleotides or polypeptides of the invention (including fragments and variants), agonists or antagonists thereof, and/or antibodies thereto, in combination with a pharmaceutically acceptable carrier type (e.g., a sterile carrier).

The Therapeutic will be formulated and dosed in a fashion consistent with
10 good medical practice, taking into account the clinical condition of the individual patient (especially the side effects of treatment with the Therapeutic alone), the site of delivery, the method of administration, the scheduling of administration, and other factors known to practitioners. The "effective amount" for purposes herein is thus determined by such considerations.

As a general proposition, the total pharmaceutically effective amount of the
15 Therapeutic administered parenterally per dose will be in the range of about 1 ug/kg/day to 10 mg/kg/day of patient body weight, although, as noted above, this will be subject to therapeutic discretion. More preferably, this dose is at least 0.01 mg/kg/day, and most preferably for humans between about 0.01 and 1 mg/kg/day for
20 the hormone. If given continuously, the Therapeutic is typically administered at a dose rate of about 1 ug/kg/hour to about 50 ug/kg/hour, either by 1-4 injections per day or by continuous subcutaneous infusions, for example, using a mini-pump. An intravenous bag solution may also be employed. The length of treatment needed to observe changes and the interval following treatment for responses to occur appears
25 to vary depending on the desired effect.

Therapeutics can be are administered orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, gels, drops or transdermal patch), buccally, or as an oral or nasal spray.
"Pharmaceutically acceptable carrier" refers to a non-toxic solid, semisolid or liquid
30 filler, diluent, encapsulating material or formulation auxiliary of any. The term "parenteral" as used herein refers to modes of administration which include

intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

Therapeutics of the invention are also suitably administered by sustained-release systems. Suitable examples of sustained-release Therapeutics are administered orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, gels, drops or transdermal patch), buccally, or as an oral or nasal spray. "Pharmaceutically acceptable carrier" refers to a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

Therapeutics of the invention are also suitably administered by sustained-release systems. Suitable examples of sustained-release Therapeutics include suitable polymeric materials (such as, for example, semi-permeable polymer matrices in the form of shaped articles, e.g., films, or microcapsules), suitable hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, and sparingly soluble derivatives (such as, for example, a sparingly soluble salt).

Sustained-release matrices include polylactides (U.S. Pat. No. 3,773,919, EP 58,481), copolymers of L-glutamic acid and gamma-ethyl-L-glutamate (Sidman et al., Biopolymers 22:547-556 (1983)), poly (2-hydroxyethyl methacrylate) (Langer et al., J. Biomed. Mater. Res. 15:167-277 (1981), and Langer, Chem. Tech. 12:98-105 (1982)), ethylene vinyl acetate (Langer et al., Id.) or poly-D-(-)-3-hydroxybutyric acid (EP 133,988).

Sustained-release Therapeutics also include liposomally entrapped Therapeutics of the invention (see generally, Langer, *Science* 249:1527-1533 (1990); Treat et al., in *Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 317-327 and 353-365 (1989)). Liposomes containing the Therapeutic are prepared by methods known per se: DE 3,218,121; Epstein et al., Proc. Natl. Acad. Sci. (USA) 82:3688-3692 (1985); Hwang et al., Proc. Natl. Acad. Sci. (USA) 77:4030-4034 (1980); EP 52,322; EP 36,676; EP 88,046; EP 143,949; EP 142,641; Japanese Pat. Appl. 83-118008; U.S. Pat. Nos.

4,485,045 and 4,544,545; and EP 102.324. Ordinarily, the liposomes are of the small (about 200-800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol. percent cholesterol, the selected proportion being adjusted for the optimal Therapeutic.

5 In yet an additional embodiment, the Therapeutics of the invention are delivered by way of a pump (*see* Langer, *supra*; Sefton, CRC Crit. Ref. Biomed. Eng. 14:201 (1987); Buchwald et al., Surgery 88:507 (1980); Saudek et al., N. Engl. J. Med. 321:574 (1989)).

Other controlled release systems are discussed in the review by Langer
10 (*Science* 249:1527-1533 (1990)).

For parenteral administration, in one embodiment, the Therapeutic is formulated generally by mixing it at the desired degree of purity, in a unit dosage injectable form (solution, suspension, or emulsion), with a pharmaceutically acceptable carrier, i.e., one that is non-toxic to recipients at the dosages and
15 concentrations employed and is compatible with other ingredients of the formulation. For example, the formulation preferably does not include oxidizing agents and other compounds that are known to be deleterious to the Therapeutic.

Generally, the formulations are prepared by contacting the Therapeutic uniformly and intimately with liquid carriers or finely divided solid carriers or both.
20 Then, if necessary, the product is shaped into the desired formulation. Preferably the carrier is a parenteral carrier, more preferably a solution that is isotonic with the blood of the recipient. Examples of such carrier vehicles include water, saline, Ringer's solution, and dextrose solution. Non-aqueous vehicles such as fixed oils and ethyl oleate are also useful herein, as well as liposomes.

25 The carrier suitably contains minor amounts of additives such as substances that enhance isotonicity and chemical stability. Such materials are non-toxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, succinate, acetic acid, and other organic acids or their salts; antioxidants such as ascorbic acid; low molecular weight (less than about ten
30 residues) polypeptides, e.g., polyarginine or tripeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as

polyvinylpyrrolidone; amino acids, such as glycine, glutamic acid, aspartic acid, or arginine; monosaccharides, disaccharides, and other carbohydrates including cellulose or its derivatives, glucose, manose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; counterions such as sodium; and/or nonionic
5 surfactants such as polysorbates, poloxamers, or PEG.

The Therapeutic is typically formulated in such vehicles at a concentration of about 0.1 mg/ml to 100 mg/ml, preferably 1-10 mg/ml, at a pH of about 3 to 8. It will be understood that the use of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of polypeptide salts.

10 Any pharmaceutical used for therapeutic administration can be sterile. Sterility is readily accomplished by filtration through sterile filtration membranes (e.g., 0.2 micron membranes). Therapeutics generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

15 Therapeutics ordinarily will be stored in unit or multi-dose containers, for example, sealed ampoules or vials, as an aqueous solution or as a lyophilized formulation for reconstitution. As an example of a lyophilized formulation, 10-ml vials are filled with 5 ml of sterile-filtered 1% (w/v) aqueous Therapeutic solution, and the resulting mixture is lyophilized. The infusion solution is prepared by
20 reconstituting the lyophilized Therapeutic using bacteriostatic Water-for-Injection.

The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the Therapeutics of the invention. Associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or
25 biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration. In addition, the Therapeutics may be employed in conjunction with other therapeutic compounds.

The Therapeutics of the invention may be administered alone or in combination with adjuvants. Adjuvants that may be administered with the
30 Therapeutics of the invention include, but are not limited to, alum, alum plus deoxycholate (ImmunoAg), MTP-PE (Biocine Corp.), QS21 (Genentech, Inc.), BCG,

and MPL. In a specific embodiment, Therapeutics of the invention are administered in combination with alum. In another specific embodiment, Therapeutics of the invention are administered in combination with QS-21. Further adjuvants that may be administered with the Therapeutics of the invention include, but are not limited to,

5 Monophosphoryl lipid immunomodulator, AdjuVax 100a, QS-21, QS-18, CRL1005, Aluminum salts, MF-59, and Virosomal adjuvant technology. Vaccines that may be administered with the Therapeutics of the invention include, but are not limited to, vaccines directed toward protection against MMR (measles, mumps, rubella), polio, varicella, tetanus/diphtheria, hepatitis A, hepatitis B, haemophilus influenzae B,

10 whooping cough, pneumonia, influenza, Lyme's Disease, rotavirus, cholera, yellow fever, Japanese encephalitis, poliomyelitis, rabies, typhoid fever, and pertussis. Combinations may be administered either concomitantly, e.g., as an admixture, separately but simultaneously or concurrently; or sequentially. This includes presentations in which the combined agents are administered together as a therapeutic

15 mixture, and also procedures in which the combined agents are administered separately but simultaneously, e.g., as through separate intravenous lines into the same individual. Administration "in combination" further includes the separate administration of one of the compounds or agents given first, followed by the second.

The Therapeutics of the invention may be administered alone or in

20 combination with other therapeutic agents. Therapeutic agents that may be administered in combination with the Therapeutics of the invention, include but not limited to, other members of the TNF family, chemotherapeutic agents, antibiotics, steroidal and non-steroidal anti-inflammatories, conventional immunotherapeutic agents, cytokines and/or growth factors. Combinations may be administered either

25 concomitantly, e.g., as an admixture, separately but simultaneously or concurrently; or sequentially. This includes presentations in which the combined agents are administered together as a therapeutic mixture, and also procedures in which the combined agents are administered separately but simultaneously, e.g., as through separate intravenous lines into the same individual. Administration "in combination"

30 further includes the separate administration of one of the compounds or agents given first, followed by the second.

In one embodiment, the Therapeutics of the invention are administered in combination with members of the TNF family. TNF, TNF-related or TNF-like molecules that may be administered with the Therapeutics of the invention include, but are not limited to, soluble forms of TNF-alpha, lymphotoxin-alpha (LT-alpha, also known as TNF-beta), LT-beta (found in complex heterotrimer LT-alpha2-beta), OPGL, FasL, CD27L, CD30L, CD40L, 4-1BBL, DcR3, OX40L, TNF-gamma (International Publication No. WO 96/14328), AIM-I (International Publication No. WO 97/33899), endokine-alpha (International Publication No. WO 98/07880), TR6 (International Publication No. WO 98/30694), OPG, and neutrokin-alpha (International Publication No. WO 98/18921, OX40, and nerve growth factor (NGF), and soluble forms of Fas, CD30, CD27, CD40 and 4-1BB, TR2 (International Publication No. WO 96/34095), DR3 (International Publication No. WO 97/33904), DR4 (International Publication No. WO 98/32856), TR5 (International Publication No. WO 98/30693), TR6 (International Publication No. WO 98/30694), TR7 (International Publication No. WO 98/41629), TRANK, TR9 (International Publication No. WO 98/56892), TR10 (International Publication No. WO 98/54202), 312C2 (International Publication No. WO 98/06842), and TR12, and soluble forms CD154, CD70, and CD153.

In certain embodiments, Therapeutics of the invention are administered in combination with antiretroviral agents, nucleoside reverse transcriptase inhibitors, non-nucleoside reverse transcriptase inhibitors, and/or protease inhibitors. Nucleoside reverse transcriptase inhibitors that may be administered in combination with the Therapeutics of the invention, include, but are not limited to, RETROVIR™ (zidovudine/AZT), VIDEX™ (didanosine/ddI), HIVID™ (zalcitabine/ddC), ZERIT™ (stavudine/d4T), EPIVIR™ (lamivudine/3TC), and COMBIVIR™ (zidovudine/lamivudine). Non-nucleoside reverse transcriptase inhibitors that may be administered in combination with the Therapeutics of the invention, include, but are not limited to, VIRAMUNE™ (nevirapine), RESCRIPTOR™ (delavirdine), and SUSTIVA™ (efavirenz). Protease inhibitors that may be administered in combination with the Therapeutics of the invention, include, but are not limited to, CRIVAN™ (indinavir), NORVIR™ (ritonavir), INVIRASE™ (saquinavir), and

VIRACEPT™ (nelfinavir). In a specific embodiment, antiretroviral agents, nucleoside reverse transcriptase inhibitors, non-nucleoside reverse transcriptase inhibitors, and/or protease inhibitors may be used in any combination with Therapeutics of the invention to treat AIDS and/or to prevent or treat HIV infection.

- 5 In other embodiments. Therapeutics of the invention may be administered in combination with anti-opportunistic infection agents. Anti-opportunistic agents that may be administered in combination with the Therapeutics of the invention, include, but are not limited to, TRIMETHOPRIM-SULFAMETHOXAZOLE™, DAPSONE™, PENTAMIDINE™, ATOVAQUONE™, ISONIAZID™, 10 RIFAMPIN™, PYRAZINAMIDE™, ETHAMBUTOL™, RIFABUTIN™, CLARITHROMYCIN™, AZITHROMYCIN™, GANCICLOVIR™, FOSCARNET™, CIDOFOVIR™, FLUCONAZOLE™, ITRACONAZOLE™, KETOCONAZOLE™, ACYCLOVIR™, FAMCICOLVIR™, PYRIMETHAMINE™, LEUCOVORIN™. NEUPOGEN™ (filgrastim/G-CSF), and LEUKINE™ 15 (sargramostim/GM-CSF). In a specific embodiment, Therapeutics of the invention are used in any combination with TRIMETHOPRIM-SULFAMETHOXAZOLE™, DAPSONE™, PENTAMIDINE™, and/or ATOVAQUONE™ to prophylactically treat or prevent an opportunistic *Pneumocystis carinii* pneumonia infection. In another specific embodiment. Therapeutics of the invention are used in any 20 combination with ISONIAZID™, RIFAMPIN™, PYRAZINAMIDE™, and/or ETHAMBUTOL™ to prophylactically treat or prevent an opportunistic *Mycobacterium avium* complex infection. In another specific embodiment, Therapeutics of the invention are used in any combination with RIFABUTIN™, CLARITHROMYCIN™, and/or AZITHROMYCIN™ to prophylactically treat or 25 prevent an opportunistic *Mycobacterium tuberculosis* infection. In another specific embodiment, Therapeutics of the invention are used in any combination with GANCICLOVIR™, FOSCARNET™, and/or CIDOFOVIR™ to prophylactically treat or prevent an opportunistic cytomegalovirus infection. In another specific embodiment, Therapeutics of the invention are used in any combination with 30 FLUCONAZOLE™, ITRACONAZOLE™, and/or KETOCONAZOLE™ to

prophylactically treat or prevent an opportunistic fungal infection. In another specific embodiment, Therapeutics of the invention are used in any combination with ACYCLOVIR™ and/or FAMCICOLVIR™ to prophylactically treat or prevent an opportunistic herpes simplex virus type I and/or type II infection. In another specific
5 embodiment, Therapeutics of the invention are used in any combination with PYRIMETHAMINE™ and/or LEUCOVORIN™ to prophylactically treat or prevent an opportunistic *Toxoplasma gondii* infection. In another specific embodiment, Therapeutics of the invention are used in any combination with LEUCOVORIN™ and/or NEUPOGEN™ to prophylactically treat or prevent an opportunistic bacterial
10 infection.

In a further embodiment, the Therapeutics of the invention are administered in combination with an antiviral agent. Antiviral agents that may be administered with the Therapeutics of the invention include, but are not limited to, acyclovir, ribavirin, amantadine, and remantidine.

15 In a further embodiment, the Therapeutics of the invention are administered in combination with an antibiotic agent. Antibiotic agents that may be administered with the Therapeutics of the invention include, but are not limited to, amoxicillin, beta-lactamases, aminoglycosides, beta-lactam (glycopeptide), beta-lactamases, Clindamycin, chloramphenicol, cephalosporins, ciprofloxacin, ciprofloxacin,
20 erythromycin, fluoroquinolones, macrolides, metronidazole, penicillins, quinolones, rifampin, streptomycin, sulfonamide, tetracyclines, trimethoprim, trimethoprim-sulfamthoxazole, and vancomycin.

Conventional nonspecific immunosuppressive agents, that may be administered in combination with the Therapeutics of the invention include, but are
25 not limited to, steroids, cyclosporine, cyclosporine analogs, cyclophosphamide methylprednisone, prednisone, azathioprine, FK-506, 15-deoxyspergualin, and other immunosuppressive agents that act by suppressing the function of responding T cells.

In specific embodiments, Therapeutics of the invention are administered in combination with immunosuppressants. Immunosuppressants preparations that may
30 be administered with the Therapeutics of the invention include, but are not limited to, ORTHOCLONE™ (OKT3), SANDIMMUNE™/NEORAL™/SANGDYA™

(cyclosporin), PROGRAF™ (tacrolimus), CELLCEPT™ (mycophenolate), Azathioprine, glucocorticosteroids, and RAPAMUNE™ (sirolimus). In a specific embodiment, immunosuppressants may be used to prevent rejection of organ or bone marrow transplantation.

5 In an additional embodiment, Therapeutics of the invention are administered alone or in combination with one or more intravenous immune globulin preparations. Intravenous immune globulin preparations that may be administered with the Therapeutics of the invention include, but not limited to, GAMMAR™, IVEEGAM™, SANDOGLOBULIN™, GAMMAGARD S/D™, and GAMIMUNE™.

10 In a specific embodiment, Therapeutics of the invention are administered in combination with intravenous immune globulin preparations in transplantation therapy (e.g., bone marrow transplant).

 In an additional embodiment, the Therapeutics of the invention are administered alone or in combination with an anti-inflammatory agent. Anti-
15 inflammatory agents that may be administered with the Therapeutics of the invention include, but are not limited to, glucocorticoids and the nonsteroidal anti-inflammatories, aminoarylcarboxylic acid derivatives, arylacetic acid derivatives, arylbutyric acid derivatives, arylcarboxylic acids, arylpropionic acid derivatives, pyrazoles, pyrazolones, salicylic acid derivatives, thiazinecarboxamides, e-
20 acetamidocaproic acid, S-adenosylmethionine, 3-amino-4-hydroxybutyric acid, amixetrine, bendazac, benzydamine, bucolome, difenpiramide, ditazol, emorfazone, guaiazulene, nabumetone, nimesulide, orgotein, oxaceprol, paranyline, perisoxal, pifoxime, proquazone, proxazole, and tenidap.

 In another embodiment, compositions of the invention are administered in
25 combination with a chemotherapeutic agent. Chemotherapeutic agents that may be administered with the Therapeutics of the invention include, but are not limited to, antibiotic derivatives (e.g., doxorubicin, bleomycin, daunorubicin, and dactinomycin); antiestrogens (e.g., tamoxifen); antimetabolites (e.g., fluorouracil, 5-FU, methotrexate, floxuridine, interferon alpha-2b, glutamic acid, plicamycin,
30 mercaptapurine, and 6-thioguanine); cytotoxic agents (e.g., carmustine, BCNU, lomustine, CCNU, cytosine arabinoside, cyclophosphamide, estramustine,

hydroxyurea, procarbazine, mitomycin, busulfan, cis-platin, and vincristine sulfate); hormones (e.g., medroxyprogesterone, estramustine phosphate sodium, ethinyl estradiol, estradiol, megestrol acetate, methyltestosterone, diethylstilbestrol diphosphate, chlorotrianisene, and testolactone); nitrogen mustard derivatives (e.g.,
5 mephalen, chorambucil, mechlorethamine (nitrogen mustard) and thiotepa); steroids and combinations (e.g., bethamethasone sodium phosphate); and others (e.g., dicarbazine, asparaginase, mitotane, vincristine sulfate, vinblastine sulfate, and etoposide).

In a specific embodiment, Therapeutics of the invention are administered in
10 combination with CHOP (cyclophosphamide, doxorubicin, vincristine, and prednisone) or any combination of the components of CHOP. In another embodiment, Therapeutics of the invention are administered in combination with Rituximab. In a further embodiment, Therapeutics of the invention are administered with Rituximab and CHOP, or Rituximab and any combination of the components of
15 CHOP.

In an additional embodiment, the Therapeutics of the invention are administered in combination with cytokines. Cytokines that may be administered with the Therapeutics of the invention include, but are not limited to, IL2, IL3, IL4, IL5, IL6, IL7, IL10, IL12, IL13, IL15, anti-CD40, CD40L, IFN-gamma and TNF-
20 alpha. In another embodiment, Therapeutics of the invention may be administered with any interleukin, including, but not limited to, IL-1alpha, IL-1beta, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IL-16, IL-17, IL-18, IL-19, IL-20, and IL-21.

In an additional embodiment, the Therapeutics of the invention are
25 administered in combination with angiogenic proteins. Angiogenic proteins that may be administered with the Therapeutics of the invention include, but are not limited to, Glioma Derived Growth Factor (GDGF), as disclosed in European Patent Number EP-399816; Platelet Derived Growth Factor-A (PDGF-A), as disclosed in European Patent Number EP-682110; Platelet Derived Growth Factor-B (PDGF-B), as
30 disclosed in European Patent Number EP-282317; Placental Growth Factor (PIGF), as disclosed in International Publication Number WO 92/06194; Placental Growth

Factor-2 (PlGF-2), as disclosed in Hauser et al., Growth Factors, 4:259-268 (1993); Vascular Endothelial Growth Factor (VEGF), as disclosed in International Publication Number WO 90/13649; Vascular Endothelial Growth Factor-A (VEGF-A), as disclosed in European Patent Number EP-506477; Vascular Endothelial Growth
5 Factor-2 (VEGF-2), as disclosed in International Publication Number WO 96/39515; Vascular Endothelial Growth Factor B (VEGF-3); Vascular Endothelial Growth Factor B-186 (VEGF-B186), as disclosed in International Publication Number WO 96/26736; Vascular Endothelial Growth Factor-D (VEGF-D), as disclosed in International Publication Number WO 98/02543; Vascular Endothelial Growth
10 Factor-D (VEGF-D), as disclosed in International Publication Number WO 98/07832; and Vascular Endothelial Growth Factor-E (VEGF-E), as disclosed in German Patent Number DE19639601. The above mentioned references are incorporated herein by reference herein.

In an additional embodiment, the Therapeutics of the invention are
15 administered in combination with hematopoietic growth factors. Hematopoietic growth factors that may be administered with the Therapeutics of the invention include, but are not limited to, LEUKINE™ (SARGRAMOSTIM™) and NEUPOGEN™ (FILGRASTIM™).

In an additional embodiment, the Therapeutics of the invention are
20 administered in combination with Fibroblast Growth Factors. Fibroblast Growth Factors that may be administered with the Therapeutics of the invention include, but are not limited to, FGF-1, FGF-2, FGF-3, FGF-4, FGF-5, FGF-6, FGF-7, FGF-8, FGF-9, FGF-10, FGF-11, FGF-12, FGF-13, FGF-14, and FGF-15.

In additional embodiments, the Therapeutics of the invention are administered in
25 combination with other therapeutic or prophylactic regimens, such as, for example, radiation therapy.

Example 24: Method of Treating Decreased Levels of the Polypeptide

The present invention relates to a method for treating an individual in need of
30 an increased level of a polypeptide of the invention in the body comprising administering to such an individual a composition comprising a therapeutically

effective amount of an agonist of the invention (including polypeptides of the invention). Moreover, it will be appreciated that conditions caused by a decrease in the standard or normal expression level of a secreted protein in an individual can be treated by administering the polypeptide of the present invention, preferably in the
5 secreted form. Thus, the invention also provides a method of treatment of an individual in need of an increased level of the polypeptide comprising administering to such an individual a Therapeutic comprising an amount of the polypeptide to increase the activity level of the polypeptide in such an individual.

For example, a patient with decreased levels of a polypeptide receives a daily
10 dose 0.1-100 ug/kg of the polypeptide for six consecutive days. Preferably, the polypeptide is in the secreted form. The exact details of the dosing scheme, based on administration and formulation, are provided in Example 23.

Example 25: Method of Treating Increased Levels of the Polypeptide

15 The present invention also relates to a method of treating an individual in need of a decreased level of a polypeptide of the invention in the body comprising administering to such an individual a composition comprising a therapeutically effective amount of an antagonist of the invention (including polypeptides and antibodies of the invention).

20 In one example, antisense technology is used to inhibit production of a polypeptide of the present invention. This technology is one example of a method of decreasing levels of a polypeptide, preferably a secreted form, due to a variety of etiologies, such as cancer. For example, a patient diagnosed with abnormally increased levels of a polypeptide is administered intravenously antisense
25 polynucleotides at 0.5, 1.0, 1.5, 2.0 and 3.0 mg/kg day for 21 days. This treatment is repeated after a 7-day rest period if the treatment was well tolerated. The formulation of the antisense polynucleotide is provided in Example 23.

Example 26: Method of Treatment Using Gene Therapy-Ex Vivo

30 One method of gene therapy transplants fibroblasts, which are capable of expressing a polypeptide, onto a patient. Generally, fibroblasts are obtained from a

subject by skin biopsy. The resulting tissue is placed in tissue-culture medium and separated into small pieces. Small chunks of the tissue are placed on a wet surface of a tissue culture flask, approximately ten pieces are placed in each flask. The flask is turned upside down, closed tight and left at room temperature over night. After 24
5 hours at room temperature, the flask is inverted and the chunks of tissue remain fixed to the bottom of the flask and fresh media (e.g., Ham's F12 media, with 10% FBS, penicillin and streptomycin) is added. The flasks are then incubated at 37 degree C for approximately one week.

At this time, fresh media is added and subsequently changed every several
10 days. After an additional two weeks in culture, a monolayer of fibroblasts emerge. The monolayer is trypsinized and scaled into larger flasks.

pMV-7 (Kirschmeier, P.T. et al., DNA, 7:219-25 (1988)), flanked by the long terminal repeats of the Moloney murine sarcoma virus, is digested with EcoRI and HindIII and subsequently treated with calf intestinal phosphatase. The linear vector is
15 fractionated on agarose gel and purified, using glass beads.

The cDNA encoding a polypeptide of the present invention can be amplified using PCR primers which correspond to the 5' and 3' end sequences respectively as set forth in Example 1 using primers and having appropriate restriction sites and initiation/stop codons, if necessary. Preferably, the 5' primer contains an EcoRI site
20 and the 3' primer includes a HindIII site. Equal quantities of the Moloney murine sarcoma virus linear backbone and the amplified EcoRI and HindIII fragment are added together, in the presence of T4 DNA ligase. The resulting mixture is maintained under conditions appropriate for ligation of the two fragments. The ligation mixture is then used to transform bacteria HB101, which are then plated onto
25 agar containing kanamycin for the purpose of confirming that the vector has the gene of interest properly inserted.

The amphotropic pA317 or GP+am12 packaging cells are grown in tissue culture to confluent density in Dulbecco's Modified Eagles Medium (DMEM) with 10% calf serum (CS), penicillin and streptomycin. The MSV vector containing the
30 gene is then added to the media and the packaging cells transduced with the vector.

The packaging cells now produce infectious viral particles containing the gene (the packaging cells are now referred to as producer cells).

Fresh media is added to the transduced producer cells, and subsequently, the media is harvested from a 10 cm plate of confluent producer cells. The spent media,
5 containing the infectious viral particles, is filtered through a millipore filter to remove detached producer cells and this media is then used to infect fibroblast cells. Media is removed from a sub-confluent plate of fibroblasts and quickly replaced with the media from the producer cells. This media is removed and replaced with fresh media. If the titer of virus is high, then virtually all fibroblasts will be infected and no
10 selection is required. If the titer is very low, then it is necessary to use a retroviral vector that has a selectable marker, such as neo or his. Once the fibroblasts have been efficiently infected, the fibroblasts are analyzed to determine whether protein is produced.

The engineered fibroblasts are then transplanted onto the host, either alone or
15 after having been grown to confluence on cytodex 3 microcarrier beads.

Example 27: Gene Therapy Using Endogenous Genes Corresponding To Polynucleotides of the Invention

Another method of gene therapy according to the present invention involves
20 operably associating the endogenous polynucleotide sequence of the invention with a promoter via homologous recombination as described, for example, in U.S. Patent NO: 5,641,670, issued June 24, 1997; International Publication NO: WO 96/29411, published September 26, 1996; International Publication NO: WO 94/12650, published August 4, 1994; Koller et al., *Proc. Natl. Acad. Sci. USA*, 86:8932-8935
25 (1989); and Zijlstra et al., *Nature*, 342:435-438 (1989). This method involves the activation of a gene which is present in the target cells, but which is not expressed in the cells, or is expressed at a lower level than desired.

Polynucleotide constructs are made which contain a promoter and targeting sequences, which are homologous to the 5' non-coding sequence of endogenous
30 polynucleotide sequence, flanking the promoter. The targeting sequence will be sufficiently near the 5' end of the polynucleotide sequence so the promoter will be

operably linked to the endogenous sequence upon homologous recombination. The promoter and the targeting sequences can be amplified using PCR. Preferably, the amplified promoter contains distinct restriction enzyme sites on the 5' and 3' ends. Preferably, the 3' end of the first targeting sequence contains the same restriction
5 enzyme site as the 5' end of the amplified promoter and the 5' end of the second targeting sequence contains the same restriction site as the 3' end of the amplified promoter.

The amplified promoter and the amplified targeting sequences are digested with the appropriate restriction enzymes and subsequently treated with calf intestinal
10 phosphatase. The digested promoter and digested targeting sequences are added together in the presence of T4 DNA ligase. The resulting mixture is maintained under conditions appropriate for ligation of the two fragments. The construct is size fractionated on an agarose gel then purified by phenol extraction and ethanol precipitation.

15 In this Example, the polynucleotide constructs are administered as naked polynucleotides via electroporation. However, the polynucleotide constructs may also be administered with transfection-facilitating agents, such as liposomes, viral sequences, viral particles, precipitating agents, etc. Such methods of delivery are known in the art.

20 Once the cells are transfected, homologous recombination will take place which results in the promoter being operably linked to the endogenous polynucleotide sequence. This results in the expression of polynucleotide corresponding to the polynucleotide in the cell. Expression may be detected by immunological staining, or any other method known in the art.

25 Fibroblasts are obtained from a subject by skin biopsy. The resulting tissue is placed in DMEM + 10% fetal calf serum. Exponentially growing or early stationary phase fibroblasts are trypsinized and rinsed from the plastic surface with nutrient medium. An aliquot of the cell suspension is removed for counting, and the remaining cells are subjected to centrifugation. The supernatant is aspirated and the pellet is
30 resuspended in 5 ml of electroporation buffer (20 mM HEPES pH 7.3, 137 mM NaCl, 5 mM KCl, 0.7 mM Na_2HPO_4 , 6 mM dextrose). The cells are recentrifuged, the

supernatant aspirated, and the cells resuspended in electroporation buffer containing 1 mg/ml acetylated bovine serum albumin. The final cell suspension contains approximately 3×10^6 cells/ml. Electroporation should be performed immediately following resuspension.

5 Plasmid DNA is prepared according to standard techniques. For example, to construct a plasmid for targeting to the locus corresponding to the polynucleotide of the invention, plasmid pUC18 (MBI Fermentas, Amherst, NY) is digested with HindIII. The CMV promoter is amplified by PCR with an XbaI site on the 5' end and a BamHI site on the 3' end. Two non-coding sequences are amplified via PCR: one
10 non-coding sequence (fragment 1) is amplified with a HindIII site at the 5' end and an Xba site at the 3' end; the other non-coding sequence (fragment 2) is amplified with a BamHI site at the 5' end and a HindIII site at the 3' end. The CMV promoter and the fragments (1 and 2) are digested with the appropriate enzymes (CMV promoter - XbaI and BamHI; fragment 1 - XbaI; fragment 2 - BamHI) and ligated together. The
15 resulting ligation product is digested with HindIII, and ligated with the HindIII-digested pUC18 plasmid.

Plasmid DNA is added to a sterile cuvette with a 0.4 cm electrode gap (Bio-Rad). The final DNA concentration is generally at least 120 μ g/ml. 0.5 ml of the cell suspension (containing approximately 1.5×10^6 cells) is then added to the cuvette,
20 and the cell suspension and DNA solutions are gently mixed. Electroporation is performed with a Gene-Pulser apparatus (Bio-Rad). Capacitance and voltage are set at 960 μ F and 250-300 V, respectively. As voltage increases, cell survival decreases, but the percentage of surviving cells that stably incorporate the introduced DNA into their genome increases dramatically. Given these parameters, a pulse time of
25 approximately 14-20 mSec should be observed.

Electroporated cells are maintained at room temperature for approximately 5 min, and the contents of the cuvette are then gently removed with a sterile transfer pipette. The cells are added directly to 10 ml of prewarmed nutrient media (DMEM with 15% calf serum) in a 10 cm dish and incubated at 37 degree C. The following
30 day, the media is aspirated and replaced with 10 ml of fresh media and incubated for a further 16-24 hours.

The engineered fibroblasts are then injected into the host, either alone or after having been grown to confluence on cytodex 3 microcarrier beads. The fibroblasts now produce the protein product. The fibroblasts can then be introduced into a patient as described above.

5

Example 28: Method of Treatment Using Gene Therapy - In Vivo

Another aspect of the present invention is using *in vivo* gene therapy methods to treat disorders, diseases and conditions. The gene therapy method relates to the introduction of naked nucleic acid (DNA, RNA, and antisense DNA or RNA) sequences into an animal to increase or decrease the expression of the polypeptide. The polynucleotide of the present invention may be operatively linked to a promoter or any other genetic elements necessary for the expression of the polypeptide by the target tissue. Such gene therapy and delivery techniques and methods are known in the art, see, for example, WO90/11092, WO98/11779; U.S. Patent NO. 5693622, 10 5705151, 5580859; Tabata et al., Cardiovasc. Res. 35(3):470-479 (1997); Chao et al., Pharmacol. Res. 35(6):517-522 (1997); Wolff, Neuromuscul. Disord. 7(5):314-318 (1997); Schwartz et al., Gene Ther. 3(5):405-411 (1996); Tsurumi et al., Circulation 15 94(12):3281-3290 (1996) (incorporated herein by reference).

The polynucleotide constructs may be delivered by any method that delivers injectable materials to the cells of an animal, such as, injection into the interstitial space of tissues (heart, muscle, skin, lung, liver, intestine and the like). The polynucleotide constructs can be delivered in a pharmaceutically acceptable liquid or aqueous carrier. 20

The term "naked" polynucleotide, DNA or RNA, refers to sequences that are free from any delivery vehicle that acts to assist, promote, or facilitate entry into the cell, including viral sequences, viral particles, liposome formulations, lipofectin or precipitating agents and the like. However, the polynucleotides of the present invention may also be delivered in liposome formulations (such as those taught in Felgner P.L. et al. (1995) Ann. NY Acad. Sci. 772:126-139 and Abdallah B. et al. 25 (1995) Biol. Cell 85(1):1-7) which can be prepared by methods well known to those skilled in the art. 30

The polynucleotide vector constructs used in the gene therapy method are preferably constructs that will not integrate into the host genome nor will they contain sequences that allow for replication. Any strong promoter known to those skilled in the art can be used for driving the expression of DNA. Unlike other gene therapies
5 techniques, one major advantage of introducing naked nucleic acid sequences into target cells is the transitory nature of the polynucleotide synthesis in the cells. Studies have shown that non-replicating DNA sequences can be introduced into cells to provide production of the desired polypeptide for periods of up to six months.

The polynucleotide construct can be delivered to the interstitial space of
10 tissues within the an animal, including of muscle, skin, brain, lung, liver, spleen, bone marrow, thymus, heart, lymph, blood, bone, cartilage, pancreas, kidney, gall bladder, stomach, intestine, testis, ovary, uterus, rectum, nervous system, eye, gland, and connective tissue. Interstitial space of the tissues comprises the intercellular fluid, mucopolysaccharide matrix among the reticular fibers of organ tissues, elastic fibers
15 in the walls of vessels or chambers, collagen fibers of fibrous tissues, or that same matrix within connective tissue ensheathing muscle cells or in the lacunae of bone. It is similarly the space occupied by the plasma of the circulation and the lymph fluid of the lymphatic channels. Delivery to the interstitial space of muscle tissue is preferred for the reasons discussed below. They may be conveniently delivered by injection
20 into the tissues comprising these cells. They are preferably delivered to and expressed in persistent, non-dividing cells which are differentiated, although delivery and expression may be achieved in non-differentiated or less completely differentiated cells, such as, for example, stem cells of blood or skin fibroblasts. *In vivo* muscle cells are particularly competent in their ability to take up and express
25 polynucleotides.

For the naked polynucleotide injection, an effective dosage amount of DNA or RNA will be in the range of from about 0.05 g/kg body weight to about 50 mg/kg body weight. Preferably the dosage will be from about 0.005 mg/kg to about 20 mg/kg and more preferably from about 0.05 mg/kg to about 5 mg/kg. Of course, as
30 the artisan of ordinary skill will appreciate, this dosage will vary according to the tissue site of injection. The appropriate and effective dosage of nucleic acid sequence

can readily be determined by those of ordinary skill in the art and may depend on the condition being treated and the route of administration. The preferred route of administration is by the parenteral route of injection into the interstitial space of tissues. However, other parenteral routes may also be used, such as, inhalation of an aerosol formulation particularly for delivery to lungs or bronchial tissues, throat or mucous membranes of the nose. In addition, naked polynucleotide constructs can be delivered to arteries during angioplasty by the catheter used in the procedure.

The dose response effects of injected polynucleotide in muscle *in vivo* is determined as follows. Suitable template DNA for production of mRNA coding for polypeptide of the present invention is prepared in accordance with a standard recombinant DNA methodology. The template DNA, which may be either circular or linear, is either used as naked DNA or complexed with liposomes. The quadriceps muscles of mice are then injected with various amounts of the template DNA.

Five to six week old female and male Balb/C mice are anesthetized by intraperitoneal injection with 0.3 ml of 2.5% Avertin. A 1.5 cm incision is made on the anterior thigh, and the quadriceps muscle is directly visualized. The template DNA is injected in 0.1 ml of carrier in a 1 cc syringe through a 27 gauge needle over one minute, approximately 0.5 cm from the distal insertion site of the muscle into the knee and about 0.2 cm deep. A suture is placed over the injection site for future localization, and the skin is closed with stainless steel clips.

After an appropriate incubation time (e.g., 7 days) muscle extracts are prepared by excising the entire quadriceps. Every fifth 15 um cross-section of the individual quadriceps muscles is histochemically stained for protein expression. A time course for protein expression may be done in a similar fashion except that quadriceps from different mice are harvested at different times. Persistence of DNA in muscle following injection may be determined by Southern blot analysis after preparing total cellular DNA and HIRT supernatants from injected and control mice. The results of the above experimentation in mice can be use to extrapolate proper dosages and other treatment parameters in humans and other animals using naked DNA.

Example 29: Transgenic Animals.

The polypeptides of the invention can also be expressed in transgenic animals. Animals of any species, including, but not limited to, mice, rats, rabbits, hamsters, guinea pigs, pigs, micro-pigs, goats, sheep, cows and non-human primates, *e.g.*,
5 baboons, monkeys, and chimpanzees may be used to generate transgenic animals. In a specific embodiment, techniques described herein or otherwise known in the art, are used to express polypeptides of the invention in humans, as part of a gene therapy protocol.

Any technique known in the art may be used to introduce the transgene (i.e.,
10 polynucleotides of the invention) into animals to produce the founder lines of transgenic animals. Such techniques include, but are not limited to, pronuclear microinjection (Paterson et al., Appl. Microbiol. Biotechnol. 40:691-698 (1994); Carver et al., Biotechnology (NY) 11:1263-1270 (1993); Wright et al., Biotechnology (NY) 9:830-834 (1991); and Hoppe et al., U.S. Pat. No. 4,873,191 (1989)); retrovirus
15 mediated gene transfer into germ lines (Van der Putten et al., Proc. Natl. Acad. Sci., USA 82:6148-6152 (1985)), blastocysts or embryos; gene targeting in embryonic stem cells (Thompson et al., Cell 56:313-321 (1989)); electroporation of cells or embryos (Lo, 1983, Mol Cell. Biol. 3:1803-1814 (1983)); introduction of the polynucleotides of the invention using a gene gun (see, e.g., Ulmer et al., Science
20 259:1745 (1993); introducing nucleic acid constructs into embryonic pluripotent stem cells and transferring the stem cells back into the blastocyst; and sperm-mediated gene transfer (Lavitrano et al., Cell 57:717-723 (1989); etc. For a review of such techniques, see Gordon, "Transgenic Animals," Intl. Rev. Cytol. 115:171-229 (1989), which is incorporated by reference herein in its entirety.

25 Any technique known in the art may be used to produce transgenic clones containing polynucleotides of the invention, for example, nuclear transfer into enucleated oocytes of nuclei from cultured embryonic, fetal, or adult cells induced to quiescence (Campell et al., Nature 380:64-66 (1996); Wilmut et al., Nature 385:810-813 (1997)).

30 The present invention provides for transgenic animals that carry the transgene in all their cells, as well as animals which carry the transgene in some, but not all their

cells, *i.e.*, mosaic animals or chimeric. The transgene may be integrated as a single transgene or as multiple copies such as in concatamers, *e.g.*, head-to-head tandems or head-to-tail tandems. The transgene may also be selectively introduced into and activated in a particular cell type by following, for example, the teaching of Lasko et al. (Lasko et al., Proc. Natl. Acad. Sci. USA 89:6232-6236 (1992)). The regulatory sequences required for such a cell-type specific activation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art. When it is desired that the polynucleotide transgene be integrated into the chromosomal site of the endogenous gene, gene targeting is preferred. Briefly, when such a technique is to be utilized, vectors containing some nucleotide sequences homologous to the endogenous gene are designed for the purpose of integrating, via homologous recombination with chromosomal sequences, into and disrupting the function of the nucleotide sequence of the endogenous gene. The transgene may also be selectively introduced into a particular cell type, thus inactivating the endogenous gene in only that cell type, by following, for example, the teaching of Gu et al. (Gu et al., Science 265:103-106 (1994)). The regulatory sequences required for such a cell-type specific inactivation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art.

Once transgenic animals have been generated, the expression of the recombinant gene may be assayed utilizing standard techniques. Initial screening may be accomplished by Southern blot analysis or PCR techniques to analyze animal tissues to verify that integration of the transgene has taken place. The level of mRNA expression of the transgene in the tissues of the transgenic animals may also be assessed using techniques which include, but are not limited to, Northern blot analysis of tissue samples obtained from the animal, *in situ* hybridization analysis, and reverse transcriptase-PCR (rt-PCR). Samples of transgenic gene-expressing tissue may also be evaluated immunocytochemically or immunohistochemically using antibodies specific for the transgene product.

Once the founder animals are produced, they may be bred, inbred, outbred, or crossbred to produce colonies of the particular animal. Examples of such breeding strategies include, but are not limited to: outbreeding of founder animals with more

than one integration site in order to establish separate lines; inbreeding of separate lines in order to produce compound transgenics that express the transgene at higher levels because of the effects of additive expression of each transgene; crossing of heterozygous transgenic animals to produce animals homozygous for a given
5 integration site in order to both augment expression and eliminate the need for screening of animals by DNA analysis; crossing of separate homozygous lines to produce compound heterozygous or homozygous lines; and breeding to place the transgene on a distinct background that is appropriate for an experimental model of interest.

10 Transgenic animals of the invention have uses which include, but are not limited to, animal model systems useful in elaborating the biological function of polypeptides of the present invention, studying diseases, disorders, and/or conditions associated with aberrant expression, and in screening for compounds effective in ameliorating such diseases, disorders, and/or conditions.

15

Example 30: Knock-Out Animals.

Endogenous gene expression can also be reduced by inactivating or "knocking out" the gene and/or its promoter using targeted homologous recombination. (*E.g.*, see Smithies et al., Nature 317:230-234 (1985); Thomas & Capecchi, Cell 51:503-
20 512 (1987); Thompson et al., Cell 5:313-321 (1989); each of which is incorporated by reference herein in its entirety). For example, a mutant, non-functional polynucleotide of the invention (or a completely unrelated DNA sequence) flanked by DNA homologous to the endogenous polynucleotide sequence (either the coding regions or regulatory regions of the gene) can be used, with or without a selectable
25 marker and/or a negative selectable marker, to transfect cells that express polypeptides of the invention *in vivo*. In another embodiment, techniques known in the art are used to generate knockouts in cells that contain, but do not express the gene of interest. Insertion of the DNA construct, via targeted homologous recombination, results in inactivation of the targeted gene. Such approaches are particularly suited in
30 research and agricultural fields where modifications to embryonic stem cells can be used to generate animal offspring with an inactive targeted gene (*e.g.*, see Thomas &

Capecchi 1987 and Thompson 1989, *supra*). However this approach can be routinely adapted for use in humans provided the recombinant DNA constructs are directly administered or targeted to the required site *in vivo* using appropriate viral vectors that will be apparent to those of skill in the art.

5 In further embodiments of the invention, cells that are genetically engineered to express the polypeptides of the invention, or alternatively, that are genetically engineered not to express the polypeptides of the invention (e.g., knockouts) are administered to a patient *in vivo*. Such cells may be obtained from the patient (i.e., animal, including human) or an MHC compatible donor and can include, but are not
10 limited to fibroblasts, bone marrow cells, blood cells (e.g., lymphocytes), adipocytes, muscle cells, endothelial cells etc. The cells are genetically engineered *in vitro* using recombinant DNA techniques to introduce the coding sequence of polypeptides of the invention into the cells, or alternatively, to disrupt the coding sequence and/or endogenous regulatory sequence associated with the polypeptides of the invention,
15 e.g., by transduction (using viral vectors, and preferably vectors that integrate the transgene into the cell genome) or transfection procedures, including, but not limited to, the use of plasmids, cosmids, YACs, naked DNA, electroporation, liposomes, etc. The coding sequence of the polypeptides of the invention can be placed under the control of a strong constitutive or inducible promoter or promoter/enhancer to achieve
20 expression, and preferably secretion, of the polypeptides of the invention. The engineered cells which express and preferably secrete the polypeptides of the invention can be introduced into the patient systemically, e.g., in the circulation, or intraperitoneally.

 Alternatively, the cells can be incorporated into a matrix and implanted in the
25 body, e.g., genetically engineered fibroblasts can be implanted as part of a skin graft; genetically engineered endothelial cells can be implanted as part of a lymphatic or vascular graft. (See, for example, Anderson et al. U.S. Patent No. 5,399,349; and Mulligan & Wilson, U.S. Patent No. 5,460,959 each of which is incorporated by reference herein in its entirety).

30 When the cells to be administered are non-autologous or non-MHC compatible cells, they can be administered using well known techniques which

prevent the development of a host immune response against the introduced cells. For example, the cells may be introduced in an encapsulated form which, while allowing for an exchange of components with the immediate extracellular environment, does not allow the introduced cells to be recognized by the host immune system.

- 5 Transgenic and "knock-out" animals of the invention have uses which include, but are not limited to, animal model systems useful in elaborating the biological function of polypeptides of the present invention, studying diseases, disorders, and/or conditions associated with aberrant expression, and in screening for compounds effective in ameliorating such diseases, disorders, and/or conditions.

10

Example 31: Production of an Antibody

a) Hybridoma Technology

The antibodies of the present invention can be prepared by a variety of methods. (See, Current Protocols, Chapter 2.) As one example of such methods, cells
15 expressing polypeptide(s) of the invention are administered to an animal to induce the production of sera containing polyclonal antibodies. In a preferred method, a preparation of polypeptide(s) of the invention is prepared and purified to render it substantially free of natural contaminants. Such a preparation is then introduced into an animal in order to produce polyclonal antisera of greater specific activity.

- 20 Monoclonal antibodies specific for polypeptide(s) of the invention are prepared using hybridoma technology. (Kohler et al., Nature 256:495 (1975); Kohler et al., Eur. J. Immunol. 6:511 (1976); Kohler et al., Eur. J. Immunol. 6:292 (1976); Hammerling et al., in: Monoclonal Antibodies and T-Cell Hybridomas. Elsevier, N.Y., pp. 563-681 (1981)). In general, an animal (preferably a mouse) is immunized
25 with polypeptide(s) of the invention or, more preferably, with a secreted polypeptide-expressing cell. Such polypeptide-expressing cells are cultured in any suitable tissue culture medium, preferably in Earle's modified Eagle's medium supplemented with 10% fetal bovine serum (inactivated at about 56°C), and supplemented with about 10 g/l of nonessential amino acids, about 1,000 U/ml of penicillin, and about 100 µg/ml
30 of streptomycin.

The splenocytes of such mice are extracted and fused with a suitable myeloma cell line. Any suitable myeloma cell line may be employed in accordance with the present invention; however, it is preferable to employ the parent myeloma cell line (SP2O), available from the ATCC. After fusion, the resulting hybridoma cells are
5 selectively maintained in HAT medium, and then cloned by limiting dilution as described by Wands et al. (Gastroenterology 80:225-232 (1981)). The hybridoma cells obtained through such a selection are then assayed to identify clones which secrete antibodies capable of binding the polypeptide(s) of the invention.

Alternatively, additional antibodies capable of binding to polypeptide(s) of the
10 invention can be produced in a two-step procedure using anti-idiotypic antibodies. Such a method makes use of the fact that antibodies are themselves antigens, and therefore, it is possible to obtain an antibody which binds to a second antibody. In accordance with this method, protein specific antibodies are used to immunize an animal, preferably a mouse. The splenocytes of such an animal are then used to
15 produce hybridoma cells, and the hybridoma cells are screened to identify clones which produce an antibody whose ability to bind to the protein-specific antibody can be blocked by polypeptide(s) of the invention. Such antibodies comprise anti-idiotypic antibodies to the protein-specific antibody and are used to immunize an animal to induce formation of further protein-specific antibodies.

20 For in vivo use of antibodies in humans, an antibody is "humanized". Such antibodies can be produced using genetic constructs derived from hybridoma cells producing the monoclonal antibodies described above. Methods for producing chimeric and humanized antibodies are known in the art and are discussed herein. (See, for review, Morrison, Science 229:1202 (1985); Oi et al., BioTechniques 4:214
25 (1986); Cabilly et al., U.S. Patent No. 4,816,567; Taniguchi et al., EP 171496; Morrison et al., EP 173494; Neuberger et al., WO 8601533; Robinson et al., WO 8702671; Boulianne et al., Nature 312:643 (1984); Neuberger et al., Nature 314:268 (1985).)

30 b) Isolation Of Antibody Fragments Directed Against
Polypeptide(s) From A Library Of scFvs

Naturally occurring V-genes isolated from human PBLs are constructed into a library of antibody fragments which contain reactivities against polypeptide(s) of the invention to which the donor may or may not have been exposed (see e.g., U.S. Patent 5,885,793 incorporated herein by reference in its entirety).

5 Rescue of the Library.

A library of scFvs is constructed from the RNA of human PBLs as described in PCT publication WO 92/01047. To rescue phage displaying antibody fragments, approximately 109 E. coli harboring the phagemid are used to inoculate 50 ml of 2xTY containing 1% glucose and 100 µg/ml of ampicillin (2xTY-AMP-GLU) and
10 grown to an O.D. of 0.8 with shaking. Five ml of this culture is used to inoculate 50 ml of 2xTY-AMP-GLU, 2 x 10⁸ TU of delta gene 3 helper (M13 delta gene III, see PCT publication WO 92/01047) are added and the culture incubated at 37°C for 45 minutes without shaking and then at 37°C for 45 minutes with shaking. The culture is centrifuged at 4000 r.p.m. for 10 min. and the pellet resuspended in 2 liters of 2xTY
15 containing 100 µg/ml ampicillin and 50 µg/ml kanamycin and grown overnight. Phage are prepared as described in PCT publication WO 92/01047.

M13 delta gene III is prepared as follows: M13 delta gene III helper phage does not encode gene III protein, hence the phage(mid) displaying antibody fragments have a greater avidity of binding to antigen. Infectious M13 delta gene III
20 particles are made by growing the helper phage in cells harboring a pUC19 derivative supplying the wild type gene III protein during phage morphogenesis. The culture is incubated for 1 hour at 37° C without shaking and then for a further hour at 37°C with shaking. Cells are spun down (IEC-Centra 8.400 r.p.m. for 10 min), resuspended in 300 ml 2xTY broth containing 100 µg ampicillin/ml and 25 µg kanamycin/ml (2xTY-
25 AMP-KAN) and grown overnight, shaking at 37°C. Phage particles are purified and concentrated from the culture medium by two PEG-precipitations (Sambrook et al., 1990), resuspended in 2 ml PBS and passed through a 0.45 µm filter (Minisart NML; Sartorius) to give a final concentration of approximately 10¹³ transducing units/ml (ampicillin-resistant clones).

30 Panning of the Library.

Immunotubes (Nunc) are coated overnight in PBS with 4 ml of either 100 $\mu\text{g/ml}$ or 10 $\mu\text{g/ml}$ of a polypeptide of the present invention. Tubes are blocked with 2% Marvel-PBS for 2 hours at 37°C and then washed 3 times in PBS. Approximately 1013 TU of phage is applied to the tube and incubated for 30 minutes at room temperature tumbling on an over and under turntable and then left to stand for another 1.5 hours. Tubes are washed 10 times with PBS 0.1% Tween-20 and 10 times with PBS. Phage are eluted by adding 1 ml of 100 mM triethylamine and rotating 15 minutes on an under and over turntable after which the solution is immediately neutralized with 0.5 ml of 1.0M Tris-HCl, pH 7.4. Phage are then used to infect 10 ml of mid-log E. coli TG1 by incubating eluted phage with bacteria for 30 minutes at 37°C. The E. coli are then plated on TYE plates containing 1% glucose and 100 $\mu\text{g/ml}$ ampicillin. The resulting bacterial library is then rescued with delta gene 3 helper phage as described above to prepare phage for a subsequent round of selection. This process is then repeated for a total of 4 rounds of affinity purification with tube-washing increased to 20 times with PBS, 0.1% Tween-20 and 20 times with PBS for rounds 3 and 4.

Characterization of Binders.

Eluted phage from the 3rd and 4th rounds of selection are used to infect E. coli HB 2151 and soluble scFv is produced (Marks, et al., 1991) from single colonies for assay. ELISAs are performed with microtitre plates coated with either 10 pg/ml of the polypeptide of the present invention in 50 mM bicarbonate pH 9.6. Clones positive in ELISA are further characterized by PCR fingerprinting (see, e.g., PCT publication WO 92/01047) and then by sequencing. These ELISA positive clones may also be further characterized by techniques known in the art, such as, for example, epitope mapping, binding affinity, receptor signal transduction, ability to block or competitively inhibit antibody/antigen binding, and competitive agonistic or antagonistic activity.

Example 32: Assays Detecting Stimulation or Inhibition of B cell Proliferation and Differentiation

Generation of functional humoral immune responses requires both soluble and cognate signaling between B-lineage cells and their microenvironment. Signals may
5 impart a positive stimulus that allows a B-lineage cell to continue its programmed development, or a negative stimulus that instructs the cell to arrest its current developmental pathway. To date, numerous stimulatory and inhibitory signals have been found to influence B cell responsiveness including IL-2, IL-4, IL-5, IL-6, IL-7, IL10, IL-13, IL-14 and IL-15. Interestingly, these signals are by themselves weak effectors but can,
10 in combination with various co-stimulatory proteins, induce activation, proliferation, differentiation, homing, tolerance and death among B cell populations.

One of the best studied classes of B-cell co-stimulatory proteins is the TNF-superfamily. Within this family CD40, CD27, and CD30 along with their respective ligands CD154, CD70, and CD153 have been found to regulate a variety of immune
15 responses. Assays which allow for the detection and/or observation of the proliferation and differentiation of these B-cell populations and their precursors are valuable tools in determining the effects various proteins may have on these B-cell populations in terms of proliferation and differentiation. Listed below are two assays designed to allow for the detection of the differentiation, proliferation, or inhibition of B-cell populations and their
20 precursors.

In Vitro Assay- Purified polypeptides of the invention, or truncated forms thereof, is assessed for its ability to induce activation, proliferation, differentiation or inhibition and/or death in B-cell populations and their precursors. The activity of the polypeptides of the invention on purified tonsillar B cells, measured qualitatively over the dose
25 range from 0.1 to 10,000 ng/mL, is assessed in a standard B-lymphocyte co-stimulation assay in which purified tonsillar B cells are cultured in the presence of either formalin-fixed Staphylococcus aureus Cowan I (SAC) or immobilized anti-human IgM antibody as the priming agent. Second signals such as IL-2 and IL-15 synergize with SAC and IgM crosslinking to elicit B cell proliferation as measured by tritiated-thymidine incorporation.
30 Novel synergizing agents can be readily identified using this assay. The assay involves isolating human tonsillar B cells by magnetic bead (MACS) depletion of CD3-positive

cells. The resulting cell population is greater than 95% B cells as assessed by expression of CD45R(B220).

Various dilutions of each sample are placed into individual wells of a 96-well plate to which are added 10^5 B-cells suspended in culture medium (RPMI 1640 containing 10%
5 FBS, 5×10^{-5} M 2ME, 100U/ml penicillin, 10ug/ml streptomycin, and 10^{-5} dilution of SAC) in a total volume of 150ul. Proliferation or inhibition is quantitated by a 20h pulse (1uCi/well) with 3H-thymidine (6.7 Ci/mM) beginning 72h post factor addition. The positive and negative controls are IL2 and medium respectively.

In Vivo Assay- BALB/c mice are injected (i.p.) twice per day with buffer only, or
10 2 mg/Kg of a polypeptide of the invention, or truncated forms thereof. Mice receive this treatment for 4 consecutive days, at which time they are sacrificed and various tissues and serum collected for analyses. Comparison of H&E sections from normal spleens and spleens treated with polypeptides of the invention identify the results of the activity of the polypeptides on spleen cells, such as the diffusion of peri-arterial lymphatic sheaths,
15 and/or significant increases in the nucleated cellularity of the red pulp regions, which may indicate the activation of the differentiation and proliferation of B-cell populations.

Immunohistochemical studies using a B cell marker, anti-CD45R(B220), are used to determine whether any physiological changes to splenic cells, such as splenic disorganization, are due to increased B-cell representation within loosely defined B-cell
20 zones that infiltrate established T-cell regions.

Flow cytometric analyses of the spleens from mice treated with polypeptide is used to indicate whether the polypeptide specifically increases the proportion of ThB+, CD45R(B220)dull B cells over that which is observed in control mice.

Likewise, a predicted consequence of increased mature B-cell representation in
25 vivo is a relative increase in serum Ig titers. Accordingly, serum IgM and IgA levels are compared between buffer and polypeptide-treated mice.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides of the invention (e.g., gene therapy), agonists, and/or
30 antagonists of polynucleotides or polypeptides of the invention.

Example 33: T Cell Proliferation Assay

A CD3-induced proliferation assay is performed on PBMCs and is measured by the uptake of ^3H -thymidine. The assay is performed as follows. Ninety-six well plates are coated with 100 μl /well of mAb to CD3 (HIT3a, Pharmingen) or isotype-matched control mAb (B33.1) overnight at 4 degrees C (1 $\mu\text{g}/\text{ml}$ in .05M bicarbonate buffer, pH 9.5), then washed three times with PBS. PBMC are isolated by F/H gradient centrifugation from human peripheral blood and added to quadruplicate wells (5×10^4 /well) of mAb coated plates in RPMI containing 10% FCS and P/S in the presence of varying concentrations of polypeptides of the invention (total volume 200 μl). Relevant protein buffer and medium alone are controls. After 48 hr. culture at 37 degrees C, plates are spun for 2 min. at 1000 rpm and 100 μl of supernatant is removed and stored -20 degrees C for measurement of IL-2 (or other cytokines) if effect on proliferation is observed. Wells are supplemented with 100 μl of medium containing 0.5 uCi of ^3H -thymidine and cultured at 37 degrees C for 18-24 hr. Wells are harvested and incorporation of ^3H -thymidine used as a measure of proliferation. Anti-CD3 alone is the positive control for proliferation. IL-2 (100 U/ml) is also used as a control which enhances proliferation. Control antibody which does not induce proliferation of T cells is used as the negative controls for the effects of polypeptides of the invention.

The studies described in this example tested activity of polypeptides of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides of the invention (e.g., gene therapy), agonists, and/or antagonists of polynucleotides or polypeptides of the invention.

Example 34: Effect of Polypeptides of the Invention on the Expression of MHC Class II, Costimulatory and Adhesion Molecules and Cell Differentiation of Monocytes and Monocyte-Derived Human Dendritic Cells

Dendritic cells are generated by the expansion of proliferating precursors found in the peripheral blood: adherent PBMC or elutriated monocytic fractions are cultured for 7-10 days with GM-CSF (50 ng/ml) and IL-4 (20 ng/ml). These dendritic cells have the characteristic phenotype of immature cells (expression of CD1, CD80, CD86, CD40 and

MHC class II antigens). Treatment with activating factors, such as TNF- α , causes a rapid change in surface phenotype (increased expression of MHC class I and II, costimulatory and adhesion molecules, downregulation of FC γ RII, upregulation of CD83). These changes correlate with increased antigen-presenting capacity and with functional
5 maturation of the dendritic cells.

FACS analysis of surface antigens is performed as follows. Cells are treated 1-3 days with increasing concentrations of polypeptides of the invention or LPS (positive control), washed with PBS containing 1% BSA and 0.02 mM sodium azide, and then incubated with 1:20 dilution of appropriate FITC- or PE-labeled monoclonal antibodies
10 for 30 minutes at 4 degrees C. After an additional wash, the labeled cells are analyzed by flow cytometry on a FACScan (Becton Dickinson).

Effect on the production of cytokines. Cytokines generated by dendritic cells, in particular IL-12, are important in the initiation of T-cell dependent immune responses. IL-
15 12 strongly influences the development of Th1 helper T-cell immune response, and induces cytotoxic T and NK cell function. An ELISA is used to measure the IL-12 release as follows. Dendritic cells (10^6 /ml) are treated with increasing concentrations of polypeptides of the invention for 24 hours. LPS (100 ng/ml) is added to the cell culture as positive control. Supernatants from the cell cultures are then collected and analyzed for
20 IL-12 content using commercial ELISA kit (e.g., R & D Systems (Minneapolis, MN)). The standard protocols provided with the kits are used.

Effect on the expression of MHC Class II, costimulatory and adhesion molecules. Three major families of cell surface antigens can be identified on monocytes: adhesion
25 molecules, molecules involved in antigen presentation, and Fc receptor. Modulation of the expression of MHC class II antigens and other costimulatory molecules, such as B7 and ICAM-1, may result in changes in the antigen presenting capacity of monocytes and ability to induce T cell activation. Increase expression of Fc receptors may correlate with improved monocyte cytotoxic activity, cytokine release and phagocytosis.

30 FACS analysis is used to examine the surface antigens as follows. Monocytes are treated 1-5 days with increasing concentrations of polypeptides of the invention or LPS

(positive control), washed with PBS containing 1% BSA and 0.02 mM sodium azide, and then incubated with 1:20 dilution of appropriate FITC- or PE-labeled monoclonal antibodies for 30 minutes at 4 degreesC. After an additional wash, the labeled cells are analyzed by flow cytometry on a FACScan (Becton Dickinson).

5

Monocyte activation and/or increased survival. Assays for molecules that activate (or alternatively, inactivate) monocytes and/or increase monocyte survival (or alternatively, decrease monocyte survival) are known in the art and may routinely be applied to determine whether a molecule of the invention functions as an inhibitor or
10 activator of monocytes. Polypeptides, agonists, or antagonists of the invention can be screened using the three assays described below. For each of these assays, Peripheral blood mononuclear cells (PBMC) are purified from single donor leukopacks (American Red Cross, Baltimore, MD) by centrifugation through a Histopaque gradient (Sigma). Monocytes are isolated from PBMC by counterflow centrifugal elutriation.

15

Monocyte Survival Assay. Human peripheral blood monocytes progressively lose viability when cultured in absence of serum or other stimuli. Their death results from internally regulated process (apoptosis). Addition to the culture of activating factors, such as TNF-alpha dramatically improves cell survival and prevents DNA fragmentation.
20 Propidium iodide (PI) staining is used to measure apoptosis as follows. Monocytes are cultured for 48 hours in polypropylene tubes in serum-free medium (positive control), in the presence of 100 ng/ml TNF-alpha (negative control), and in the presence of varying concentrations of the compound to be tested. Cells are suspended at a concentration of 2×10^6 /ml in PBS containing PI at a final concentration of 5 μ g/ml, and then incubated at room
25 temperature for 5 minutes before FACScan analysis. PI uptake has been demonstrated to correlate with DNA fragmentation in this experimental paradigm.

Effect on cytokine release. An important function of monocytes/macrophages is their regulatory activity on other cellular populations of the immune system through the
30 release of cytokines after stimulation. An ELISA to measure cytokine release is performed as follows. Human monocytes are incubated at a density of 5×10^5 cells/ml with

increasing concentrations of the a polypeptide of the invention and under the same conditions, but in the absence of the polypeptide. For IL-12 production, the cells are primed overnight with IFN (100 U/ml) in presence of a polypeptide of the invention. LPS (10 ng/ml) is then added. Conditioned media are collected after 24h and kept frozen until
5 use. Measurement of TNF-alpha, IL-10, MCP-1 and IL-8 is then performed using a commercially available ELISA kit (e.g., R & D Systems (Minneapolis, MN)) and applying the standard protocols provided with the kit.

Oxidative burst. Purified monocytes are plated in 96-w plate at 2×10^5 cell/well.
10 Increasing concentrations of polypeptides of the invention are added to the wells in a total volume of 0.2 ml culture medium (RPMI 1640 + 10% FCS, glutamine and antibiotics). After 3 days incubation, the plates are centrifuged and the medium is removed from the wells. To the macrophage monolayers, 0.2 ml per well of phenol red solution (140 mM NaCl, 10 mM potassium phosphate buffer pH 7.0, 5.5 mM dextrose, 0.56 mM phenol red
15 and 19 U/ml of HRPO) is added, together with the stimulant (200 nM PMA). The plates are incubated at 37°C for 2 hours and the reaction is stopped by adding 20 µl 1N NaOH per well. The absorbance is read at 610 nm. To calculate the amount of H_2O_2 produced by the macrophages, a standard curve of a H_2O_2 solution of known molarity is performed for each experiment.

20 The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polypeptides, polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

25 Example 35: Biological Effects of Polypeptides of the Invention

Astrocyte and Neuronal Assays

Recombinant polypeptides of the invention, expressed in *Escherichia coli* and purified as described above, can be tested for activity in promoting the survival, neurite outgrowth, or phenotypic differentiation of cortical neuronal cells and for inducing the
30 proliferation of glial fibrillary acidic protein immunopositive cells, astrocytes. The selection of cortical cells for the bioassay is based on the prevalent expression of FGF-1

and FGF-2 in cortical structures and on the previously reported enhancement of cortical neuronal survival resulting from FGF-2 treatment. A thymidine incorporation assay, for example, can be used to elucidate a polypeptide of the invention's activity on these cells.

Moreover, previous reports describing the biological effects of FGF-2 (basic FGF) on cortical or hippocampal neurons *in vitro* have demonstrated increases in both neuron survival and neurite outgrowth (Walicke et al., "Fibroblast growth factor promotes survival of dissociated hippocampal neurons and enhances neurite extension." *Proc. Natl. Acad. Sci. USA* 83:3012-3016. (1986). assay herein incorporated by reference in its entirety). However, reports from experiments done on PC-12 cells suggest that these two responses are not necessarily synonymous and may depend on not only which FGF is being tested but also on which receptor(s) are expressed on the target cells. Using the primary cortical neuronal culture paradigm, the ability of a polypeptide of the invention to induce neurite outgrowth can be compared to the response achieved with FGF-2 using, for example, a thymidine incorporation assay.

15

Fibroblast and endothelial cell assays

Human lung fibroblasts are obtained from Clonetics (San Diego, CA) and maintained in growth media from Clonetics. Dermal microvascular endothelial cells are obtained from Cell Applications (San Diego, CA). For proliferation assays, the human lung fibroblasts and dermal microvascular endothelial cells can be cultured at 5,000 cells/well in a 96-well plate for one day in growth medium. The cells are then incubated for one day in 0.1% BSA basal medium. After replacing the medium with fresh 0.1% BSA medium, the cells are incubated with the test proteins for 3 days. Alamar Blue (Alamar Biosciences, Sacramento, CA) is added to each well to a final concentration of 10%. The cells are incubated for 4 hr. Cell viability is measured by reading in a CytoFluor fluorescence reader. For the PGE₂ assays, the human lung fibroblasts are cultured at 5,000 cells/well in a 96-well plate for one day. After a medium change to 0.1% BSA basal medium, the cells are incubated with FGF-2 or polypeptides of the invention with or without IL-1 α for 24 hours. The supernatants are collected and assayed for PGE₂ by EIA

kit (Cayman, Ann Arbor, MI). For the IL-6 assays, the human lung fibroblasts are cultured at 5,000 cells/well in a 96-well plate for one day. After a medium change to 0.1% BSA basal medium, the cells are incubated with FGF-2 or with or without polypeptides of the invention IL-1 α for 24 hours. The supernatants are collected and
5 assayed for IL-6 by ELISA kit (Endogen, Cambridge, MA).

Human lung fibroblasts are cultured with FGF-2 or polypeptides of the invention for 3 days in basal medium before the addition of Alamar Blue to assess effects on growth of the fibroblasts. FGF-2 should show a stimulation at 10 - 2500 ng/ml which can be used to compare stimulation with polypeptides of the invention.

10

Parkinson Models.

The loss of motor function in Parkinson's disease is attributed to a deficiency of striatal dopamine resulting from the degeneration of the nigrostriatal dopaminergic projection neurons. An animal model for Parkinson's that has been extensively
15 characterized involves the systemic administration of 1-methyl-4 phenyl 1,2,3,6-tetrahydropyridine (MPTP). In the CNS, MPTP is taken-up by astrocytes and catabolized by monoamine oxidase B to 1-methyl-4-phenyl pyridine (MPP⁺) and released. Subsequently, MPP⁺ is actively accumulated in dopaminergic neurons by the high-affinity reuptake transporter for dopamine. MPP⁺ is then concentrated in mitochondria by the
20 electrochemical gradient and selectively inhibits nicotinamide adenine disphosphate: ubiquinone oxidoreductionase (complex I), thereby interfering with electron transport and eventually generating oxygen radicals.

It has been demonstrated in tissue culture paradigms that FGF-2 (basic FGF) has trophic activity towards nigral dopaminergic neurons (Ferrari et al., Dev. Biol. 1989).
25 Recently, Dr. Unsicker's group has demonstrated that administering FGF-2 in gel foam implants in the striatum results in the near complete protection of nigral dopaminergic neurons from the toxicity associated with MPTP exposure (Otto and Unsicker, J. Neuroscience, 1990).

Based on the data with FGF-2, polypeptides of the invention can be evaluated to
30 determine whether it has an action similar to that of FGF-2 in enhancing dopaminergic

neuronal survival *in vitro* and it can also be tested *in vivo* for protection of dopaminergic neurons in the striatum from the damage associated with MPTP treatment. The potential effect of a polypeptide of the invention is first examined *in vitro* in a dopaminergic neuronal cell culture paradigm. The cultures are prepared by dissecting the midbrain floor
5 plate from gestation day 14 Wistar rat embryos. The tissue is dissociated with trypsin and seeded at a density of 200,000 cells/cm² on polyorthinine-laminin coated glass coverslips. The cells are maintained in Dulbecco's Modified Eagle's medium and F12 medium containing hormonal supplements (N1). The cultures are fixed with paraformaldehyde after 8 days *in vitro* and are processed for tyrosine hydroxylase, a specific marker for
10 dopaminergic neurons, immunohistochemical staining. Dissociated cell cultures are prepared from embryonic rats. The culture medium is changed every third day and the factors are also added at that time.

Since the dopaminergic neurons are isolated from animals at gestation day 14, a developmental time which is past the stage when the dopaminergic precursor cells are
15 proliferating, an increase in the number of tyrosine hydroxylase immunopositive neurons would represent an increase in the number of dopaminergic neurons surviving *in vitro*. Therefore, if a polypeptide of the invention acts to prolong the survival of dopaminergic neurons, it would suggest that the polypeptide may be involved in Parkinson's Disease.

The studies described in this example tested activity of a polypeptide of the
20 invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 36: The Effect of Polypeptides of the Invention on the Growth of
25 **Vascular Endothelial Cells**

On day 1, human umbilical vein endothelial cells (HUVEC) are seeded at 2.5×10^4 cells/35 mm dish density in M199 medium containing 4% fetal bovine serum (FBS), 16 units/ml heparin, and 50 units/ml endothelial cell growth supplements (ECGS, Biotechnology, Inc.). On day 2, the medium is replaced with M199 containing 10% FBS, 8
30 units/ml heparin. A polypeptide having the amino acid sequence of SEQ ID NO:Y, and positive controls, such as VEGF and basic FGF (bFGF) are added, at varying

concentrations. On days 4 and 6, the medium is replaced. On day 8, cell number is determined with a Coulter Counter.

An increase in the number of HUVEC cells indicates that the polypeptide of the invention may proliferate vascular endothelial cells.

- 5 The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

10 **Example 37: Stimulatory Effect of Polypeptides of the Invention on the Proliferation of Vascular Endothelial Cells**

- For evaluation of mitogenic activity of growth factors, the colorimetric MTS (3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)2H-tetrazolium) assay with the electron coupling reagent PMS (phenazine methosulfate) was performed (CellTiter 96 AQ, Promega). Cells are seeded in a 96-well plate (5,000 cells/well) in 0.1 mL serum-supplemented medium and are allowed to attach overnight. After serum-starvation for 12 hours in 0.5% FBS, conditions (bFGF, VEGF₁₆₅ or a polypeptide of the invention in 0.5% FBS) with or without Heparin (8 U/ml) are added to wells for 48 hours. 20 mg of MTS/PMS mixture (1:0.05) are added per well and allowed to incubate for 1 hour at 37°C before measuring the absorbance at 490 nm in an ELISA plate reader. Background absorbance from control wells (some media, no cells) is subtracted, and seven wells are performed in parallel for each condition. See, Leak *et al.* *In Vitro Cell. Dev. Biol.* 30A:512-518 (1994).
- 15
20

- 25 The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 38: Inhibition of PDGF-induced Vascular Smooth Muscle Cell Proliferation Stimulatory Effect

HAoSMC proliferation can be measured, for example, by BrdUrd incorporation.

5 Briefly, subconfluent, quiescent cells grown on the 4-chamber slides are transfected with CRP or FITC-labeled AT2-3LP. Then, the cells are pulsed with 10% calf serum and 6 mg/ml BrdUrd. After 24 h, immunocytochemistry is performed by using BrdUrd Staining Kit (Zymed Laboratories). In brief, the cells are incubated with the biotinylated mouse anti-BrdUrd antibody at 4 degrees C for 2 h after being exposed to denaturing solution and

10 then incubated with the streptavidin-peroxidase and diaminobenzidine. After counterstaining with hematoxylin, the cells are mounted for microscopic examination, and the BrdUrd-positive cells are counted. The BrdUrd index is calculated as a percent of the BrdUrd-positive cells to the total cell number. In addition, the simultaneous detection of the BrdUrd staining (nucleus) and the FITC uptake (cytoplasm) is performed for

15 individual cells by the concomitant use of bright field illumination and dark field-UV fluorescent illumination. See, Hayashida et al., J. Biol. Chem. 6:271(36):21985-21992 (1996).

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to

20 test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 39: Stimulation of Endothelial Migration

This example will be used to explore the possibility that a polypeptide of the

25 invention may stimulate lymphatic endothelial cell migration.

Endothelial cell migration assays are performed using a 48 well microchemotaxis chamber (Neuroprobe Inc., Cabin John, MD: Falk, W., et al., J. Immunological Methods 1980;33:239-247). Polyvinylpyrrolidone-free polycarbonate filters with a pore size of 8 um (Nucleopore Corp. Cambridge, MA) are coated with 0.1% gelatin for at least 6 hours

30 at room temperature and dried under sterile air. Test substances are diluted to appropriate concentrations in M199 supplemented with 0.25% bovine serum albumin (BSA), and 25

ul of the final dilution is placed in the lower chamber of the modified Boyden apparatus. Subconfluent, early passage (2-6) HUVEC or BMEC cultures are washed and trypsinized for the minimum time required to achieve cell detachment. After placing the filter between lower and upper chamber, 2.5×10^5 cells suspended in 50 ul M199 containing 1%
5 FBS are seeded in the upper compartment. The apparatus is then incubated for 5 hours at 37°C in a humidified chamber with 5% CO₂ to allow cell migration. After the incubation period, the filter is removed and the upper side of the filter with the non-migrated cells is scraped with a rubber policeman. The filters are fixed with methanol and stained with a Giemsa solution (Diff-Quick, Baxter, McGraw Park, IL). Migration is quantified by
10 counting cells of three random high-power fields (40x) in each well, and all groups are performed in quadruplicate.


The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the
15 invention.

Example 40: Stimulation of Nitric Oxide Production by Endothelial Cells

Nitric oxide released by the vascular endothelium is believed to be a mediator of vascular endothelium relaxation. Thus, activity of a polypeptide of the invention can be
20 assayed by determining nitric oxide production by endothelial cells in response to the polypeptide.

Nitric oxide is measured in 96-well plates of confluent microvascular endothelial cells after 24 hours starvation and a subsequent 4 hr exposure to various levels of a positive control (such as VEGF-1) and the polypeptide of the invention. Nitric oxide in
25 the medium is determined by use of the Griess reagent to measure total nitrite after reduction of nitric oxide-derived nitrate by nitrate reductase. The effect of the polypeptide of the invention on nitric oxide release is examined on HUVEC.

Briefly, NO release from cultured HUVEC monolayer is measured with a NO-specific polarographic electrode connected to a NO meter (Iso-NO, World Precision
30 Instruments Inc.) (1049). Calibration of the NO elements is performed according to the following equation:



The standard calibration curve is obtained by adding graded concentrations of KNO_2 (0, 5, 10, 25, 50, 100, 250, and 500 nmol/L) into the calibration solution containing KI and H_2SO_4 . The specificity of the Iso-NO electrode to NO is previously determined by measurement of NO from authentic NO gas (1050). The culture medium is removed and HUVECs are washed twice with Dulbecco's phosphate buffered saline. The cells are then bathed in 5 ml of filtered Krebs-Henseleit solution in 6-well plates, and the cell plates are kept on a slide warmer (Lab Line Instruments Inc.) To maintain the temperature at 37°C. The NO sensor probe is inserted vertically into the wells, keeping the tip of the electrode 2 mm under the surface of the solution, before addition of the different conditions. S-nitroso acetyl penicillamin (SNAP) is used as a positive control. The amount of released NO is expressed as picomoles per 1×10^6 endothelial cells. All values reported are means of four to six measurements in each group (number of cell culture wells). See, Leak *et al. Biochem. and Biophys. Res. Comm.* 217:96-105 (1995).

The studies described in this example tested activity of polypeptides of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

20 **Example 41: Effect of Polypeptides of the Invention on Cord Formation in Angiogenesis**

Another step in angiogenesis is cord formation, marked by differentiation of endothelial cells. This bioassay measures the ability of microvascular endothelial cells to form capillary-like structures (hollow structures) when cultured *in vitro*.

25 CADMEC (microvascular endothelial cells) are purchased from Cell Applications, Inc. as proliferating (passage 2) cells and are cultured in Cell Applications' CADMEC Growth Medium and used at passage 5. For the *in vitro* angiogenesis assay, the wells of a 48-well cell culture plate are coated with Cell Applications' Attachment Factor Medium (200 µl/well) for 30 min. at 37°C. CADMEC are seeded onto the coated wells at 7,500 cells/well and cultured overnight in Growth Medium. The Growth Medium is then replaced with 300 µg Cell Applications' Chord Formation Medium containing control

buffer or a polypeptide of the invention (0.1 to 100 ng/ml) and the cells are cultured for an additional 48 hr. The numbers and lengths of the capillary-like chords are quantitated through use of the Boeckeler VIA-170 video image analyzer. All assays are done in triplicate.

- 5 Commercial (R&D) VEGF (50 ng/ml) is used as a positive control. b-esteradiol (1 ng/ml) is used as a negative control. The appropriate buffer (without protein) is also utilized as a control.

 The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to
10 test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 42: Angiogenic Effect on Chick Chorioallantoic Membrane

 Chick chorioallantoic membrane (CAM) is a well-established system to examine
15 angiogenesis. Blood vessel formation on CAM is easily visible and quantifiable. The ability of polypeptides of the invention to stimulate angiogenesis in CAM can be examined.

 Fertilized eggs of the White Leghorn chick (*Gallus gallus*) and the Japanese qual (*Coturnix coturnix*) are incubated at 37.8°C and 80% humidity. Differentiated CAM of
20 16-day-old chick and 13-day-old qual embryos is studied with the following methods.

 On Day 4 of development, a window is made into the egg shell of chick eggs. The embryos are checked for normal development and the eggs sealed with cellotape. They are further incubated until Day 13. Thermanox coverslips (Nunc, Naperville, IL) are cut into disks of about 5 mm in diameter. Sterile and salt-free growth factors are dissolved in
25 distilled water and about 3.3 mg/ 5 ml are pipetted on the disks. After air-drying, the inverted disks are applied on CAM. After 3 days, the specimens are fixed in 3% glutaraldehyde and 2% formaldehyde and rinsed in 0.12 M sodium cacodylate buffer. They are photographed with a stereo microscope [Wild M8] and embedded for semi- and ultrathin sectioning as described above. Controls are performed with carrier disks alone.

30 The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to

test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 43: Angiogenesis Assay Using a Matrigel Implant in Mouse

5 *In vivo* angiogenesis assay of a polypeptide of the invention measures the ability of an existing capillary network to form new vessels in an implanted capsule of murine extracellular matrix material (Matrigel). The protein is mixed with the liquid Matrigel at 4 degree C and the mixture is then injected subcutaneously in mice where it solidifies. After 7 days, the solid "plug" of Matrigel is removed and examined for the presence of new
10 blood vessels. Matrigel is purchased from Becton Dickinson Labware/Collaborative Biomedical Products.

When thawed at 4 degree C the Matrigel material is a liquid. The Matrigel is mixed with a polypeptide of the invention at 150 ng/ml at 4 degrees C and drawn into cold 3 ml syringes. Female C57Bl/6 mice approximately 8 weeks old are injected with the
15 mixture of Matrigel and experimental protein at 2 sites at the midventral aspect of the abdomen (0.5 ml/site). After 7 days, the mice are sacrificed by cervical dislocation, the Matrigel plugs are removed and cleaned (i.e., all clinging membranes and fibrous tissue is removed). Replicate whole plugs are fixed in neutral buffered 10% formaldehyde, embedded in paraffin and used to produce sections for histological examination after
20 staining with Masson's Trichrome. Cross sections from 3 different regions of each plug are processed. Selected sections are stained for the presence of vWF. The positive control for this assay is bovine basic FGF (150 ng/ml). Matrigel alone is used to determine basal levels of angiogenesis.

The studies described in this example tested activity of a polypeptide of the
25 invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 44: Rescue of Ischemia in Rabbit Lower Limb Model

To study the in vivo effects of polynucleotides and polypeptides of the invention on ischemia, a rabbit hindlimb ischemia model is created by surgical removal of one femoral arteries as described previously (Takeshita *et al.*, *Am J. Pathol* 147:1649-1660 (1995)). The excision of the femoral artery results in retrograde propagation of thrombus and occlusion of the external iliac artery. Consequently, blood flow to the ischemic limb is dependent upon collateral vessels originating from the internal iliac artery (Takeshita *et al.*, *Am J. Pathol* 147:1649-1660 (1995)). An interval of 10 days is allowed for post-operative recovery of rabbits and development of endogenous collateral vessels. At 10 day post-operatively (day 0), after performing a baseline angiogram, the internal iliac artery of the ischemic limb is transfected with 500 mg naked expression plasmid containing a polynucleotide of the invention by arterial gene transfer technology using a hydrogel-coated balloon catheter as described (Riessen *et al.*, *Hum Gene Ther.* 4:749-758 (1993); Leclerc *et al.*, *J. Clin. Invest.* 90: 936-944 (1992)). When a polypeptide of the invention is used in the treatment, a single bolus of 500 mg polypeptide of the invention or control is delivered into the internal iliac artery of the ischemic limb over a period of 1 min. through an infusion catheter. On day 30, various parameters are measured in these rabbits: (a) BP ratio - The blood pressure ratio of systolic pressure of the ischemic limb to that of normal limb; (b) Blood Flow and Flow Reserve - Resting FL: the blood flow during undilated condition and Max FL: the blood flow during fully dilated condition (also an indirect measure of the blood vessel amount) and Flow Reserve is reflected by the ratio of max FL: resting FL; (c) Angiographic Score - This is measured by the angiogram of collateral vessels. A score is determined by the percentage of circles in an overlaying grid that with crossing opacified arteries divided by the total number in the rabbit thigh; (d) Capillary density - The number of collateral capillaries determined in light microscopic sections taken from hindlimbs.

The studies described in this example tested activity of polynucleotides and polypeptides of the invention. However, one skilled in the art could easily modify the exemplified studies to test the agonists, and/or antagonists of the invention.

Example 45: Effect of Polypeptides of the Invention on Vasodilation

Since dilation of vascular endothelium is important in reducing blood pressure, the ability of polypeptides of the invention to affect the blood pressure in spontaneously hypertensive rats (SHR) is examined. Increasing doses (0, 10, 30, 100, 300, and 900 mg/kg) of the polypeptides of the invention are administered to 13-14 week old spontaneously hypertensive rats (SHR). Data are expressed as the mean +/- SEM. Statistical analysis are performed with a paired t-test and statistical significance is defined as $p < 0.05$ vs. the response to buffer alone.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 46: Rat Ischemic Skin Flap Model

The evaluation parameters include skin blood flow, skin temperature, and factor VIII immunohistochemistry or endothelial alkaline phosphatase reaction. Expression of polypeptides of the invention, during the skin ischemia, is studied using in situ hybridization.

The study in this model is divided into three parts as follows:

- a) Ischemic skin
- b) Ischemic skin wounds
- c) Normal wounds

The experimental protocol includes:

- a) Raising a 3x4 cm. single pedicle full-thickness random skin flap (myocutaneous flap over the lower back of the animal).
- b) An excisional wounding (4-6 mm in diameter) in the ischemic skin (skin-flap).
- c) Topical treatment with a polypeptide of the invention of the excisional wounds (day 0, 1, 2, 3, 4 post-wounding) at the following various dosage ranges: 1mg to 100 mg.
- d) Harvesting the wound tissues at day 3, 5, 7, 10, 14 and 21 post-wounding for histological, immunohistochemical, and in situ studies.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

5

Example 47: Peripheral Arterial Disease Model

Angiogenic therapy using a polypeptide of the invention is a novel therapeutic strategy to obtain restoration of blood flow around the ischemia in case of peripheral
10 arterial diseases. The experimental protocol includes:

a) One side of the femoral artery is ligated to create ischemic muscle of the hindlimb, the other side of hindlimb serves as a control.

b) a polypeptide of the invention, in a dosage range of 20 mg - 500 mg, is delivered intravenously and/or intramuscularly 3 times (perhaps more) per week for 2-3
15 weeks.

c) The ischemic muscle tissue is collected after ligation of the femoral artery at 1, 2, and 3 weeks for the analysis of expression of a polypeptide of the invention and histology. Biopsy is also performed on the other side of normal muscle of the contralateral hindlimb.

20 The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

25 **Example 48: Ischemic Myocardial Disease Model**

A polypeptide of the invention is evaluated as a potent mitogen capable of stimulating the development of collateral vessels, and restructuring new vessels after coronary artery occlusion. Alteration of expression of the polypeptide is investigated in
30 situ. The experimental protocol includes:

a) The heart is exposed through a left-side thoracotomy in the rat. Immediately, the left coronary artery is occluded with a thin suture (6-0) and the thorax is closed.

b) a polypeptide of the invention, in a dosage range of 20 mg - 500 mg, is delivered intravenously and/or intramuscularly 3 times (perhaps more) per week for 2-4
5 weeks.

c) Thirty days after the surgery, the heart is removed and cross-sectioned for morphometric and in situ analyzes.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to
10 test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 49: Rat Corneal Wound Healing Model

15 This animal model shows the effect of a polypeptide of the invention on neovascularization. The experimental protocol includes:

- a) Making a 1-1.5 mm long incision from the center of cornea into the stromal layer.
- b) Inserting a spatula below the lip of the incision facing the outer corner of
20 the eye.
- c) Making a pocket (its base is 1-1.5 mm from the edge of the eye).
- d) Positioning a pellet, containing 50ng- 5ug of a polypeptide of the invention, within the pocket.
- e) Treatment with a polypeptide of the invention can also be applied topically
25 to the corneal wounds in a dosage range of 20mg - 500mg (daily treatment for five days).

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the
invention.

Example 50: Diabetic Mouse and Glucocorticoid-Impaired Wound Healing Models

A. Diabetic db+/db+ Mouse Model.

5 To demonstrate that a polypeptide of the invention accelerates the healing process, the genetically diabetic mouse model of wound healing is used. The full thickness wound healing model in the db+/db+ mouse is a well characterized, clinically relevant and reproducible model of impaired wound healing. Healing of the diabetic wound is dependent on formation of granulation tissue and re-epithelialization rather than contraction (Gartner, M.H. *et al.*, *J. Surg. Res.* 52:389 (1992); Greenhalgh, D.G. *et al.*, *Am. J. Pathol.* 136:1235 (1990)).

The diabetic animals have many of the characteristic features observed in Type II diabetes mellitus. Homozygous (db+/db+) mice are obese in comparison to their normal heterozygous (db+/+m) littermates. Mutant diabetic (db+/db+) mice have a single autosomal recessive mutation on chromosome 4 (db+) (Coleman *et al.* *Proc. Natl. Acad. Sci. USA* 77:283-293 (1982)). Animals show polyphagia, polydipsia and polyuria. Mutant diabetic mice (db+/db+) have elevated blood glucose, increased or normal insulin levels, and suppressed cell-mediated immunity (Mandel *et al.*, *J. Immunol.* 120:1375 (1978); Debray-Sachs, M. *et al.*, *Clin. Exp. Immunol.* 51(1):1-7 (1983); Leiter *et al.*, *Am. J. of Pathol.* 114:46-55 (1985)). Peripheral neuropathy, myocardial complications, and microvascular lesions, basement membrane thickening and glomerular filtration abnormalities have been described in these animals (Norido, F. *et al.*, *Exp. Neurol.* 83(2):221-232 (1984); Robertson *et al.*, *Diabetes* 29(1):60-67 (1980); Giacomelli *et al.*, *Lab Invest.* 40(4):460-473 (1979); Coleman, D.L., *Diabetes* 31 (Suppl):1-6 (1982)). These homozygous diabetic mice develop hyperglycemia that is resistant to insulin analogous to human type II diabetes (Mandel *et al.*, *J. Immunol.* 120:1375-1377 (1978)).

The characteristics observed in these animals suggests that healing in this model may be similar to the healing observed in human diabetes (Greenhalgh, *et al.*, *Am. J. of Pathol.* 136:1235-1246 (1990)).

30 Genetically diabetic female C57BL/KsJ (db+/db+) mice and their non-diabetic (db+/+m) heterozygous littermates are used in this study (Jackson Laboratories). The

animals are purchased at 6 weeks of age and are 8 weeks old at the beginning of the study. Animals are individually housed and received food and water ad libitum. All manipulations are performed using aseptic techniques. The experiments are conducted according to the rules and guidelines of Human Genome Sciences, Inc. Institutional
5 Animal Care and Use Committee and the Guidelines for the Care and Use of Laboratory Animals.

Wounding protocol is performed according to previously reported methods (Tsuboi, R. and Rifkin, D.B., *J. Exp. Med.* 172:245-251 (1990)). Briefly, on the day of wounding, animals are anesthetized with an intraperitoneal injection of Avertin (0.01
10 mg/mL), 2,2,2-tribromoethanol and 2-methyl-2-butanol dissolved in deionized water. The dorsal region of the animal is shaved and the skin washed with 70% ethanol solution and iodine. The surgical area is dried with sterile gauze prior to wounding. An 8 mm full-thickness wound is then created using a Keyes tissue punch. Immediately following wounding, the surrounding skin is gently stretched to eliminate wound expansion. The
15 wounds are left open for the duration of the experiment. Application of the treatment is given topically for 5 consecutive days commencing on the day of wounding. Prior to treatment, wounds are gently cleansed with sterile saline and gauze sponges.

Wounds are visually examined and photographed at a fixed distance at the day of surgery and at two day intervals thereafter. Wound closure is determined by daily
20 measurement on days 1-5 and on day 8. Wounds are measured horizontally and vertically using a calibrated Jameson caliper. Wounds are considered healed if granulation tissue is no longer visible and the wound is covered by a continuous epithelium.

A polypeptide of the invention is administered using at a range different doses, from 4mg to 500mg per wound per day for 8 days in vehicle. Vehicle control groups
25 received 50mL of vehicle solution.

Animals are euthanized on day 8 with an intraperitoneal injection of sodium pentobarbital (300mg/kg). The wounds and surrounding skin are then harvested for histology and immunohistochemistry. Tissue specimens are placed in 10% neutral buffered formalin in tissue cassettes between biopsy sponges for further processing.
30

Three groups of 10 animals each (5 diabetic and 5 non-diabetic controls) are evaluated: 1) Vehicle placebo control, 2) untreated group, and 3) treated group.

Wound closure is analyzed by measuring the area in the vertical and horizontal axis and obtaining the total square area of the wound. Contraction is then estimated by establishing the differences between the initial wound area (day 0) and that of post treatment (day 8). The wound area on day 1 is 64mm², the corresponding size of the dermal punch. Calculations are made using the following formula:

$$[\text{Open area on day 8}] - [\text{Open area on day 1}] / [\text{Open area on day 1}]$$

Specimens are fixed in 10% buffered formalin and paraffin embedded blocks are sectioned perpendicular to the wound surface (5mm) and cut using a Reichert-Jung microtome. Routine hematoxylin-eosin (H&E) staining is performed on cross-sections of bisected wounds. Histologic examination of the wounds are used to assess whether the healing process and the morphologic appearance of the repaired skin is altered by treatment with a polypeptide of the invention. This assessment included verification of the presence of cell accumulation, inflammatory cells, capillaries, fibroblasts, re-epithelialization and epidermal maturity (Greenhalgh, D.G. *et al.*, *Am. J. Pathol.* 136:1235 (1990)). A calibrated lens micrometer is used by a blinded observer.

Tissue sections are also stained immunohistochemically with a polyclonal rabbit anti-human keratin antibody using ABC Elite detection system. Human skin is used as a positive tissue control while non-immune IgG is used as a negative control. Keratinocyte growth is determined by evaluating the extent of reepithelialization of the wound using a calibrated lens micrometer.

Proliferating cell nuclear antigen/cyclin (PCNA) in skin specimens is demonstrated by using anti-PCNA antibody (1:50) with an ABC Elite detection system. Human colon cancer can serve as a positive tissue control and human brain tissue can be used as a negative tissue control. Each specimen includes a section with omission of the primary antibody and substitution with non-immune mouse IgG. Ranking of these sections is based on the extent of proliferation on a scale of 0-8, the lower side of the scale reflecting slight proliferation to the higher side reflecting intense proliferation.

Experimental data are analyzed using an unpaired t test. A p value of < 0.05 is considered significant.

B. Steroid Impaired Rat Model

The inhibition of wound healing by steroids has been well documented in various *in vitro* and *in vivo* systems (Wahl, Glucocorticoids and Wound healing. In: Anti-Inflammatory Steroid Action: Basic and Clinical Aspects. 280-302 (1989); Wahl *et al.*, *J. Immunol.* 115: 476-481 (1975); Werb *et al.*, *J. Exp. Med.* 147:1684-1694 (1978)). Glucocorticoids retard wound healing by inhibiting angiogenesis, decreasing vascular permeability (Ebert *et al.*, *An. Intern. Med.* 37:701-705 (1952)), fibroblast proliferation, and collagen synthesis (Beck *et al.*, *Growth Factors.* 5: 295-304 (1991); Haynes *et al.*, *J. Clin. Invest.* 61: 703-797 (1978)) and producing a transient reduction of circulating monocytes (Haynes *et al.*, *J. Clin. Invest.* 61: 703-797 (1978); Wahl, "Glucocorticoids and wound healing", In: Antiinflammatory Steroid Action: Basic and Clinical Aspects, Academic Press, New York, pp. 280-302 (1989)). The systemic administration of steroids to impaired wound healing is a well establish phenomenon in rats (Beck *et al.*, *Growth Factors.* 5: 295-304 (1991); Haynes *et al.*, *J. Clin. Invest.* 61: 703-797 (1978); Wahl, "Glucocorticoids and wound healing", In: Antiinflammatory Steroid Action: Basic and Clinical Aspects, Academic Press, New York, pp. 280-302 (1989); Pierce *et al.*, *Proc. Natl. Acad. Sci. USA* 86: 2229-2233 (1989)).

To demonstrate that a polypeptide of the invention can accelerate the healing process, the effects of multiple topical applications of the polypeptide on full thickness excisional skin wounds in rats in which healing has been impaired by the systemic administration of methylprednisolone is assessed.

Young adult male Sprague Dawley rats weighing 250-300 g (Charles River Laboratories) are used in this example. The animals are purchased at 8 weeks of age and are 9 weeks old at the beginning of the study. The healing response of rats is impaired by the systemic administration of methylprednisolone (17mg/kg/rat intramuscularly) at the time of wounding. Animals are individually housed and received food and water *ad libitum*. All manipulations are performed using aseptic techniques. This study is conducted according to the rules and guidelines of Human Genome Sciences, Inc. Institutional Animal Care and Use Committee and the Guidelines for the Care and Use of Laboratory Animals.

The wounding protocol is followed according to section A, above. On the day of wounding, animals are anesthetized with an intramuscular injection of ketamine (50 mg/kg) and xylazine (5 mg/kg). The dorsal region of the animal is shaved and the skin washed with 70% ethanol and iodine solutions. The surgical area is dried with sterile gauze prior to wounding. An 8 mm full-thickness wound is created using a Keyes tissue punch. The wounds are left open for the duration of the experiment. Applications of the testing materials are given topically once a day for 7 consecutive days commencing on the day of wounding and subsequent to methylprednisolone administration. Prior to treatment, wounds are gently cleansed with sterile saline and gauze sponges.

Wounds are visually examined and photographed at a fixed distance at the day of wounding and at the end of treatment. Wound closure is determined by daily measurement on days 1-5 and on day 8. Wounds are measured horizontally and vertically using a calibrated Jameson caliper. Wounds are considered healed if granulation tissue is no longer visible and the wound is covered by a continuous epithelium.

The polypeptide of the invention is administered using at a range different doses, from 4mg to 500mg per wound per day for 8 days in vehicle. Vehicle control groups received 50mL of vehicle solution.

Animals are euthanized on day 8 with an intraperitoneal injection of sodium pentobarbital (300mg/kg). The wounds and surrounding skin are then harvested for histology. Tissue specimens are placed in 10% neutral buffered formalin in tissue cassettes between biopsy sponges for further processing.

Four groups of 10 animals each (5 with methylprednisolone and 5 without glucocorticoid) are evaluated: 1) Untreated group 2) Vehicle placebo control 3) treated groups.

Wound closure is analyzed by measuring the area in the vertical and horizontal axis and obtaining the total area of the wound. Closure is then estimated by establishing the differences between the initial wound area (day 0) and that of post treatment (day 8). The wound area on day 1 is 64mm², the corresponding size of the dermal punch. Calculations are made using the following formula:

$$[\text{Open area on day 8}] - [\text{Open area on day 1}] / [\text{Open area on day 1}]$$

Specimens are fixed in 10% buffered formalin and paraffin embedded blocks are sectioned perpendicular to the wound surface (5mm) and cut using an Olympus microtome. Routine hematoxylin-eosin (H&E) staining is performed on cross-sections of bisected wounds. Histologic examination of the wounds allows assessment of whether the healing process and the morphologic appearance of the repaired skin is improved by treatment with a polypeptide of the invention. A calibrated lens micrometer is used by a blinded observer to determine the distance of the wound gap.

Experimental data are analyzed using an unpaired t test. A p value of < 0.05 is considered significant.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 51: Lymphadema Animal Model

The purpose of this experimental approach is to create an appropriate and consistent lymphedema model for testing the therapeutic effects of a polypeptide of the invention in lymphangiogenesis and re-establishment of the lymphatic circulatory system in the rat hind limb. Effectiveness is measured by swelling volume of the affected limb, quantification of the amount of lymphatic vasculature, total blood plasma protein, and histopathology. Acute lymphedema is observed for 7-10 days. Perhaps more importantly, the chronic progress of the edema is followed for up to 3-4 weeks.

Prior to beginning surgery, blood sample is drawn for protein concentration analysis. Male rats weighing approximately ~350g are dosed with Pentobarbital. Subsequently, the right legs are shaved from knee to hip. The shaved area is swabbed with gauze soaked in 70% EtOH. Blood is drawn for serum total protein testing. Circumference and volumetric measurements are made prior to injecting dye into paws after marking 2 measurement levels (0.5 cm above heel, at mid-pt of dorsal paw). The intradermal dorsum of both right and left paws are injected with 0.05 ml of 1% Evan's

Blue. Circumference and volumetric measurements are then made following injection of dye into paws.

Using the knee joint as a landmark, a mid-leg inguinal incision is made circumferentially allowing the femoral vessels to be located. Forceps and hemostats are
5 used to dissect and separate the skin flaps. After locating the femoral vessels, the lymphatic vessel that runs along side and underneath the vessel(s) is located. The main lymphatic vessels in this area are then electrically coagulated suture ligated.

Using a microscope, muscles in back of the leg (near the semitendinosus and adductors) are bluntly dissected. The popliteal lymph node is then located. The 2
10 proximal and 2 distal lymphatic vessels and distal blood supply of the popliteal node are then and ligated by suturing. The popliteal lymph node, and any accompanying adipose tissue, is then removed by cutting connective tissues.

Care is taken to control any mild bleeding resulting from this procedure. After lymphatics are occluded, the skin flaps are sealed by using liquid skin (Vetbond) (AJ
15 Buck). The separated skin edges are sealed to the underlying muscle tissue while leaving a gap of ~0.5 cm around the leg. Skin also may be anchored by suturing to underlying muscle when necessary.

To avoid infection, animals are housed individually with mesh (no bedding). Recovering animals are checked daily through the optimal edematous peak, which
20 typically occurred by day 5-7. The plateau edematous peak are then observed. To evaluate the intensity of the lymphedema, the circumference and volumes of 2 designated places on each paw before operation and daily for 7 days are measured. The effect plasma proteins on lymphedema is determined and whether protein analysis is a useful testing perimeter is also investigated. The weights of both control and edematous limbs are
25 evaluated at 2 places. Analysis is performed in a blind manner.

Circumference Measurements: Under brief gas anesthetic to prevent limb movement, a cloth tape is used to measure limb circumference. Measurements are done at the ankle bone and dorsal paw by 2 different people then those 2 readings are averaged. Readings are taken from both control and edematous limbs.

30 Volumetric Measurements: On the day of surgery, animals are anesthetized with Pentobarbital and are tested prior to surgery. For daily volumetrics animals are under

brief halothane anesthetic (rapid immobilization and quick recovery), both legs are shaved and equally marked using waterproof marker on legs. Legs are first dipped in water, then dipped into instrument to each marked level then measured by Buxco edema software(Chen/Victor). Data is recorded by one person, while the other is dipping the
5 limb to marked area.

Blood-plasma protein measurements: Blood is drawn, spun, and serum separated prior to surgery and then at conclusion for total protein and Ca²⁺ comparison.

Limb Weight Comparison: After drawing blood, the animal is prepared for tissue collection. The limbs are amputated using a quillitinc, then both experimental and control
10 legs are cut at the ligature and weighed. A second weighing is done as the tibio-cacaneal joint is disarticulated and the foot is weighed.

Histological Preparations: The transverse muscle located behind the knee (popliteal) area is dissected and arranged in a metal mold, filled with freezeGel, dipped into cold methylbutane, placed into labeled sample bags at - 80EC until sectioning. Upon
15 sectioning, the muscle is observed under fluorescent microscopy for lymphatics..

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

20

Example 52: Suppression of TNF alpha-induced adhesion molecule expression by a Polypeptide of the Invention

The recruitment of lymphocytes to areas of inflammation and angiogenesis involves specific receptor-ligand interactions between cell surface adhesion molecules
25 (CAMs) on lymphocytes and the vascular endothelium. The adhesion process, in both normal and pathological settings, follows a multi-step cascade that involves intercellular adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule-1 (VCAM-1), and endothelial leukocyte adhesion molecule-1 (E-selectin) expression on endothelial cells (EC). The expression of these molecules and others on the vascular endothelium
30 determines the efficiency with which leukocytes may adhere to the local vasculature and extravasate into the local tissue during the development of an inflammatory response. The

local concentration of cytokines and growth factor participate in the modulation of the expression of these CAMs.

Tumor necrosis factor alpha (TNF- α), a potent proinflammatory cytokine, is a stimulator of all three CAMs on endothelial cells and may be involved in a wide variety of inflammatory responses, often resulting in a pathological outcome.

The potential of a polypeptide of the invention to mediate a suppression of TNF- α induced CAM expression can be examined. A modified ELISA assay which uses ECs as a solid phase absorbent is employed to measure the amount of CAM expression on TNF- α treated ECs when co-stimulated with a member of the FGF family of proteins.

To perform the experiment, human umbilical vein endothelial cell (HUVEC) cultures are obtained from pooled cord harvests and maintained in growth medium (EGM-2; Clonetics, San Diego, CA) supplemented with 10% FCS and 1% penicillin/streptomycin in a 37 degree C humidified incubator containing 5% CO₂.

HUVECs are seeded in 96-well plates at concentrations of 1×10^4 cells/well in EGM medium at 37 degree C for 18-24 hrs or until confluent. The monolayers are subsequently washed 3 times with a serum-free solution of RPMI-1640 supplemented with 100 U/ml penicillin and 100 mg/ml streptomycin, and treated with a given cytokine and/or growth factor(s) for 24 h at 37 degree C. Following incubation, the cells are then evaluated for CAM expression.

Human Umbilical Vein Endothelial cells (HUVECs) are grown in a standard 96 well plate to confluence. Growth medium is removed from the cells and replaced with 90 μ l of 199 Medium (10% FBS). Samples for testing and positive or negative controls are added to the plate in triplicate (in 10 μ l volumes). Plates are incubated at 37 degree C for either 5 h (selectin and integrin expression) or 24 h (integrin expression only). Plates are aspirated to remove medium and 100 μ l of 0.1% paraformaldehyde-PBS(with Ca⁺⁺ and Mg⁺⁺) is added to each well. Plates are held at 4°C for 30 min.

Fixative is then removed from the wells and wells are washed 1X with PBS(+Ca,Mg)+0.5% BSA and drained. Do not allow the wells to dry. Add 10 μ l of diluted primary antibody to the test and control wells. Anti-ICAM-1-Biotin, Anti-VCAM-1-Biotin and Anti-E-selectin-Biotin are used at a concentration of 10 μ g/ml (1:10 dilution

of 0.1 mg/ml stock antibody). Cells are incubated at 37°C for 30 min. in a humidified environment. Wells are washed X3 with PBS(+Ca,Mg)+0.5% BSA.

Then add 20 µl of diluted ExtrAvidin-Alkaline Phosphatase (1:5,000 dilution) to each well and incubated at 37°C for 30 min. Wells are washed X3 with

5 PBS(+Ca,Mg)+0.5% BSA. 1 tablet of p-Nitrophenol Phosphate pNPP is dissolved in 5 ml of glycine buffer (pH 10.4). 100 µl of pNPP substrate in glycine buffer is added to each test well. Standard wells in triplicate are prepared from the working dilution of the ExtrAvidin-Alkaline Phosphatase in glycine buffer: 1:5,000 (10^0) > $10^{-0.5}$ > 10^{-1} > $10^{-1.5}$. 5 µl of each dilution is added to triplicate wells and the resulting AP content in each well is

10 5.50 ng, 1.74 ng, 0.55 ng, 0.18 ng. 100 µl of pNPP reagent must then be added to each of the standard wells. The plate must be incubated at 37°C for 4h. A volume of 50 µl of 3M NaOH is added to all wells. The results are quantified on a plate reader at 405 nm. The background subtraction option is used on blank wells filled with glycine buffer only. The template is set up to indicate the concentration of AP-conjugate in each standard well [

15 5.50 ng; 1.74 ng; 0.55 ng; 0.18 ng]. Results are indicated as amount of bound AP-conjugate in each sample.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists. and/or antagonists of the

20 invention.

It will be clear that the invention may be practiced otherwise than as particularly described in the foregoing description and examples. Numerous modifications and variations of the present invention are possible in light of the above teachings and, therefore, are within the scope of the appended claims.


25 The entire disclosure of each document cited (including patents, patent applications, journal articles, abstracts, laboratory manuals, books, or other disclosures) in the Background of the Invention, Detailed Description, and Examples is hereby incorporated herein by reference. Further, the hard copy of the sequence listing submitted herewith and the corresponding computer readable form are both

30 incorporated herein by reference in their entireties.

307

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>49</u> . line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit 26 April 1999	Accession Number 203957
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
Europe In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g. "Accession Number of Deposit")	
For receiving Office use only <input checked="" type="checkbox"/> This sheet was received with the international application  Authorized officer	For International Bureau use only <input type="checkbox"/> This sheet was received by the International Bureau on: Authorized officer

ATCC Deposit No.: 203957

CANADA

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

NORWAY

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

AUSTRALIA

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

FINLAND

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

UNITED KINGDOM

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

ATCC Deposit No.: 203957

DENMARK

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

SWEDEN

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

NETHERLANDS

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

What Is Claimed Is:

1. An isolated nucleic acid molecule comprising a polynucleotide having a nucleotide sequence at least 95% identical to a sequence selected from the group consisting of:
 - (a) a polynucleotide fragment of SEQ ID NO:X or a polynucleotide fragment of the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;
 - (b) a polynucleotide encoding a polypeptide fragment of SEQ ID NO:Y or a polypeptide fragment encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;
 - (c) a polynucleotide encoding a polypeptide domain of SEQ ID NO:Y or a polypeptide domain encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;
 - (d) a polynucleotide encoding a polypeptide epitope of SEQ ID NO:Y or a polypeptide epitope encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;
 - (e) a polynucleotide encoding a polypeptide of SEQ ID NO:Y or the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X, having biological activity;
 - (f) a polynucleotide which is a variant of SEQ ID NO:X;
 - (g) a polynucleotide which is an allelic variant of SEQ ID NO:X;
 - (h) a polynucleotide which encodes a species homologue of the SEQ ID NO:Y;
 - (i) a polynucleotide capable of hybridizing under stringent conditions to any one of the polynucleotides specified in (a)-(h), wherein said polynucleotide does not hybridize under stringent conditions to a nucleic acid molecule having a nucleotide sequence of only A residues or of only T residues.

2. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises a nucleotide sequence encoding a secreted protein.
- 5 3. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises a nucleotide sequence encoding the sequence identified as SEQ ID NO:Y or the polypeptide encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X.
- 10 4. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises the entire nucleotide sequence of SEQ ID NO:X or the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X.
- 15 5. The isolated nucleic acid molecule of claim 2, wherein the nucleotide sequence comprises sequential nucleotide deletions from either the C-terminus or the N-terminus.
- 20 6. The isolated nucleic acid molecule of claim 3, wherein the nucleotide sequence comprises sequential nucleotide deletions from either the C-terminus or the N-terminus.
- 25 7. A recombinant vector comprising the isolated nucleic acid molecule of claim 1.
8. A method of making a recombinant host cell comprising the isolated nucleic acid molecule of claim 1.
9. A recombinant host cell produced by the method of claim 8.
- 30 10. The recombinant host cell of claim 9 comprising vector sequences.

11. An isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence selected from the group consisting of:
- (a) a polypeptide fragment of SEQ ID NO:Y or the encoded sequence
5 included in ATCC Deposit No:Z;
 - (b) a polypeptide fragment of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z, having biological activity;
 - (c) a polypeptide domain of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;
 - 10 (d) a polypeptide epitope of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;
 - (e) a secreted form of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;
 - (f) a full length protein of SEQ ID NO:Y or the encoded sequence included in
15 ATCC Deposit No:Z;
 - (g) a variant of SEQ ID NO:Y;
 - (h) an allelic variant of SEQ ID NO:Y; or
 - (i) a species homologue of the SEQ ID NO:Y.
12. The isolated polypeptide of claim 11, wherein the secreted form or the
20 full length protein comprises sequential amino acid deletions from either the C-terminus or the N-terminus.
13. An isolated antibody that binds specifically to the isolated polypeptide of claim 11.
25
14. A recombinant host cell that expresses the isolated polypeptide of claim 11.
15. A method of making an isolated polypeptide comprising:
30 (a) culturing the recombinant host cell of claim 14 under conditions such that said polypeptide is expressed; and

(b) recovering said polypeptide.

16. The polypeptide produced by claim 15.

5 17. A method for preventing, treating, or ameliorating a medical condition, comprising administering to a mammalian subject a therapeutically effective amount of the polypeptide of claim 11 or the polynucleotide of claim 1.

10 18. A method of diagnosing a pathological condition or a susceptibility to a pathological condition in a subject comprising:

(a) determining the presence or absence of a mutation in the polynucleotide of claim 1; and

(b) diagnosing a pathological condition or a susceptibility to a pathological condition based on the presence or absence of said mutation.

15

19. A method of diagnosing a pathological condition or a susceptibility to a pathological condition in a subject comprising:

(a) determining the presence or amount of expression of the polypeptide of claim 11 in a biological sample; and

20 (b) diagnosing a pathological condition or a susceptibility to a pathological condition based on the presence or amount of expression of the polypeptide.

20. A method for identifying a binding partner to the polypeptide of claim 11 comprising:

25 (a) contacting the polypeptide of claim 11 with a binding partner; and

(b) determining whether the binding partner effects an activity of the polypeptide.

21. The gene corresponding to the cDNA sequence of SEQ ID NO:Y.

30

22. A method of identifying an activity in a biological assay, wherein the method comprises:

- (a) expressing SEQ ID NO:X in a cell;
- (b) isolating the supernatant;
- 5 (c) detecting an activity in a biological assay; and
- (d) identifying the protein in the supernatant having the activity.

23. The product produced by the method of claim 20.

1

<110> Human Genome Sciences, Inc.

<120> 48 Human Secreted Proteins

<130> PS526PCT

<140> Unassigned

<141> 2000-03-22

<150> 60/126,595

<151> 1999-03-26

<150> 60/171,549

<151> 1999-12-22

<160> 144

<170> PatentIn Ver. 2.0

<210> 1

<211> 733

<212> DNA

<213> Homo sapiens

<400> 1

```

gggatccgga gccc aaatct tctgacaaaa ctcacacatg cccaccgtgc ccagcacctg      60
aattcgaggg tgcaccgtca gtcttctctt tcccccaaaa acccaaggac accctcatga      120
tctcccggac tcctgagggt acatgctgtg tgggtggacgt aagccacgaa gaccctgagg      180
tcaagttcaa ctggtacgtg gacggcgtgg aggtgcataa tgccaagaca aagccgcggg      240
aggagcagta caacagcacg taccgtgtgg tcagcgtcct caccgtcctg caccaggact      300
ggctgaatgg caaggagtac aagtgc aagg tctccaacaa agccctccca acccccatcg      360
agaaaacccat ctccaaagcc aaagggcagc cccgagaacc acaggtgtac accctgcccc      420
catcccggga tgagctgacc aagaaccagg tcagcctgac ctgcctggtc aaaggcttct      480
atccaagcga catcgccgtg gagtgggaga gcaatgggca gccggagAAC aactacaaga      540
ccacgcctcc cgtgctggac tccgacggct ccttcttctt ctacagcaag ctcaccgtgg      600
acaagagcag gtggcagcag gggaacgtct tctcatgctc cgtgatgcat gaggctctgc      660
acaaccacta cagcagaag agcctctccc tgtctccggg taaatgagtg cgacggccgc      720
gactctagag gat                                     733

```

<210> 2

<211> 5

<212> PRT

<213> Homo sapiens

<220>

<221> Site

<222> (3)

<223> Xaa equals any of the twenty naturally occurring L-amino acids

<400> 2

Trp Ser Xaa Trp Ser

1

5

<210> 3
<211> 86
<212> DNA
<213> Homo sapiens

<400> 3
gcgcctcgag atttccccga aatctagatt tccccgaaat gatttccccg aaatgatttc 60
cccgaaatat ctgccatctc aattag 86

<210> 4
<211> 27
<212> DNA
<213> Homo sapiens

<400> 4
gcggcaagct ttttgcaaag cctaggc 27

<210> 5
<211> 271
<212> DNA
<213> Homo sapiens

<400> 5
ctcgagattt ccccgaaatc tagatttccc cgaaatgatt tccccgaaat gatttccccg 60
aaatatctgc catctcaatt agtcagcaac catagtcccc cccctaactc cgcccatccc 120
gccctaact ccgcccagtt ccgcccattc tccgccccat ggctgactaa ttttttttat 180
ttatgcagag gccgaggccg cctcggcctc tgagctattc cagaagtagt gaggaggctt 240
ttttggaggc ctaggctttt gcaaaaagct t 271

<210> 6
<211> 32
<212> DNA
<213> Homo sapiens

<400> 6
gcgctcgagg gatgacagcg atagaacccc gg 32

<210> 7
<211> 31
<212> DNA
<213> Homo sapiens

<400> 7
gcgaagcttc gcgactcccc ggatccgcct c 31

<210> 8
<211> 12
<212> DNA
<213> Homo sapiens

<400> 8

ggggactttc cc

12

<210> 9

<211> 73

<212> DNA

<213> Homo sapiens

<400> 9

gcggcctcga ggggactttc ccggggactt tccggggact ttccgggact ttccatcctg
ccatctcaat tag

60

73

<210> 10

<211> 256

<212> DNA

<213> Homo sapiens

<400> 10

ctcgagggga ctttcccggg gactttccgg ggactttccg ggactttcca tctgccatct 60
caattagtca gcaaccatag tccgcccct aactccgccc atcccgcccc taactccgcc 120
cagttccgcc cattctccgc cccatggctg actaattttt ttattttatg cagaggccga 180
ggcgccctcg gcctctgagc tattccagaa gtagtgagga ggcttttttg gaggcctagg 240
cttttgcaaa aagctt 256

<210> 11

<211> 1626

<212> DNA

<213> Homo sapiens

<400> 11

ggcacgagct gcagtctccc tagcatctgt tattttattga ctttttaata acagccattc 60
tgaccgctgt gaaatggtat ctcatatgg ttttgatttg catttctcta attgttagtg 120
atgtggaaca ttttttcata tgtttgttgg ctccctgtat gtcttctttt gagaagtgtc 180
tgttcatgtc ttttgcccag ttttaaatgg gatttgtttt ttgcttggtc acttggtcac 240
actttttttt tttttataga ttctggatgt tagaccttg tcagatgcat aatttgcgaa 300
cattttttct attttgtagg ttgtctgttt actccactga aagtttcttt tgtagtgcag 360
aagctcttta attaggtctc acttgccaat ttttattttt gttgcagttg cttttaagca 420
cttagtcata aattctttcc cagagccgat atctagaatg gtgtttccta ggttttcttg 480
taaaattctt atagtttgag gtcttacact taaatattta atccatcttg agttaatttt 540
tgtatatggt gaaaggtagg ggtcctttca ccatagaata aaacggtgtt tcattctttt 600
gcatatggct agccagctat ctacgcacca tttactgaat agggaatcct ttccccattg 660
cctatttttt gttgactttg tcaaagaaca ggtggctgta ggtgtacagc tttatttctg 720
ggttctcaat tctgttccct tggctcgttt gtctgctttt gtaccagtac catgctgttt 780
gggttactgt agctttatag tatagtataa agtcaggtaa tgtgatgcct cagctttgtt 840
ctttttgctt gggattgctt tggttatttg ggttcttttt tggttccata tgaatttcag 900
aatagttttt tctagtcttg tgaaaaatga cactgggtcat ttgataggaa taacattgaa 960
tctatagatt gctttgagaa gtatagccat ttttaacaata ttgattcttg taatccatga 1020
gcatggaatg tttttccatt tgtctgtgtc atctgtgatt tctttcagca gtgtttccta 1080
gttctctttg taaagatcct tcacctcctt ggttagatgt atttctaggt actttgtttt 1140
tttgatggct atcgtaaaca ggattgtgtt cttcatttgg ctctctagct tggatgttat 1200
tgggttatag aaatgctact gatttttcta cattgatctt gtatcctgaa actttaccaa 1260
agttgtatgt cagttccagg agccttttgg tggagtctt aggggtttccc atgtgcagaa 1320

tctcatgatt ccttaaaagc atgcatttct acttaaacca tcatgtttac ttttctagag	1380
agcaattaac ttggagggtgg gtgccgggga ggtaggttg cttttgtaat attaatggat	1440
gtacaccaag aatattgctt ctgagaatga tcttatcctc attgggaaag atttttctgt	1500
ttttagttga aattgagatg aaatacatct tattataaat aaattttgac tcttactaat	1560
gattacagga ttgtagacaa ttaactgtct tcctcatgct gagtacataa aaaaaaaaaa	1620
aaaaaa	1626

<210> 12
 <211> 2397
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (155)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1195)
 <223> n equals a,t,g, or c

<400> 12	
cttttttttt ttttttttag gtggaatcaa agcaagttgt cttcagagac tgggatccga	60
gatagaaaac acacagtga gtttaatcag gaaccaaac tccggtcctc tgctacaacc	120
acggaaacgg ctccaaactt gaggggggac cccnaacgc ctgcttttgg cccaaagctc	180
tgccctccag cctcctcat acccactggc cacctaggac caggaaagg gggtagagcc	240
ctgagaattc tgggtctggg gtcaccagct cccacacctg tgctccccgg cccacacac	300
atgatgcccc ggggtgggca atccctgaca gcggtggcgg gcacttggga gctcctgctc	360
agccacctgc cacggccccc cctggggggtc cggcaggagc cagggcagt catggcagca	420
taaggccccc ctgcagatcg actgccttca gaaacaaaa gtcccggcgc aaaggcggtc	480
ccggagtggc agcctggcct gcacccacgc tgtgctgccc ctgcagagcc ccagcagcga	540
gscacaccca ggtcagggga gggggcttgg gtaccagggg cctcactggc tcttcaccag	600
gaccctgtag agtgagaaag tgaggactgc ggccacggcg gccccgacaa cccccagcag	660
ccccggagc cagaaggga agggatgcag ctctgcgtgg accaaatgtg ggaaggcggc	720
catggtggcg agctgggtga agatgggtgt gctgggctcg gctgggcccag cacaggagaa	780
cggcacggga gcgggtagcc ggtgcttgcg gcaaaactcg gccggtgat ggcagacac	840
cgcgacactt cgggcaggct ggccctggag gagacaaaga ggcagggggt ctgcccgctc	900
atgtaattgt gcttgttagc gctggcacaa tgtgcaaagg actttgggtc actgccatca	960
aacatcaagc aggaacgtc acaggtggcg tccagcgatg tggccagcag accatctgtg	1020
cccacctcac agaggatcaa gtacttctcc tgcccattga cctgcaccgt gtcgatggcg	1080
tagccgggag gctgctccct cgtgtcctgg tgcccaggc cgcggccgag aaaggcctgc	1140
aggaaggcag acttgcccac tccacgggccc cctaccacct tgcacaggag gamgnrccgc	1200
tgctctgtc ccttctcctg gtccagcctc ttctcacgag tgactgtgat ggcattggcc	1260
tggtcctgct cacagagggg ggggtagccc aggtagccta ggtgtccaag gcagctccgg	1320
acgtccaggt aggtcaccag ggtccactgg cagaggatc cgtgcagggg caaccggcgg	1380
gcctctgtgc ggactgtgcg tgggagctcg gggccccagg gcgctgtctg gaacacactg	1440
aaaaggcttt gcagctccac gggcgagagg gcgcgctcgc ggtcctggtc gtgcttctca	1500
aacactctct gcacaaactg gttagccaag ttgttgagct ccgtgctgca gccggggggc	1560
acgtggatca gaggggagag atagtccgca gtcagctcca gggcatcgct gtagccgaag	1620
cgcgcagga tgggtccaggt ggtctcgtgc cgcccgcgct ggatgaagag cgtgttcagg	1680
aagaggaaac ctgcaccgag acaagcagg gggagcacca ggcacagccc cctccccaca	1740
ctccccaggc acaggcgggc acccggcctc accatccagg gtcagcyggt cctcccgcac	1800
gccgcccggc acattcctgc acaccaccgt cttcacgtcc tccagggcct gcggggccag	1860

ggggtgccya	aagcaggatt	tctggaaagc	gttgagctct	tcgtcactga	gcgcctggtc	1920
caggtcctga	tctgagagcc	tgaagatgcg	cgtcagcgcc	tgggcgcacg	cgggcctcaa	1980
ctgcttgccc	tcagggtcat	agaggggggc	tgtgggatgc	aggacggcct	tctgggcgta	2040
gtagaacagc	tctgagatgt	tcctcagggt	cttggccgaa	cactccacgc	aggtctcaat	2100
ctcgggaaac	tgggtcatga	tggggagcac	ggcctccatg	gagctccccg	accgcaggtc	2160
tgacttggtg	cccactagga	tgatgggcac	cctggggccc	tgctggtgcc	ccccattcac	2220
cagtggtgatc	cacttagttc	gaatcttctc	aatggtggcc	tcctcagaga	cgtcatacac	2280
cacacacacc	acgtttgcct	tgtggatctc	ctcccgagc	tcctcgtccg	tctgctcggc	2340
ttctgagtag	tccacgatgt	gggtgggcac	cttctccggg	gtgacgtccc	tcgtgcc	2397

<210> 13
 <211> 814
 <212> DNA
 <213> Homo sapiens

<400> 13	
ccacgcgtcc	gccgggcggg
gccgcggg	agccgctgca
caagtgcgcg	ccgaggtcac
cgcagggaga	aggaagaggt
aagcacaaagt	tcacgaaagc
agaccacagc	aagattcttt
accctggaca	tcaaaggagg
cacgtcttgg	tgcttctctt
attctcgtaa	acgcagcctc
gctgctctga	gatttttgagt
aaaacctgct	gaacatttta
ataattccag	aaagtcacac
caggatgaga	atctaagata
aaaaaaaaaa	aaaaaaaaaa
	aaaa
	60
	120
	180
	240
	300
	360
	420
	480
	540
	600
	660
	720
	780
	814

<210> 14
 <211> 641
 <212> DNA
 <213> Homo sapiens

<400> 14	
ggcagcaggt	tccttgctac
caaatgaaaa	cattgcaaga
gcagcatggt	gctgacattg
gtacacttag	ttgctagtgt
ttgtttttgc	tcagggttct
gtctacatta	tgtctgaagg
aatgtttgta	aatattttat
gggcggctgg	ggtggaattg
gtatgtaaca	gtgatttgct
agtggttgtc	ttgactgggt
aataaaaaaa	ctgcatactt
	caaaaaaaaa
	aaaaaaaaaa
	a
	60
	120
	180
	240
	300
	360
	420
	480
	540
	600
	641

<210> 15
 <211> 2163
 <212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (2118)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (2136)

<223> n equals a,t,g, or c

<400> 15

```

tttgcgtgcc ggctgcctak ggtctgggaa gctcgggcac cctccctctc cggggctcct      60
gctccacccc ctccggcccc ccacccgctc cgctccctcca ggctgggcct gtggccgcgg      120
tgcttttttaa ttttccccc gctcagaatc ttgctgctcg gccccagga gagcaacaac      180
tcaacgggaa cgatgtggaa ggtgtcagct ctgctcttcg ttttgggaag cgcgctcgctc      240
tgggctcctgg cagaaggagc cagcacaggc cagccagaag atgacactga gactacaggt      300
ttggaaggcg gcgttgccat gccagggtgc gaagatgatg tggtgactcc aggaaccagc      360
gaagaccgct ataagtctgg cttgacaact ctgggtggcaa caagtgtcaa cagtgttaaca      420
ggcattcgca tcgaggatct gccaaacttca gaaagcacag tccacgcgca agaacaaggt      480
ccaagcgcca cagcctcaaa cgctggccacc agtcactcca cggagaaagt ggatggagac      540
acacagacaa cagttgagaa agatggtttg tcaacagtga ccctggttgg aatcatagtt      600
ggggctcttac tagccatcgg yttcattggg ggaatcatcg ttgtggttat gcgaaaaatg      660
tcgggaaggc cctaaagagc tgaagggtta cgccctgctg ccaacgtgct taaaaaaga      720
ccgtttctga ctctgtgccc tgtccctgag ctcggtggag aagatgaccc gtggaacact      780
tgccctggccc actcagaatc cacggtgacc tctccgcttg ccaaataaac cgaagaaaga      840
ccgttcacca gacttggtc ctctaaacat ttgctgttca aacatgtttt tgaatataca      900
ttctataaaa gattatttga aagacaaat tcatagaaaa tggagcaaaa ctgtataaac      960
tgatttgtaa ctaacactgg accattggat cgatattaya tgctgtaacc atgtgtctcc     1020
gtctgaccat tcttgttatt gttaaaatgc agaggaatct ggaaatattt atatccacgg     1080
agtccttggg tccagtgtca cgtcagtaaa tagcaccagc attttgcaat tgctgatctg     1140
ctgaaatgta cacattctgg tctagtttgg tctatctttt aaagcctgat ctggtgtgaa     1200
taatcaacta ggaaatctaa acttgataaa cacgtgggtga acaactgcct ttagctggtc     1260
cagattaatc atttcaaaga catccatttt agatcacaa caggaagtcg atagtctcaa     1320
aggcactttg tttctcccaa gtaggccacc aggcagctc tagagttgct ttacccaaat     1380
ccttctccag ccattgactg gtgactctaa gcttgctccc acctgcccc tccacttccc     1440
tcagatgatg aggagccagg gctaaggggg cagccttctc tcttcccagt gatgcacatc     1500
cttcacattg gctgctttgt tctggaatat ggatatctca gcctggatgc cgaggaagct     1560
gctggatgct taatgggtgt agaggctcaa gtgtgtttga aaccaagagc cagttgtccc     1620
ccatgcagaa agaaaatctg tgtgagcctc tggatgaga aataaaatct gccagtttta     1680
taacattcac tttctgcctc tgaggaaaga tacagggaac aaaaatcaat ttgtacagtc     1740
ttaatattaa aagcagcttg actaaatacc tgatttaaaa atagaagaca tccccagtcc     1800
tcatgacata cgcgaaatat ctgtggggtc ctgttgaaaa gaacaaaata aaggagccca     1860
aggggtcatt ctgtctcagc accatccagc ctggcacttc tcttcccata tatecattgg     1920
atTTTTTTTT ttttttccct aaacaaagtt ttacactga gcagatgctc tgtcatgatg     1980
gcggttgtgc aattctggta tcctctaaat ttgtaagcat tcataaaaaa aaaaaaaaaa     2040
aactcgaggg ggggcccggw cccaattgcc ctataggag tcgtattaca attcactgsc     2100
cgcgttttac aacgtcgnga ctgggaaaac cctgnggtta cccaacttaa tcgccttgca     2160
gaa

```

<210> 16

<211> 1312

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (1168)

<223> n equals a,t,g, or c

<400> 16

ttgcaggaat tgcgcacagt cacggcatcc cactctgttc tgccgaggac gtgaatcttt	60
cctttgttca gtatatccca tgctgtatac actacctgcc tgtcagtcac tcagtagcgg	120
cctatgttat cagattggct gatcattgta ttgcagtgtc atgttcagggt aacacttatt	180
ttacttatag tgggtcccaag gtgcaagagt agtgatgctg acatattatt ataacttcta	240
ttttattatt ctattgtta atcttactat gcctaattta taaattaaac ttaatcatag	300
gtatatatat ataggaaaaa catagtgtta tatagggttc aatactagcc ccagtttcag	360
ttgggggtct tgggaatgtgc cccccacaga taagagggga ctatatctct atagaagaag	420
agtttgattt tttaggacgt tatttcctgt tcaaaccatt gacctactct ttgagataag	480
ttattctgaa gctttcatat aagtaaaagc aattaaattt tgccaccagc accatcctca	540
accattctga attataaagt gttaaaatta agaacaggaa gaatggaagt attcaataaa	600
aaatgaagta ttttcagttc catattgtca tcaaataagg gtgggaaagg caaaggagaa	660
aatcaataag tatggagaag aaagaaatga gtagatagaa caaaaaagat gagaaaggaa	720
gccatattga agtcaggaga gagtgcagag caagaaagga tgggcagaaa aaaagaaaac	780
aagaaaggag tttaatgagt gagtagaatg tgctcttgtt gcattctact gccagtctaa	840
gccatggaat agttcactat tctgggaagt attcttaaga tcatcagtga ccttcaggtc	900
atgaaatcta ataagcattc tttcagacct gttcttattt gacctctcag tgggtgttga	960
cactctcaac caccaactct ttgtgaaacc tttttttccc tgaaagctgt gaggtcgtgc	1020
tttctcggtt ctcttctcat gtctctggtg actcctgctt tgtctacttt gggttttttg	1080
ttttgttttg ktttttgktc ttaagagtt gggatcttgc tctgctgcct aggcctggagt	1140
gcagtggcgc aatcatagct cactgcanc cccacactyct gggcttaagt gatctacctg	1200
ccttggccty ctaagtaact gtgttcattt gacacagttt aggaactaaa gatcaggtaa	1260
aaccagccag ttcttgagct ggaattgaat ctgttcagaa tgtctgggaa ct	1312

<210> 17

<211> 1726

<212> DNA

<213> Homo sapiens

<400> 17

ggcacgagtt gaggaggagg gagcgcttga aggggactgg cctggcgtgc actccgcacc	60
tcggggacat tattgcgcgt ggaacggctg cttttggaag gcacaacttc ctgaatggac	120
catgactccc accaaagatc cctgtctctg attcaccaaa cagcttcaac cctgaaacca	180
ggacgagaag ttgacaacat ctgagtggac agctaattga cctaagactt cagaccagge	240
ctgtatgac tcctgtctaa acattcttag agtattatat ttactttggg gactattggc	300
cttgtctgct ttgactcaga ttataggata tataacctgg ttaatgtttc tgtacacatg	360
atggccacct atgtatatac acttgacttt tcagggtctc tgcctgggat ggaaaaatag	420
ttcattagcc aaactctcct aaagtgtggc aatggaggca gtctctcaga ttgctggatt	480
tgtcaccatc ttccaagggc cattcaagat cattacatgc tacttgttgc accagagatt	540
gatttttctg acaactccaa tgttaccata tacttacatc agttccctcc aaatgtcact	600
ttccaaattc aaattcagct tctaaagcca ggccatattt tcattctgtg cttagtaata	660
agccctaagt ctcacccttc cccaccgccc ccaggtctat ctatcagtaa gccaaatagg	720
gacaaattct ctaacttate tagaacacaa acttaattcc tcacttttcc caggctccctg	780
cacattctct gcctcccacc gttactgtca taactggcaa ctgaatgtcc aaaggagaaa	840
aagcaaagca gattgtgcca gcacgtctca tttgttatat atcatgtcta tgcaggcctt	900
tgttaaggcct gcagtaaaac caatccatgg ctcaactcatg ttttaaacaa cttacctatt	960
caaactatta tgaatgaata gttaagagat acatttttgtt atgtggcatt gggccacacc	1020

caccaactca	aggatgggca	catagtgtaa	atagctggca	aaatagctgg	caaattgaag	1080
ggcgcttttt	tttttttttt	ttttgaggga	gagtccttgc	ctttcgccca	ggttgtagtg	1140
ggtgggtgtga	catggctcac	tacagccttg	acctcctggg	ctcagggatc	ctcctacctt	1200
agccttccga	gtactgggcc	aaggagctctg	tgtagtagca	tcaagagcct	gtgccttaca	1260
tcaacacctc	gggtgaagta	gagacttgca	cagaaagaat	ttccaagcaa	gctaaatggg	1320
tacagaggca	caaactactg	atcttctcaa	tgactattta	ttggcttcc	atttcttcca	1380
gtctgcagtt	caaccctttt	ttcatgggta	aagtgccgac	gtcataacaa	ggtttgaggg	1440
aggcactttc	accgtgtgaa	tgcccagtc	tcacacttac	aaatgacaaa	aggattcaaa	1500
ccttgatata	gtctctatcc	tactttctta	ctatcactta	aattactcat	ggtatttata	1560
accacttacg	taagtctatt	gctaaaacca	aattataggt	caggtgcagt	ggctcacatt	1620
caatcctagc	acttggggagg	ccaaggtggg	aagactgctt	gaggctagga	gtttgagacc	1680
agcgtgggca	atgcagtga	accctatctt	gataaaaaaa	aaaaaa		1726

<210> 18

<211> 2006

<212> DNA

<213> Homo sapiens

<400> 18

ggcacagctg	aggaactgaa	aagaaatgct	gagacaggaa	atctgcctca	ttcgtaccgg	60
ctcatcagtg	ttgtcagtc	cattggtagc	acttcttctt	caggtcatta	cattagtgat	120
gtatatgaca	ttaagaagca	agcgtgggtt	acttacaatg	acctggagggt	atcaaaaatc	180
caagaggctg	ccgtgcaagt	gatcgagatc	ggagtggcta	catcttcttt	tatatgcaca	240
aggagatctt	tgatgagctg	ctggaaacag	aaaagaactc	tcagtcactt	agcacgggaag	300
tggggaagac	taccctgcag	scctcgtgag	gaacaaactc	ctgggttggc	agcatgcact	360
gcatatttgt	tactgtgcc	cacctcacct	ttctctgct	gaaggagaat	ttggaattct	420
acttgatgcg	ggagcaacaa	acagctcagg	gccaaaccaa	aagacaaaaa	ttggagtaac	480
gtagaatgct	ccatgctatt	ttatggaaac	tttgggtctca	catccgtagc	tgattatcct	540
ctttttctcc	tatgagtggc	acttcttttg	tcttaggaat	acatgttgta	aatatatatc	600
tgtgtatgtg	tgtatacaca	cacacagaca	cacacacaca	cacacgggat	gaatggagcc	660
ttaaagagtt	aggatgagcc	accagaatat	gcctgctcaa	aattaatagc	acagcagttt	720
ggagaagaaa	tgaagggtgc	aaagagtcca	ttcacctgag	aaatgtgtga	agacatactt	780
atcagttggc	ttttagcttt	tatgttcctt	gagttagttc	actcaagtct	gtaacctttt	840
gtgtttcctt	attagtaaaa	ttcactggaa	agccagctct	tcagtgtaca	ctaataacag	900
tttgttctct	ttgcaagaga	ggggcattac	tgtaacctga	cttgaggagc	tgttttgttg	960
ttgttggtgt	ctgcaaattt	catgaatttg	tgatgtcttt	gctgtttaca	tgagtcacca	1020
agaaatggat	tgttgtgtgt	ttggaatatg	ttacagtcct	acatttgata	tttcttatat	1080
actttgtttt	ctctaaggag	atttcttcac	acagtatgtt	catcatatat	catcatcatt	1140
attatggtgg	taaagataga	atcttttttc	ttttttgtca	ttctgscatg	gagcagcatt	1200
accctaattg	attgcaacca	aaactttaaa	caagtagaaa	gataatattt	ctccaattgg	1260
gactccccag	caggataact	tagggataag	gaagaatgct	agcatctctg	tctctcarac	1320
atagggagga	taagaagagt	gktcttctgg	kaaagctaaa	attctggacc	actgaagcta	1380
aaagccctat	tgcaagtatg	aaattaagta	cttgagctat	aggacaaacc	ttgggcattt	1440
aaccatttac	tgtctggctt	tgcccttaaa	ataggggtgc	aattaaaatg	tgattggctt	1500
aggtaatccc	aaaaactaac	aaataacaaa	ggtgcataat	ttatttatct	actttttagg	1560
tgctctgagt	tgaggcaaa	tagagcggca	acattaagt	ctatgctagt	cacttagctg	1620
acgtaaccag	cttggttaag	cagcttatga	aaccatataa	agaattcttt	tgaggatgga	1680
attctgtcca	caaaataatt	ttgtgagccc	agatatcatt	aggatcacac	agagttaaat	1740
atagaaaaat	gaaaccatca	ttatatctct	tcgtgttttt	tcttttatta	taaacaaggg	1800
gattattctt	tagttctcag	aggtagggac	aaaaccacat	cagggttttca	gaaggaaaaa	1860
acattttaaa	accaccatc	acatgagaga	atcacttgaa	cccaggaggc	agagggttga	1920
gtgagctgag	atcgcatcat	tgactgcag	tctgagtgac	agagtggagac	tccatctcat	1980
taaaaaaaaa	aaaaaaaaaa	ctcgag				2006

<210> 19
 <211> 1607
 <212> DNA
 <213> Homo sapiens

<400> 19
 ggcacgagcg gcacgaggaa gcagccacgc ctggaacaa ttaaccagcg tatttctggc 60
 tttgtggatg gatggatgat ggcacccata tggagctttg gacctaaact ttatgtggat 120
 aagagttggc tttttggtct tcaagacccc tggcctgcgt actccagcag caggagagcg 180
 gatttacaac atctcaggga atggcagccc tcttgctgac agcaaagaga tcttcctcac 240
 tgtgccagtg ggcggcggag agagcctgcg attattggcc agtgacttgc agaggcacag 300
 tattgcccag ctggatccag aggccttggg aaacattaaag aaagtctcca accgtctcgc 360
 ccaaactctgc agcagcatatc ggacccacaa atgagacacc aaagttagaca ggatggactt 420
 ttaattggga cttctgggac cctgaagaga cttcttccct tcaggcttat tgtttgagtg 480
 tgaagtcca gagcaaggag ccatgttccct ctaagggaat tcaggaattc agacgtgcta 540
 gtccacacacc agttaggtag agctgtctgt tcacctccc atcccagctg atcccagtca 600
 ctgcttgctg gggccatgcc atggaagctt cccatcagtc tcccagctga atcctccctg 660
 ctctctgagc tgctgccttt tgcctcctgc aactcaacat cctcttcacc ctgccctgcc 720
 tgcagttgag ggggcgaaga agaaccctgt gttctcagga agactgcctc caccaccgct 780
 acccagagaa cctctgcac tggcatttct gctctctatg cttgagaccg ggagggttag 840
 gctcagataa ctgagctctg ggccatgaga gggtaggtcc agaaggtggg gggaactgta 900
 cagatcagca gagcaggaca gttggcagca gtgacctcag tagggaacat gtccgtctac 960
 cctctcgac tcatgacacc tccccctacc agccctcctc ttcctcctcc tctcctcct 1020
 gtggaggtgg tcagtgggac ttagggatct ttcacctgct gtgcccagta gttctgaagt 1080
 ctgcttgctg agcagtgttt tatgtttatc cctgtttact gaagaccaa tactggtttg 1140
 gagacaactt ccatgtcttg ctcttctacc tcctagtta gtggaaattt ggataagggg 1200
 actgtagggc ccagattctg gaggttttat gtcattggcc acagaataac tgtctctaag 1260
 ctatccatgg tccagtggtc cctgccaaagt ctgtagactt cagagagcac ttctctctta 1320
 tggggttcac gggaacaggg ggcgggtgtga cttgcttggg ggccctattc catgtgtgcc 1380
 tgtgcctggg gcatggactt tgttaagcag agtcagcagt gaggtcctca ttctccagcc 1440
 agcctctctg ccctggagaa tcatgtgcta tgttctaaga atttgagaac tagagtctc 1500
 atccccaggc ttgaaggcac atggctttct catgtagggc tctctgtggt atttgttatt 1560
 attttgaac aagaccattt tagtaaaaca aaaaaaaaa aaaaaaa 1607

<210> 20
 <211> 1402
 <212> DNA
 <213> Homo sapiens

<400> 20
 ggcacgagac gctctggacg agcgaccagc aggcagacga tggcggcgaa ggcaacaatt 60
 aaggccccag gggaactggc agcgcacgcg gatgctacta ctgcagtctt tatttttttc 120
 ccatgagttg ggggtcgggt gggggaggga aaggaggga tgaccttccc agggagaaac 180
 ccacgacctg tctgtctttt gatcgctctt ttgacatttt tgccaaaata ccactagtgg 240
 aaagtcaggc tagctgtgct ggtattggaa tagcagcctc acactggcgt ctggactgtt 300
 ctgtagattc atgcaagtgg agctgtctgt ctctaattta acttattgct agataatagg 360
 gttttcagat gaaaagaaaa cttaaaggag aatggccctc attcagtaag ttctgtgggt 420
 ccagtaagga tttttatgta catacgctct cgtctctcgt tttgggtact ttctatctca 480
 tctgtctcgg ctctgcatgt tttccagggt gtacccata gacatggaac agtgtaaatc 540
 ccagactgac agacttagaa cctgaggtct cattcatcct tatggttttag gccttgccag 600
 ttttccgaag tctctgatta gttgacagta ttaacactaa attgcagttt acagtatttc 660
 tacattacag ccatatgtaa catcaagcca tcgattgtgt acttttctt tgctagtgtt 720
 ttgggcttta acatccttat tcagccttat ccaggttggg tttgctgttg atcgggtctc 780

```

taggctaaat gagaatgaaa gcgacttcag gtttttgggt tcataagggt ctcggcaagg      840
tggctgtggg aatttttttt ttttggctct tctttcctct taacgtaaat ccaccaccaa      900
aattattaat cctcttgaaa agaaaacgtg aaacgccaca aaaatagaga aaattcaggt      960
ctgtatgtca tggatcgtgt tggatatttc agagaacatc ccgcttctga agctgctgca     1020
gctccctcct cagggatcac actgccgtca cccactctgc actggggcgt ttcctactgc     1080
gcctcgtgct ggcggacgca gctgggtgca gaagctgtgg ggtcggagag gcgtttggag     1140
aaggctctgt gtgcagtgtg tgaaaattca ggtgctagaa gcctactggt agaaaaaccc     1200
aaaaggaaga gctatacctt taaccattct gtccaatttc gggagccttg tcagtgtgtc     1260
agtttttctt ccccgaaagac actccttccc caagtaattg taggaagata aaaaaactgt     1320
taccagataa caaacactga actcctattt gaccagaact ttttcctctc aaaaaaaaaa     1380
aaaaacaaaa aaaaaaaaaa aa                                          1402

```

<210> 21
 <211> 1221
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (1220)
 <223> n equals a,t,g, or c

```

<400> 21
ggatggcgtc ccgcaaggaa ggtaccggct ctactgccac ctcttcacag tccaccgccg      60
gcgcacaggg aaaggcaaaag gcaaaggcgg ctcgggagat tcagccgtga agcaagtgca     120
gatagatggc cttttccaca gcccttgga cgtgcttagg atgcatacca gggtagaagc     180
ttaggaagtg tgggcaccca ccccgaccca tggacctgtg tgctccctgc gttgagcctg     240
agatgccatg cttctctcta ctctctctac cactgtggga tcccttggtg attttggttt     300
tctgtagta aacttctgtg tttgggatct cgttgcttag acaaataacc accctcatcc     360
ctgccatctg ccagtgctt cacaccctcg ctacacacag tgtggtcctc ctgtcaggtg     420
tcagcatcac tggagaatat tagaaatgca gagtcttagg ccctatccca gatctaatag     480
cttcaaatga atcttcattt ttataagatc ccgagagacc ctgtgcccac tgaagtttga     540
gaaatgctcc caacttatta ctgccactcc acagctgtca cctcctcagc tccaccaaac     600
ttccggagtt attgtcatca ccatttcaca agtgaggaaa tgtccattga gattaagagg     660
cagagagtaa gagggactca gatcgagatc tgctgccttc caccaggggt ccctgaaata     720
ccttgccctg actttccctg tgggtaccag gcaaaaggac acttcgaaga gttcactga     780
gcaatcgacc tgcccaggcc tcttgagacr tgcaatttam cawwactgtg ggtcacctca     840
tccacacctc taacaaagtc acaaggggaa ctggagaaat tcaagatgcc agcaaagaaa     900
taactgccgg gctgaaagac ttcataattg ccagatggct gaagtgttca gagagtcgga     960
gctatggaga agggctaatt gatcttgaca tgggctaaga aacaacatya cacctgcggc     1020
aatgactgca agcttttcta gtcccaagag ggaaggsaa aattctgctc ccaacaatga     1080
gagagaaatg gtctccaccc ccagtcagtg gcaacacaga gaggaggtag aagcacaggc     1140
tgatctgctt gtgaattgtc accacactta actagtcagc tagcctgggg aaagttctga     1200
tttattctga gacgtagttt t
                                          1221

```

<210> 22
 <211> 1255
 <212> DNA
 <213> Homo sapiens

```

<400> 22
aactagtggg atcccccggg ctgcaggaat tccgatacat tgggtctttaa acccaaacc     60
cctttaaggc atgttggtct caaggggatt tgggattgaa gcrcagcttt ctgttgagga     120

```

aacctaggtg	tcacttttta	gaacattaaa	aagaaagttg	taggggtggc	ttttccattt	180
aaaggaatgt	gatgatccta	ggttctgatg	agtgataggt	ggctctcgtt	tataaccctt	240
cttcttttgg	cacctgtaag	ttccggtagt	ggccatctta	cattctcatg	tcctgctgga	300
agtgcctagt	tgcttgcaaa	gccagctaag	gcttattatt	tcaaaagaag	attattttaaa	360
acatgagtga	caggtagtca	gaagagaaca	aaaggacgca	agatactcta	tagccaagtc	420
agcttggagg	caggatgggt	tgctgagtga	agtcgccgct	cacttttggg	ttcttatgga	480
ctgtgagtta	gtcttccctc	tacacggagt	cacaggaagg	gtataaatgc	atgttcctga	540
ggtgccctcc	cccaaagaat	gtacctgcac	tcaaaccagg	atctgttttt	gctgttttaa	600
tcataaatag	actagttagt	agaagacttt	tgaagaacaa	agtaaaactt	ttttttttca	660
ttaaaagatg	tccagagga	aaggccctgt	gcagccagta	tattctaata	actgcctggr	720
ccatgtccta	atawgggtgg	ttttaagttg	ttgggccaaa	aatcctttta	agacatacga	780
aacatctgcc	aacttttttag	caacccca	gagcccgca	mamgctcgct	ttcttccccg	840
ccctgcccc	ttagtccccg	ctctggaags	cccaggcagt	ttaggtgtaa	atasgtatct	900
tttatggttt	ccaaatgaat	tatttgtgtg	agagtaatta	aatctgtaag	aaaacctgtt	960
gagattcttc	acwatgaatt	atgacttcta	caacatgtat	tttagcaaaa	acacgatgct	1020
ggcctccact	ggatagctca	gtatgctgat	tgccagtgat	agttctgtac	gcgttaccaa	1080
cagcgtcttt	attaacccctc	tcccacatcc	agtggaaatc	attgctaggc	ggatattgtt	1140
ggttggtgtg	tagcttttgc	ttatgatctt	atgtttcttt	taaagggtgt	tttgcatggt	1200
gaatattaaa	tttttttttt	ctgtgtmaaa	aaaaaaaaaa	aaaaaatctt	cggcg	1255

<210> 23
 <211> 1642
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (1369)
 <223> n equals a,t,g, or c

<400> 23						
tcgacccacg	cgtccggtgg	aggggaccct	gtgggttagca	gcagctatcg	cagcgtcgga	60
tgttcagagc	agcagaagcc	ggcgctcgctg	gatgttgtgt	tgcccgccac	catgagctac	120
acaggctttg	tccagggatc	tgaaaccact	ttgcagtcga	catactcgga	taccagcgct	180
cagccacact	gtgattatgg	atatggaact	tggaactctg	ggacaaatag	aggctacgag	240
ggctatggct	atggctatgg	ctatggccag	gataacacca	ccaactatgg	gtatggtagt	300
gccacttcac	actcttggga	aatgcctagc	tctgacacaa	atgcaaacac	tagtgccctg	360
ggtagcgcca	gtgccgattc	cgttttatcc	agaattaacc	agcgcttaga	tatgggtgccg	420
catttgagga	cagacatgat	gcaaggaggc	gtgtacggct	cagggtggaga	aaggtatgac	480
tcttatgagt	cctgcgactc	gagggccgct	ctgagttagc	gcgacctgta	ccggtcaggc	540
tatgactaca	gcgagcttga	ccctgagatg	gaaatggcct	atgagggcca	atacgatgcc	600
taccgcgacc	agttccgcat	gcgtggcaac	gacaccttcg	gtcccagggc	acagggctgg	660
gcccgggatg	cccggagcgg	ccggccaatg	gcctcaggct	atgggcgcat	gtgggaagac	720
cccattgggg	cccggggcca	gtgcatgtct	ggtgcctctc	ggctgcccct	cctcttctcc	780
cagaacatca	tccccgagta	cggcattgtc	cagggcatgc	gaggtggggg	cgccttcccc	840
ggcggctccc	gcttttggtt	cggttttggc	aatggcatga	agcagatgag	gcggacctgg	900
aagacctgga	ccacagccga	cttccgagtg	agtggaggca	gccttcccct	ctgggaagct	960
tagttcccac	tggggcggag	ctaagggccg	ggtgccatgc	accctgacac	ggcttcccc	1020
ccttatgacc	cagaccaaga	agaagaagag	aaagcagggc	ggcagtcctg	atgagccaga	1080
tagcaaaagg	accgcgacgg	actgctcgga	caacagcgac	tcagacaatg	atgagggcac	1140
cgagggggaa	gccacagagg	gccttgaagg	caccgaggct	gtggagaagg	gctccagagt	1200
ggttaagtgg	tctggctggg	cattttctgt	ctgcaggcag	ggcgagctgg	cctaaagagt	1260
ggtgctctcc	ctggggggcca	gaggcaatgc	tgtgtcagct	ctgcgtgcat	ggtgaccagc	1320
atggggaggc	agtgtggtct	ggcasaaagc	acaggctgac	tagctggang	cacagtttcc	1380

ttgttgga	atgggatgag	cccctgtgca	ccaggtagtg	ggaggatgtg	aggtttmtgg	1440
cacatagccg	ggccttggtca	tttaaagcca	ttgtaaataa	ctgggctact	ctggggccacc	1500
ctgcctatag	gatagccctg	ctctgtctat	ggagcaagca	gctgttttac	tatataccgt	1560
tgctctaata	aacttggttt	cttttaaaaa	aaaaaaaaaa	aaaaaaaaaa	ctcgaggggg	1620
ggccccgtac	ccaattcgcc	cg				1642

<210> 24
 <211> 932
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (929)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (932)
 <223> n equals a,t,g, or c

<400> 24						
ttccactcct	gctttttggt	accagctgtg	atgtttggca	agttaataaa	cctctcaaaa	60
cctcacctgt	aaaatatgga	taacagtaca	ataagggttc	akcaaatagt	agatgttgtc	120
aataatgctt	tgttttcttt	ggaacatgat	aatcttacta	gtggcttctt	cggcctattc	180
tggttggtgac	cttgcccttc	ctggaacttc	ggcggttatga	ctgtwcttaa	ctgctgaagr	240
atggctgrat	gtctggaaat	ggraaaaatct	gtscgtgtgga	tgaaatctta	ttaatagatg	300
tggragmcac	taattagamc	accacaactt	aaaagagtgt	ggatgaatgc	ttaatgtctc	360
tttaagtcat	ggagatgggt	ttctgggaaa	gaggtgagtg	tagtgggggt	atgatggcat	420
ctgactccct	gttaccact	tcctgcagct	agatacactg	tcagatcctt	tggcatccgg	480
agaaatgaaa	agattgctgt	ccactgcaca	gttcgagggg	ccaaggcaga	agaaatcttg	540
gagaaggggc	taaaggtgag	cctaattccc	taatggagtg	atattgatca	gcactccttt	600
agtaacacat	gtagataagt	tacatttaat	gttctgttct	ttggtgtctt	gatatttatt	660
tacttaagct	tctaaaaggc	tttttctaca	atcagcaggg	ttaaactggt	cttggtgggt	720
taaaagatgc	ttgaggctgg	gcacgggtgg	tcaacgcctg	taatcccaac	actttgggag	780
gccaaaggcg	ttggatcatt	tgtggccagg	agttcgagac	catcctggcc	aacatgggtga	840
aacaccatct	ctactaaaaa	aagataaaaa	ttagccgggc	ttggtggcgg	gstcgrgtag	900
tccaagcaag	agagttccag	ttacggtgng	cn			932

<210> 25
 <211> 1064
 <212> DNA
 <213> Homo sapiens

<400> 25						
ggcacgagga	agagtcagcc	ttcttctttt	cctggcctag	gtagtagagc	tcatatagaa	60
aaagtgtgag	aatatgggtg	caaaaactaca	ttattttattg	cttcactga	aatgtcaaga	120
ggcagcaggt	gaggtcatgag	gatgggcagt	tctcagaagt	tttcctgaac	ctacagggtt	180
atgttaattt	ttttatgtat	aatttgtctt	ccttgtttat	gatctcattt	ctagtctgcc	240
atgtaacccc	ttctcaaaact	ttaaaaggac	ctcccttgag	ctggagctaa	cgagaccatt	300
tcttgtctgc	ttacaatttt	aaaaaaaaag	ctattttgcaa	gtaatttttc	tcattatgat	360
gctgttatca	taaagtgtga	ttccagtagc	caggggtgtca	agggatggta	tatggacagt	420
gcaactttga	cttactttac	tctacttagt	caaattttaa	ctattttctg	gttcctttca	480

agctgaaaaa	ttgtcttcaa	atttacagca	ccatttcagt	ttgtcatctg	tttattcaca	480
ttacttttct	gacactggaa	ttccagaagt	gaccacctgt	cattcccga	gtgccataac	540
tgtggattat	atthttctact	ctgcagaaaa	ggaagatggt	gctgggcacc	caggagctga	600
agttgctttg	gttgggtggc	tgaaacttct	agctagactg	tcacctctta	cagaacaaga	660
cttatggact	gttaatggac	ttccaaacga	aaataactct	tcagatcatc	tgcttttatt	720
ggcaaagtct	agacttgagc	tctgactctc	tttgatcaca	tactaatttt	ctttccaatt	780
tgtattgttt	ttcaaagaat	gtaaagtctt	taagtgtatg	catgttggtt	atthttgcac	840
tgtggagatt	ctgaagcgg	tatgttagat	gctttgaaac	tccatatacag	aagaaataac	900
tttataacaa	ttttttttaa	taatgaaaaa	tattttcctg	acaagtgagc	tctaaattct	960
ctttattgta	aaagagatgt	aaagggttta	tattctaaat	cctagtaaaa	ttgacagtga	1020
tttttaataa	taatgcatct	tcctttgtct	gcttagtaaa	aaatttcatt	tcataatttt	1080
ggcaagctct	gtagtggatc	caaagtatct	ttgagtctct	gcaaactaca	agttgtttcc	1140
tttccagaag	gcttgatttc	attaggagac	ccctctattg	agttctaaat	agtttatctt	1200
agaaagcctt	gggtcattca	cagggtatcca	accagccatt	gtttagtctg	tttttgaagg	1260
ggtttgataa	tgctttttaa	gttgtagaca	atgcttaatc	catcttatta	ctgtcctgag	1320
ccatgtaata	tgctgcatc	gtgttgggga	aatgtttggg	aaatataagc	cagcataacg	1380
tgtaaaagctc	actctttcac	cctggaacag	acaagagggtg	ggcttaatag	aggcagagac	1440
tggggatata	cctttgtttc	cctagcattt	ttattttatt	atthtttatt	tattttattt	1500
tttgagatgg	agtttcactc	ttgttgccca	ggctgggggt	gcaatggcgc	aatctttgct	1560
cactgcaacc	cctgcctccc	gggtcgaagc	gattctctctg	cctcagccctc	tcgagttagct	1620
gggattacag	gcattgcgtca	ccactcccag	ctaattttgt	atthttcagta	gagacaggg	1680
ttccccatgt	tggttaggct	ggctctgaac	tcccaacctc	aggtgatccg	cccacctcag	1740
ctctcaaagt	gctgggatta	caggcgggag	ccactgcaca	tggcagcatt	ttttaaaaat	1800
tcagttttaa	gatctctggg	ttaggggaga	gatttttatt	tactgaacca	gttctataga	1860
aatthttttt	attaggaaat	ttgtctttgg	aacaaagtgg	cagctataaa	attattttttg	1920
ttaagcctca	gaaatatgag	gaagcctgta	aaactctagt	ggggagatat	taacttggag	1980
acctaattgt	ctgtaaatag	tcattttaa	tggttggttt	agtgggtttg	tttctaaaat	2040
gttttctttt	accgtgatgc	accagtatga	gatgggtgctg	acacttttcta	tgaagtgggt	2100
atgagcagg	ttataaaatc	ctctatacaa	gtgtttgaat	caatttttaa	acataaaaag	2160
tggaaaatttc	cttttttgta	gacagtagga	caagggaatta	tatgcatttt	tactaagtag	2220
taatttttaca	ctgaattgta	aatgttttta	cagtgaagtt	tattaataga	atgcttcacc	2280
ttaaattgga	aaacaataat	agtcttgga	taagtctttg	tactaaagca	tttgctataa	2340
ttattttttaa	aaaaacaaac	agatgaaaac	ctcagagaag	gcatgtggat	tataagattt	2400
gtctagtaaa	aatgttaatt	gaatatgttt	aaatatttta	tttctcattt	tgggggggtt	2460
ttattttttt	atthttttaga	tggtgtctca	cgctatcgcc	caggctggag	tgcatgtggc	2520
cgaactcggc	tcactgcaac	ctccgcctcc	tgggttcaag	caattctcct	tcctcagcct	2580
cccaagtagc	tgggattaca	ggcatgcacc	accatgcccc	gctaattttt	gtatttttag	2640
tagagatgga	atthttgcat	gttggtccagg	ctgggtctga	actcctgacc	tcagggggtga	2700
tccaccgcc	tcgggctccc	aaagtgtgt	gattacaggc	atgagccacc	atgcctggcc	2760
tttgggggga	ttttaattac	agtatttaatt	atagttctag	gatttcccac	atthttatagt	2820
agtagtttgc	aggatattat	gtgccctaatt	tagcagatat	agacattatc	aaattataat	2880
gatagtataa	ttatcccttt	ttaatatttg	ggaaagaaaa	atgaaaattc	attagttaat	2940
tactgctttg	tggtgtgtga	atthttttaac	aaataacatt	attacatata	ggtcattgtt	3000
aacaaacaaa	catccctgaa	aacctctgtg	aaaattttaat	tttttatatc	ctgattaata	3060
tattgtgact	ttaggcccat	ttttcatgtg	cttcactttg	atagagttaa	tccataaaat	3120
tgctctttac	tttagcttat	caaatagaat	attattttgt	ggactggagg	ccaaaaagtc	3180
aatgtgagct	tctcacaggt	ttttaaagct	ccactaaaag	taattatcca	cttgtcttta	3240
cttttggtga	ccagaatagt	tggttaactct	gccagagcct	gtacttacct	gccaaaaaca	3300
attaaatctg	gttaatgcct	gaaaccaa	ctctcagctc	caagtgttat	actatccaag	3360
ttttaaatgg	aaaggtaaac	tgtggagtaa	tgaatttttg	gttttactgt	accttttgct	3420
atcaagataa	tattcatgtt	tgaaatcttg	cttttatttg	gaatttagtt	actgtctgct	3480
tttaaccttt	gctttctcaa	agaaagtgtg	agatccagag	agttcaaggg	attggggaaa	3540
gagagggctc	aagtcatttg	cactttgtac	ctgtaagtta	ggtaataaac	tattatactc	3600
gtaaaaaaa	aaaaaaaaa					3620

<210> 28
 <211> 1896
 <212> DNA
 <213> Homo sapiens

<400> 28
 ggcacgagtg aagatggacc agaagtcctt gatgaggaag gaactcaaga agacctagag 60
 tacaagttga agggattaat tgacctaacct ctggataaga gtgcgaagac aaggcaagca 120
 gctcttgaag gtattaaaaa tgcactggct tcaaaaatgc tgtatgaatt tattctggaa 180
 aggagaatga ctttaactga tagcattgaa cgctgcctga aaaaaggtaa gaggatgag 240
 caacgtgcag ctgcagcgtt agcatctgtt ctttgtattc agctgggccc tgggaattgaa 300
 agtgaagaga ttttgaaaac tcttggaaca atcctaaaga aaatcatttg tgatgggtca 360
 gctagtatgc aggttaggca aacttggtgca acttgctttg gtgtttgctg ttttattgcc 420
 acagatgaca ttactgaact atactcaact ctggaatgtt tggaaaatat cttcactaaa 480
 tcctatctca aagagaaaga cactactgtt atttgcagca ctctaatac agtgcttcat 540
 atcagctctc ttcttgcag gacactactg ctgaccatat gcccaatcaa tgaagtgaag 600
 aaaaagcttg agatgcattt ccataagctt ccaagcctcc tctcttgtga tgatgtaaac 660
 atgagaatag ctgctgggtga atctttggca cttctctttg aattggccag aggaatagag 720
 agtgactttt tttatgaaga catggagtcc ttgacgcaga tgcttagggc cttggcaaca 780
 gatggaaata aacaccgggc caaagtggac aagagaaagc agcggtcagt tttcagagat 840
 gtcttgaggc agtggaggaa cgggattttc caacagaaac cattaaattt ggtcctgaac 900
 gcatgtatat tgattgctgg gtaaaaaaac acacctatga cacctttaag gaagtctctg 960
 gatcagggat gcagaccac ttgcagtcaa atgaattcct tcgaaatgtt tttgaacttg 1020
 gacccccagt gatgcttgat gctgcacgct taaacgatga agattctcgt tcgaaaggca 1080
 tttatataac tctgcagcct tcaaagctcg aaccaaagct agaagcaaatt gtcgagataa 1140
 gagagcagat gttggagaat tcttctagat tttcagaact tgaagactat tttctaattt 1200
 ctattttttt ttctatttca atgtatttaa actctagaca cagtttttat cttggattaa 1260
 cttagataac tttttagca gtggttataat tgcttataat ttaattgaca atactattga 1320
 aactgggtgag ttctgattat taaatattct ctgtaaatca gtaaacatgt ataaagtatt 1380
 tgtaatgttt ggtcataatt tatttatgaa gacagcaaaa gactgatttc atgatgggga 1440
 aaacaattag ccaaagttaa atttcttaca ctgtggttgt caagaatact gatttactat 1500
 aatgatatat acatgcaaga tatttaactt aatatcttag acaagagttc tgggtacaat 1560
 tttgggatct agttccctcg gaaaagctgc tgtattttta atttttaatg gaatgtagct 1620
 tttaaaatcc tgtcactggc atcaacaaaa ggaattatac catgagacct tatagctgta 1680
 cttaaaagcc attcagttca gctattggga gttcatgatg aattagcata tgccagaaag 1740
 gttgctaacc ttaacatctg agagcagtaa cactgatatt atctgctgta tgagactttg 1800
 tgcattttac tttgaaataa agattttttt cccactgaaa aaaaaaaaaa aaaaaaaaaa 1860
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaa 1896

<210> 29
 <211> 1361
 <212> DNA
 <213> Homo sapiens

<400> 29
 agcatttggg ggactttctga gcgggtgtcca ccttttagct gtgatgaacg gtgctgttgt 60
 gaacacttgt gtacaagttt cagtgtggac gtgtgtcttc gtttctctcg ggtatatacc 120
 taagggtgaa tcaactgggac tatggttaact gtgtttactc atttgagcag ccgccagcct 180
 gctttcccaa gtgtctgcac catctcacgt cccttcctgc agccttggga gttctgattt 240
 cgccacatct ttgccatcac ttgtttttct ctgacttttt gtttccagcc catgttgggt 300
 gtaaagcagt gtcttgtggt tttcatctgc attttcatga tgactgataa tgttgaacat 360
 ctttttttgt ctttcttggc gatctgtatt tcttcttttg agaaatgtct attccgattc 420
 ttgccacctt ttagtgtgat tattgtcttt ttgttgtga tttatgagtt ctttacatca 480

tctagataga	agtgttcct	cacatatatg	attcgcaaat	atcttctccc	attctgtgag	540
atgtcatttc	actttcttga	tgggtgcctt	tgaggcacia	aacttttaag	tttttacgaa	600
gtccaattta	ttttctttt	gtcacttggg	cttttgctgt	catatgtaag	aatcttttgc	660
caaatccaag	gttatgaaaa	agaaaacctt	caaaagtttt	tttatttttag	ctcttttatt	720
taagagctaa	aataaaaaatt	ttaatagctg	catgtggtgt	gaagtaggga	tcccagttcg	780
cccttttgcg	tgtatctgtt	cgttggccca	gcaccgtttg	tgaaggctg	ctctttcctc	840
actgaacagt	ctcgacattc	ttgtcaaaac	tgaagtgcct	gtaggttatg	agttttattcc	900
agaaccctca	gttccactgc	accgatgcgt	gtgtctgtcc	ttgcgccact	cccacactgt	960
cctgttacca	tcgcattgta	agtcttaaaa	ctgggaatgt	aagtcctttg	acttttttct	1020
aatccaatac	tatgaacaaa	atagatggca	gatgaaattg	aaaatttcct	agaaagggtc	1080
aaactactca	aactgattca	acaagaaata	gacaatgtga	atagatatat	aacaagtga	1140
gaggttgaat	tagtaatcca	aaaacaccca	taaagaaaag	ccaaggacca	gagagcttcc	1200
cagctaaatt	ctaccaagca	tttaagaat	taaaaccaat	tctcaciaac	tctttcaaaa	1260
gactaggaga	gtgggaaatg	cttctcaatt	catttttatga	gaccaatatt	actctgatac	1320
cataatcaga	caaagacatt	gccaaaaaaa	aaaaaaaaaa	a		1361

<210> 30

<211> 1323

<212> DNA

<213> Homo sapiens

<400> 30

aggcccagca	ccacgggcct	ttccccaggg	cagggggagg	accacctaag	gattcaaggg	60
cagctcctgt	tttcttggtt	ctgtgaacac	tcgagtctga	gccagcccct	caggaattgc	120
ctcaaaagag	aaaaacaaaa	aaaagtcctc	cttcccagg	cctgctactc	caaggtttgg	180
ctccatccct	tgcctttggg	tccgtccctat	ttcccactc	ctggtctctt	atctttgggg	240
ccaccagtgg	ggagtacccc	gggcccacat	ccctctaagg	cgctaagttg	aaggaggcct	300
tcccagagtg	actattgggtg	ccaaagtccc	agttcctgtt	ggacttgggg	taaaaacagg	360
agatgggtgag	tgggtgtaag	gcccaaatgc	ccagagaagt	taactcgaac	ccatgggacc	420
tgtcccagcc	tgtcagtccc	tgatgagtgt	aacttccttc	ccctgggggc	ctggccyttc	480
tctccaaccc	agtggccatg	ctttctcacc	cagccttggtg	cccggcctgc	atttctgtat	540
atattgctgt	gtattgtgtg	tatgtatgta	ttcctggaca	agtgtgttca	tctgcagccc	600
ttgcctgagg	ataaggttta	ggattgggta	aagatcagaa	taccagggcc	agctaaggca	660
acgactccct	ccccaaaccc	ttgggacctc	agccagtccc	aaggctgccc	tgacaatcag	720
gcaggctccc	caccgtgagg	ccaagcctcc	tctgccactg	ccagcatggc	ccaagggagg	780
cttggccttg	ggcttgccag	cctcagctct	gccctgacaa	gggtcttgta	tccagggcag	840
aggcctgagg	tgaccagggc	ttgctttgtg	gctgatgcca	gcaggcttgg	ttctagtggg	900
caccactggt	gggcaacctc	cataactggc	ccttagggcc	taccttccta	cacagctagg	960
ctataatggg	cctgagttag	agggtagctt	ccccagcccc	aagcacaggc	agaggggtgg	1020
agagcaattt	ttggttttat	ttttgtttct	gaagtgggtg	ctgtacctcc	agccccagg	1080
gggccttccc	tggccacact	tctctgcccc	accagggcat	cgccatccca	gcactttgct	1140
ccatgtcacc	cgtaagatgc	cctttgctga	atgtacctga	gtgtatgtat	ttaaaaggac	1200
tcacatgggc	atcagagaat	ttatggctct	gtatccaata	aaaaagatgg	tgaaactggw	1260
maaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaactc	1320
gag						1323

<210> 31

<211> 2083

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (1781)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (2057)

<223> n equals a,t,g, or c

<400> 31

ggtgtccttc	taatgccatc	tctgtagaaa	atgtgccctg	taagtcagtt	tcttgggaagc	60
tcttctgtgt	gttgtccctt	ctccagcagt	gggtcgattg	tcagggagcc	caggatggaa	120
gctaagtgc	ctggtcattg	gttgttcttt	cagtgccctt	cagacagccc	ttgccctggt	180
ggtcttgtgc	cctccctgtc	tgtttgggtg	ctctttttata	aattagtgt	gacttcagga	240
aatggctctg	gatttcaa	agctattcct	ggagacattc	taattctgtg	gtttaaacct	300
taaaacaaac	aaaccaaagt	aattccctgg	atattgggtg	ctactgggtg	gaagagcatg	360
gtgcggcgcc	tgttacttgg	atgagctttg	atcaaagaat	ggcatcaaat	gataacagac	420
attggaggta	taagtgatta	caaggagaat	catagatcta	aataaaaaatg	gaatgggtgt	480
taatacttta	attgatcgag	atgatacagc	aataattttta	ttcatatatac	agtacaatat	540
ttaactttta	aaggaagtga	tattcatctt	agcagaggtc	tcttagcacc	atatttgcaa	600
cattggatgt	tatccattga	gccttgtttt	ggggaggaaa	aaagacacca	actttcttga	660
gtaaatgtgt	ctctgaaggt	gttttacaca	ggaatacaaaa	tttgcttgaa	ctcaaaaggg	720
tcttgtttac	agtactttta	tcttgttttt	cacttcataa	gccctctgta	aactgaaata	780
cagagctaca	ggcaaaccctc	attttattgc	acttagcttt	attgctcttt	gaagatactc	840
tgkttttttt	tttttattca	aattacagat	ttgtggtaac	cctgcctcaa	acaagtctgt	900
tggtgcatg	tttccaataa	cagggtgctca	ctttttgtct	ctgtgtcaca	ttttagtcat	960
tatctcaata	tttcagactt	tttcattact	ggtatatctg	gtatgggtgac	ctgtgggtccc	1020
agatctttga	tgttactatk	gtcattgktt	tggggcccca	taamccatgc	ccatataacg	1080
tggcaamcct	tatcaataag	ttttgtgtgk	tccgattgct	ycatgaaacc	agctgttccc	1140
cctctctctc	cctcttctcg	ggcctcccta	ttgcctgaga	caccaaatat	tgagattatg	1200
ccagttaata	accctgcaat	gcctctaaat	gttcaaatga	aaggaagaat	ctcatgtctc	1260
tcactttaaa	taaaattcta	gaaaggatta	aacttggtga	gaaaggcatg	ttgaaagcca	1320
aaatacgctg	aaggctagac	ctcttgccac	aaacagccaa	gttgtgaatg	caatgaaaag	1380
ttcttgaagg	aaattaaaaag	tgttacttca	tagaacacat	gaataataag	gaaaacacct	1440
tatttctgat	atagagagag	ttttaatggg	ctggatagaa	gatcaaatca	gccacaacgt	1500
ttccttaagc	cgaagccaga	gcaaggctct	aactctcttc	aattctctga	aagctgagac	1560
aggtgaggga	gctgcagcca	aaaagctgga	agctagctga	gggtgggttta	tgagggttaa	1620
gaaaagaagc	tatctccgta	acataaaagt	actagacgaa	gcagcaagtg	ctgatgtaga	1680
agctgcagtg	agttatccag	aagatctagc	tgggataatt	gataaagcta	rtacactaa	1740
acaacacatt	ttcactgtaa	accaarkagc	catctyggaa	naaagatgct	gtctaggact	1800
ttcttagcta	gagagtacga	ggcatcgctt	tttcttaaa	cttgaaagga	caagctgacc	1860
cttggttagga	gctaattgcag	ctagtgcac	taaattgaag	ttagtgtctca	ttttcccttc	1920
tcaaaatcct	acggcccttc	agaattatcc	ttaaatctac	tctgcctgtg	ctctaggaat	1980
ggaactacaa	aggctagata	acagcacatc	tgctgacagc	atgatttact	gactatttta	2040
agcccactat	taagacntac	tgcttagagg	caaaaaaaaaa	aaa		2083

<210> 32

<211> 1738

<212> DNA

<213> Homo sapiens

<400> 32

ggcacgagga	aggcttaaat	taagtatgtt	gacaataccc	caaactactg	gatgatattt	60
gtaatgaatt	tcaccctata	tcagtagaga	aggggtgcac	agttcagcaa	aacaaaagag	120
aatacgcttt	gttaaacctc	tactttctga	atttttaggac	agtgaggaaa	tttaacattt	180

tcaatttttt	tttcttttct	tgactgaaaa	gaaagtcaag	ccagcaatat	gtttctgaga	240
gagcagtgat	gcatttcaca	acactgttaa	ctgtctgctt	ggctttttga	ggcttccaga	300
gttcagaatt	gtcttctctt	gaataggtca	gtgcattttt	ttccttcagt	ttttctcctt	360
aagcagcaaa	acagaccatt	taacttccaa	atattttacag	cttgcaaaaca	gataaaacttc	420
ccaaatctgt	ttttttttta	tgaaaaggaa	aacgatcagc	cacaataatc	tataatatga	480
tatatttgaa	tcaaagttat	tagatgccct	agggtctttt	catggcagat	tttatatatc	540
accaccatta	ataaatctgt	tatcagaatt	atgtctttct	ctctgctgat	agttattttt	600
agactaacat	attcatacct	ccttctgatg	aaaaacatta	aaatttgaat	aaggcatatt	660
agaaaacct	aaagctctgt	atttacacaa	aggagactca	taaatatcgg	tttttcaggg	720
tgaagcattg	tgtgttattc	cattttgtac	cacagggaaa	gcctagtcac	acatggggcc	780
tcattaaaag	aggatctaaa	gaaatattta	atgggtgaaa	tataaggctct	tattctgaat	840
atctaccttc	actttataat	aatagaaact	gaactgaaaa	gattcagtaa	gtgattttaga	900
acatccactc	attttaaaag	taatgtctag	gcctaggaaa	gtgacatcat	gttccaaatg	960
ttacaaatcc	agcgtttttc	cttgatgtct	ttataaatac	attgtttaca	gtttttattc	1020
tcctccatat	atgatgcccc	ttttctaaga	ttatttcgta	ggtatatcct	ttaatgagag	1080
aacctcataa	ataaacttcc	tgaatttgaa	aacgagttag	aggagacttc	aagtttgtgg	1140
actgagctaa	acatgtgtct	actctccctt	caaacatccc	atggaaatgg	cagtaaagac	1200
agaacaaaga	gaatacattt	ctgaatatac	tgaggaaacat	atactccaga	gagcagaaac	1260
aggaagaggg	gcctctgcta	aagctgaagg	agtcttcagg	aggtaaccca	gctgggctct	1320
gtgttctgg	tggcagatac	agagagtagc	caagggtcaa	agcagacaaa	cagaaagtga	1380
ggtgattggc	cacaggtttg	gaaatggctc	aggtttgctc	tttccaaccc	cttgcattgc	1440
actgcacatt	tacctccaga	agtaaagcag	gagaactctg	caaagggaac	caacaatctg	1500
catgggaggg	cattgcctcc	aagtgtgtgg	gctggcgacc	aagtgcagcc	ctctgcccta	1560
gttatgggag	ttggagaggg	acttactgg	ttgcatgttc	catgcaagag	ctaaagcaaa	1620
gcacatctcg	gagaagttag	ccaggaaagt	ggatcaatcc	atgccacaaa	cctatattaa	1680
ttaacctgat	aattaatatg	tatataaagt	gaaaaaaaag	aaaaaaaaaa	aaaaaaaa	1738

<210> 33

<211> 917

<212> DNA

<213> Homo sapiens

<400> 33

ggcagcaggg	tttccgggtac	aggatacagc	tgtgcttctc	ctactttgta	tttgccatat	60
acggatagat	acaaaacctc	atgactgatt	tttccctatt	tattttgagt	ggctctgtat	120
accattttca	tgtattactt	catttttctg	tttttcagtt	atgttctctg	ttttcgtata	180
tttttgaag	ctagttctaa	gtcatgtttt	gtaggaaata	agcaatctta	aatacatgca	240
taggggattt	ctttcttctg	aggatccctta	atttcttcta	tttttaagat	tcaaattgaa	300
tgattaatca	gtaacagttt	atgttttaaa	taaaagtctt	taaaatgtta	aatatcagcc	360
tttcatttct	gatatattgg	ctttgaagag	gaaacataat	gcaatagtaa	ttcataatag	420
tggagggttc	tttccctaca	tccttgaaa	ccaccagtct	tattcttcag	cctggctctt	480
gagctattgc	tgtattaatt	ttaaataggt	tgtgatagca	taagctgatg	gaagcctgca	540
gagatctcac	tttgaaatgg	tgatacatta	catgggaaaa	gattagagag	gtgttttata	600
ctgcacatcg	gtgaggccta	ataagaaagt	agaatagacg	taaaccattg	ttttcatcat	660
cctttaagac	agggatttca	gactcaagaa	ccaggcagct	acctatttaa	atgagtcaag	720
tggggcctaa	tgtaagacaa	tgtgatgggt	aaggattgat	tctgagggaa	caggagcagg	780
catgccctct	ctaacagagg	ccgtcagttc	tgtgtacct	gaaatgtggg	cattgttgc	840
ggcgattcca	cacttacatc	taaaatttcc	tgatttttta	agaatgacaa	taaccaat	900
aaaaaaaaaa	aaaaaaa					917

<210> 34

<211> 1772

<212> DNA

<213> Homo sapiens

<400> 34

aatttctaaaa	taccaatgta	tttttaggtt	gtagctaattg	ttgtattcac	tttcaattct	60
cagttgtcca	cactggtgat	ataagaggaa	caaatcagaa	tcattaaata	ctttgtaattg	120
ccatcataaa	ctcatatatt	catcctcaaa	ctcccttggt	taatgctaatt	tgggtggcctg	180
gaacttcact	gagatgcaaa	atcaagaact	gaagcctagt	tgctagataa	caaaaagcta	240
taaatgttta	tgtatgtgaa	ttttaaatta	gaataaccgt	cttaaaactcc	tacttgccat	300
ttctaaggca	aagcattcat	tttaattattg	tactttgcct	tttcattcag	ttagtggagt	360
aagtcatgaa	acccttagga	agaaaaacaa	gttatgactt	attcactaaa	attgatgcaa	420
gacagtgggt	tctagatgac	catggccatg	tggtcatcat	ataaaacctt	cagttctctc	480
tatggtgctt	ggctggagat	tgacatgtga	ggatgtgcca	atcatattaa	atggatttgg	540
tctatgtggg	tgatattgtg	cctgaatgta	actgtgatag	actgaaattt	gttcttagct	600
ctcaaatcc	actgaagaag	tcaagtgaag	tggggtaaaa	tagggagatt	agtgacaact	660
ttgtgcaaaa	ttttttaaaa	aatggaagca	ggtagccaat	attagaatga	taatttaagg	720
gtgtgggtga	attttagtta	gttgctacat	agttattgaa	cctcatatgc	tcagtgtctg	780
gggaatcaaa	catggaagag	gtatggctcc	tgcccctaatt	gagaacaagg	gggaaaaatc	840
cagatataat	ctaaatgcta	ggttatgtca	gggtatagga	acacagagaa	tgggggacct	900
gtaagaactg	gaagagtcag	agagggtccc	attgaagagg	tcaaacataa	ttccggaaag	960
aattaggtag	tgaggagatt	gtgccaggaa	aataagtggg	aaaggccaca	gttatgtttc	1020
ctttgaatgg	aagagagaca	aagctatcag	ctatagatca	ttgttttctt	aagacagcca	1080
aactggccct	ttgaaaccat	tcaaattacc	ccagtttagc	tccctacctt	ttagtctccg	1140
tgaggaagac	aagctgttgc	attatcatat	tcctctgtgc	tgagcagctc	aagactcagc	1200
cacaatatgc	aaattgcttt	aatgccatat	tacggcagtt	gatttagaca	tttgccagtg	1260
caccaaacca	tgagagattg	tcccgacctc	atgccacctg	gcagatgtgt	acccagagat	1320
ttttctgtag	ctccatgttt	cccataaagg	gcattggaaa	tgacacagat	aagatcttcc	1380
tttggaaacca	ggcacatttg	gccccttctc	agtgactgca	ctgtgggaac	tcttcttaag	1440
aaaatattga	aaacagctta	atgctttcat	atagtgcacc	acatttagtt	gaaaactact	1500
gctgcatagc	aaatattgtg	actcttcacg	tgtccacagg	agctcttctg	tgggttttaa	1560
gctatgaagt	gtattcacat	tgtggaagttt	taattatctt	tattgaaatt	aattgtgtaa	1620
aaatggtatg	tgctctatta	ggtattcagt	ttgtatgtga	attctatata	gaaagtgggt	1680
tttgttcttt	gagtttgttt	tatttcttga	agattacaat	aaatatctaa	gagactatat	1740
tcctgaaaaa	aaaaaaaaaa	aaactcgagg	gg			1772

<210> 35

<211> 799

<212> DNA

<213> Homo sapiens

<400> 35

tgctggtggt	gttctgcttg	ctcgcaggcc	tggtgctggc	cgtagtatca	cagtgtgag	60
cggcgcgcc	tacctgcaag	tgggcaggct	tttgtgcaag	agtgtgtggt	atgcaaggct	120
tggggcgctt	tggecgctgat	cacgccccgc	gttattactt	ctacttctct	gcaggcttgc	180
tcacctatgg	cgtcacagcc	tttatcgcat	taaactgtgat	catcaagaag	ggcaaacact	240
tcttggtatg	ctcgccggcg	ccggccactc	agatgtgtgt	gtacctgacg	gttggttatt	300
ggtagtgatc	gtactgttca	cgattccaag	ttccaagaaa	gcgcgttaca	ttctttcgat	360
tacgccggcg	atttcattat	tggccgcgta	tatcttcgtg	gatcgagtc	agcgctttgc	420
aagtaccgcg	gacaagttgc	tgaagttctg	cctcaactta	cctatggctg	gcctgggcat	480
gttggtactt	gcctttatct	atggcctgta	tgacgcgacg	ccgctgcggc	ctaactatct	540
tggcgcggtg	gccgggttga	tcagccttat	tgccattcgc	tcttgggtga	cgtcgcggtt	600
tcatgcacat	ccacaccggg	agttcgtact	gttggtgttt	ggtgtgccc	ccttcctggt	660
cctggatatg	tttttcttca	attcgattac	ttaccacctg	gaattggccg	atgagccgac	720
acccaagttt	ctgccttatt	ggttcttggt	ggcgcggtct	atcgagttt	taactttcgt	780
ttggcgaccc	acaaggcac					799

<210> 36
 <211> 1730
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (455)
 <223> n equals a,t,g, or c

<400> 36
 gggtcgaccc acgcgtccgg aaggatggaa ataggaccct tgagccgatt actccgtgat 60
 ggctcagact gcatgcaaag actaggatgg ggctcttgct ctggctcagt gttgggcata 120
 ctccccctca gaaggccccc gccaaagagc ttagattttg gcttgggaaa aacattaccc 180
 ctcttcagta accctgaagc tctgtatttg gtatttggga ttcaggtagg tcagctgctc 240
 atgttgmtcg gcccaagtgt gtaagaacaa acagtaatgc cagtcatttt cccactaaga 300
 tgttccagtg ggaagggggg ctggtatgaa aaagagaatt tttttttctc tgtgtaatga 360
 taactttgtt cacgtagtaa gaattcagtt cttactattg gtgtgaatag ggggtaaata 420
 ttatttttat ttaaaagcaa aattaaatac tttcntgaac ctcatccatg tttgcaagta 480
 gatgtctact gtggttgctt tttttcctca agagaatatt ttaataaact tgaagtaat 540
 tttgttacat ttttctgtct gcctgtgtac tatgtattaa aactcacatg ggggctttca 600
 tgataaaaag ataaactgtt aagcagttgg aaattttcag tgttcttcca gtggacacct 660
 gccttggggc aggagcttct ttgtagtcac tattgataga atgggggtcac acacattgtg 720
 ctctcgcat aagggcagct ccaaggtttg gcatgagact atgcatgtgt gtggacacgg 780
 aggtttctca gtgagaaaga gtcctaagac agtgaagtgg aacgargcct taaaaatcat 840
 ctatgcagct gacttccagt ttcaggttct caggctcctt ttggtatttt aggaagccca 900
 cattggacta gagagagact tcagaccaag atatcatgtt tatgttcttt tagctagaat 960
 tgtgttaagg caatgactat ctctacagc ttagaagttc tgaagtacat ggccaggagc 1020
 ggtggctcam acctgcaatc ccmgcacttt gggaggccga rgcggttga tcaccgagc 1080
 tcaggagtgt gagaccagcc tgggcaagat ggtgaaaccc tgtctctact aaaactgtga 1140
 aaaacacatt agccgagcat ggtgatgcac gcctgtaatc ccagctactt gagaggcttg 1200
 ggcacaagaa tcacttgagc ctgggaggca gaggttacag tgagccaaga tgggtgccagt 1260
 ggactccagc ctaggcaaca gagcaagact ctgtctcaaa aaaaaaaaaa aaagttctga 1320
 agtacagtta tatatgtatt tgattaatac aaaagtagag aagttaatca ctcttaaaa 1380
 aatgcagttt gggaggccag gcacaatggc ttatgcctgt aatcccagta ctttggaagg 1440
 ccgaggaaag gcggatcact tgagtccagg agtaggagac cagcctgagc aacatggtga 1500
 aaccccatct ctacaaaaaa tacaaaaatt agctgggtct ggtgacgtgt gcctgtggtc 1560
 ccagctactt gggagactga ggtaggaggg tcacgtgggc ccaggagatt gagacttcag 1620
 tgagccatga ttacaccact gcacttcagc ctgggcagca gagtaagacc ctgtctcaaa 1680
 aaaaaaaaaa aagggcggcc gctcttaaaa gatccaagct ttacgtttcc 1730

<210> 37
 <211> 1020
 <212> DNA
 <213> Homo sapiens

<400> 37
 ggtggaatg aaataaatgt gtacatgtaa agcacaagaa catggaaggt gcttactaaa 60
 tgttacttat ctttactttc tctgccttgg tctctcata cccactcctg attttaggtg 120
 attgggtgga aatggccatt gaacatcata ctctactaac aaagaccatt tgagagttag 180
 attaatctct tccccgttcc aacaacagga agaagcccca caaatcaagt atttcccttg 240
 ttctatacct tgtcattttg ttgctactcc caccagccaa agagggagga aagtttcttg 300

gtataattaa	aatgttatag	gctgggtcgg	gggggctcat	gcctgtcatc	ccagcacttt	360
gggaggctaa	gacgggagga	tcgcttgagg	ctaggagtgc	aagaccaggc	tgggcaacat	420
agtgaagccc	atctctacaa	aaaaaaaaag	atagccaggc	atgatggcat	ccatctgtag	480
tcccagctac	ttgggaggct	gaggcaggag	gatcacttga	gcccaggggt	ttcaggctgc	540
aggaagctat	gatcatgcca	ctacactcca	gcctgggcaa	tagagcaaga	ctgtctctct	600
aaaaacccaa	aattgttata	gaatatagag	ttgaataact	tttctggaat	gagaaagctc	660
tcattttaga	tatccattca	ttcattcatt	caatagtgtg	ctggatgcca	rgaatttaat	720
ggtgargaaa	atagacatga	tctctgcctt	ctgargctca	aratcctccc	tctattttta	780
aaaatcaggt	ttattgaagt	ataattgatg	tacagtaaaa	wttactgttt	ttagtggaaa	840
cttctgtggg	ttttggcaaa	tgtgtaacca	acacaattaa	gatcyagaac	atcctgtctc	900
tcccctccca	attttcttgt	gctcctttgg	agtcaacaac	tctcccccaa	ccccatggcc	960
tccatgactt	tttcaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aagggcggcc	1020

<210> 38

<211> 957

<212> DNA

<213> Homo sapiens

<400> 38

cccacgcgtc	cggagactga	gtggcttaaa	caaaagacat	ttattttctc	acagtccttg	60
aggctggaag	tccaagacca	agggtgtcgc	agggttggtt	tcttctaagg	cctctctcct	120
tggcttgagg	atggccctgg	ttggctctct	gagtgcagg	gtccctggtg	tctctttgtg	180
tgtccagatt	ttctcttata	aggacactgg	tgagattgga	tgaggggccc	cctgacggcc	240
tcattctaat	gtcatcactt	ctcttatctc	catatccatt	cacatacggg	ccattgcgag	300
ctaccgggga	ttagggggtc	aacacataaa	ttgggggtgg	gggggtgcag	ctcagcccat	360
aaacatgccc	cctctggctc	gctctcccag	ggcatccatc	gtagcactta	gaaaaatgat	420
cacttctttt	ttggctttgt	ggtggcgtgt	gtgtgtgtgt	atttgtatat	acacacatat	480
atatataaat	acacagacac	acatatatac	aaacacacgc	attcattttc	gtccacagtt	540
cctggctcat	aactcccaca	gcccttggtt	cactcttttg	ttacaacatt	gggtgtgtca	600
ggcctcagga	gacaatcact	ctaacctcct	gcccttctct	ttacctgccc	aagacaggac	660
tctaactctc	cctacctttc	tgatgggtgg	tcataaaaact	cattccagag	acggtcccac	720
cccatatcct	gctagaagga	atgctgctgt	catgaagctt	ccataaaaaa	caaggggact	780
ggattcagag	agcttccaga	taactgaaca	tacagagggt	ctagaagggt	ggtgcgcccc	840
gggagggcac	aggaagctcc	atgcccttcc	ttcatacctc	accctatgca	tctctttatc	900
tgtatctttt	ataatatcct	ttataataaa	ccagtaaatg	taaaaaaaaa	aaaaaaa	957

<210> 39

<211> 1034

<212> DNA

<213> Homo sapiens

<400> 39

cccacgcgtc	cgttttatct	aattttatact	taagtttgaa	tagccacaca	tggctagtgg	60
ctatcatact	ggtcaacatg	ggctatctaa	ttctgatgta	cagtctctgc	atcatcatc	120
cattttctaa	agtgggtttc	tcccacagca	gttggttaaaa	caaggatttg	gatacaaaaga	180
gattatttaa	gaggtgatcc	tagacagcag	gagagggatt	gaggagggtga	gatggagaag	240
gtgaatacta	gtaaaagact	acattgatat	gctgtaggca	cttgagggtt	ctgtttgttt	300
gtttgtttgt	ttttttggaa	accttcgaga	gacttatgga	aaacactgta	taattgtttc	360
attgaggagc	acggaactga	tcattgtttat	tcaccaactc	tcattgccttt	ttggttgaag	420
gatgtccta	ggggcattca	cacctgctct	gaactccttc	ttcatctcca	gcctgcctcc	480
cttatgggcc	atttcatgcc	actgtacctc	gataaagcct	tagggccaag	agattcaggt	540
gcttgaggct	gggtctatcc	ttctgagctg	caaataactt	ccagggtgga	tcaagggaat	600
gtggggtcag	gatagattat	attgcctctc	atttacaacc	aaccatatta	gctcatatta	660

gtgtttttatc	tggaatagga	ggagaagcat	ttgagactgt	tagaaaaaag	gttctgctat	720
ttaaaaaaaaa	agttttgaaa	ctgtagctct	aaactacatt	cctacctgtt	ttcctgacca	780
cctcaattac	taacttgtat	cgtgttccat	attatttcta	gcacagcatg	gattctctaa	840
cattcttgca	gtctgggccc	tgcaaggtag	ctgtaaaaat	gctgccctgg	gactttctca	900
gtggagacat	aagaagactc	aagcattcca	gaaagtgttt	gtttggttgg	cttttattga	960
tgtatataaa	aagattagag	acagcaagag	aagaaaatca	gagagaacct	atacttggtt	1020
aaaaaaaaaa	aaaa					1034

<210> 40

<211> 1761

<212> DNA

<213> Homo sapiens

<400> 40

ccacgcgtcc	gaagagacct	tgggtaataa	gcttgccact	gagtggctgc	tttatgtcct	60
caagggattg	taccatatca	agcactcagc	attggtgtct	gttgctaata	gtcaggcatt	120
cagtagtggg	ggtagctaga	tcagccttgg	agagagagag	cccatcattt	caggccatca	180
cgactaatcc	attcatttat	gggccctttg	taaggattga	gatggctgag	gacaggggct	240
gacaggcatt	cattagacca	gtcactcctg	ccatatgctt	attcagtgtc	tcttctgcca	300
cacgtgctct	ctgggtgggct	ctagtgtgag	acacaaagat	caacacatta	tgtgcccatt	360
cttatagctc	cagccacatg	cctcttcttc	agacatgctt	ggtttcaate	ctctagtgtt	420
gttcccttga	ggcccttgac	caagcaacca	agccattctc	caccacctag	aagtctgtgt	480
atattctttac	ttttggccgc	ttctctccag	acacaaagca	gatgaccact	ggacttgaat	540
tggcaccag	agttattttg	ggtgtgtctt	tagtgcagca	ccagtccatc	tttttagctc	600
acgccagcat	atcatgctag	cctaatacctt	ataaagccct	tttctgctc	cttttctatt	660
ctgtcaactg	tctgtggaga	aatccccaag	gggccatagg	tattatgtct	ggaattgggt	720
ctctccgagg	ggttcttggg	ctcgttgact	tcaagaatga	agccatggac	cctcgcagtg	780
agtgttgtag	ttcttaagaa	tgggtgtgtc	ggagtttgtg	ttttcagatg	ttcagatgtg	840
tctggaggag	tttcttcctt	ctgggtgggct	cgtggtctcg	ctgacttcag	gagtgaagcc	900
acagaccttc	acagttagca	ttacagctct	taaagtggtg	gcgtccagag	ttgtttgttc	960
ctcccggtgg	gtttgtgggc	tcactggctt	caggagtga	gctgcagacc	tttgcagtga	1020
gtgttacagc	tcataaagg	agtgtggacc	caaagagtga	gcagcagcaa	gatttattgt	1080
gaagagtgat	agaaaaaagc	ttccacagcg	tgggaagggga	cctgagaagg	ttgccgccac	1140
tggctcgggt	ggccagcttt	tattccctta	tttggccccg	ccctcatcct	cgtgattggg	1200
ccattttaca	gggtgctgat	tgccccattt	tacagagtgc	tgattgggtc	gtttttacag	1260
agtgttgatt	ggtgcgttta	caaaccttta	gttagacaca	gagtgttaat	tggtgcattt	1320
ttacagagtg	ctgattgggtg	cattttacaaa	cctttggcta	gacacagagc	gcttattggg	1380
gcattttaca	tccttttagct	aggcagaaaa	gttctccaag	tccccacca	accagaaggt	1440
ccagctggct	tcacttctca	atcctccttc	taaacaggac	accacaagtg	ttgttgggaa	1500
ttggccgatg	accgtcttag	ctatttcctg	ctggataggg	gcaaagaagg	ggccctgcag	1560
ttgtagtgtc	ctccagaggg	gaactcttta	ggccagtga	agggccagca	ggttgggtctg	1620
gggtcctcag	tagaagttgt	tagtttagct	catttggggg	tccatttgta	agaccatctg	1680
tagcttgatg	gcctcaattc	tagaggaaac	aaatttgaca	aggagattaa	aaatacaggg	1740
tccaaaggca	aaaaaaaaaa	a				1761

<210> 41

<211> 616

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (610)

<223> n equals a,t,g, or c

<400> 41

ccacgcgtcc	gaaaattagt	taggcgtggt	ggcacacacc	tgtaatccca	gctacttcgg	60
aggetgaggc	acaagaatcg	cttgagcccg	ggaggtgcag	gttgacgtga	gccaagatca	120
caccactgca	ctccagcctg	ggcagcagag	tgagaccgtg	tcccaaaaaa	gagaaggaga	180
aacagagatc	atgtggaaaa	agttattttt	tatttattta	cttagttttc	agtttggttt	240
gagactcgtg	ttttaaacca	gagggcatgg	ttactgaggg	ataacatcaa	tagaactcct	300
ataattgagg	ggataattat	caaggtatta	gatgattcac	tggtatttac	aaagaacaca	360
gaaattatga	aacctgggtc	tgtaacttat	agtttttcat	attattttta	taccatggat	420
aactcttcta	tgtgtattca	taggtgtaag	attactggca	gtgtcatatg	aaacaacata	480
ttacattttt	taagcctgga	aagcatctag	tatggctgtg	cacgtagtga	tgacattgac	540
tttttttact	taaagaagag	ctaccacttc	aaataagaaa	aaaaaaaaaa	aaaaaaaaaa	600
aaaaaaaaan	aaaaaa					616

<210> 42

<211> 573

<212> DNA

<213> Homo sapiens

<400> 42

gctccaaact	gttttccata	gtgggtgtac	tgatttagat	ttccaccaag	agatatgtat	60
gagagtcccc	ttttctccac	attcttgcca	gcatttggtta	ttgcctgtct	tttgataaaa	120
agccatttga	actagggtga	gatgacatct	tattgtagtt	ttgatttgca	tttctctgat	180
gataattatg	ttgagcactg	atatggtttg	gatctgtgtc	cccaccaa	ctcatgttga	240
attgtagctc	ccagtgttgg	aggtggggtc	tgctggaggg	tggttggttt	tctcatgaat	300
agtttcatac	tgctctcttg	gtgctgttct	tgtgatagtg	agttctcatg	agatttggtt	360
atthaaagc	gtacagcacc	tccccgctca	ctctctcttg	ctcctgcttc	cgccgtctaa	420
gatgccttac	tccctctttg	ctttcttgcca	tgattggcag	tttcttgagg	cctctccaga	480
agcagaagct	gctgtgcttc	ctgtacagcc	tacaaaactg	taagccaatt	aaaactcttt	540
tctttataaa	aaaaaaaaaa	aaagggcggc	cgc			573

<210> 43

<211> 1384

<212> DNA

<213> Homo sapiens

<400> 43

cccacgcgtc	cggaggaatt	gctattttaag	tggaaggtgc	ttgaacagcc	aatgagcac	60
ctgtcatgtt	tatttttact	ttgaaaagta	gaaacatgac	agtatagttt	ccactgccta	120
tccaatatga	acttctggca	ttcttggtat	cttctcagga	actgcactgt	ttgcttcttg	180
tgttcaattt	tttttttttt	tccagggatg	aggatgtatc	tctcctcttg	aaactttcac	240
tctaaaatat	gcccccaaat	actgacattt	tcaagcttgt	gctttttttac	ttgatatctg	300
atthttttta	aatttttatt	taataggag	aaggtcatga	cttatggaat	ttactctaag	360
gatagggatc	tgagcagggt	cctgagtact	tgaaaggctt	tatttacctc	ttacactttt	420
ttgttacatc	cagggttttg	ctcattttct	tatcatttgt	gtcttgaaact	ttacctgaaa	480
tgcatgtctt	gagtgcttac	taagggtccag	gcattgtgct	ggcagatttg	atatattttt	540
gtttattctt	tattacagga	ctattacata	gtagactctc	attctgacac	tcactttttt	600
tttccagaaa	aaaaaattaa	tgagatttga	ttgctaattg	gtacagggct	gcttttaagg	660
ctatgaaaat	gctctaaaaat	tagtttggtg	tgatggctgc	acatctctga	acgtactaaa	720
atacattgaa	ttatatactt	tcttttttct	tcgttttgca	gcttttgctc	aaaggagaa	780
tagatattta	aaatgagtga	attgtatggt	acatgaatta	gatctcaata	aaactatttt	840
aaaaaaagaa	actaaagctt	agagaagtat	aatatcttgt	tcaatatgac	actacctggt	900

aaagcttttg	tccaaatttt	tctgagacca	aagctattct	ttcttccact	gacgcatgct	960
atctctacta	attatatagc	cacggtatcc	tttttcttta	aaaatgtagg	aaaaaatggc	1020
tgggtgcggt	ggctcacgcc	tgtaatccca	gcactttggg	aggccaaggc	ggacggatct	1080
cttgagggtca	ggagtttgag	accagcctgg	ccaacatggg	gataccccac	ctctactaaa	1140
aaaaaaaaaa	aaaaaaaaaa	aattagccag	gtgtgggtgg	gtgcacctgt	aatcccagct	1200
actcgggagg	ctgaggcatg	ataatcactt	gaacctggga	ggcagagggt	gcagcgagtc	1260
aagattgcac	cactgcactc	cagcctgggc	aacagagtga	gactccgtct	caaaaaaaaaa	1320
aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	1380
aaaa						1384

<210> 44

<211> 706

<212> DNA

<213> Homo sapiens

<400> 44

gccacgcgt	ccgtgcagag	agactgggtg	ctccgagctc	cactcagggtg	aaagaatttg	60
cggcaattgt	tgacgtgaaa	gaagaatctc	attacatctt	ggatccaaag	caagcactga	120
tgaagctcac	cctaggtact	gcaggcagtt	tatttcccca	agcattgtac	attttgcttg	180
acttcatatg	ggtaaatttt	attgatggct	ctcattacat	ttagttgtgg	ggtgatgtca	240
ccttcgtagc	tcattttaag	tctttagacc	accatcagtc	ataattttca	aagaagctaa	300
ttttgtctat	taaatggaac	agaaacttcc	tactctgaa	ttttggataa	gtttgtcatt	360
tagcccatgg	tgggggtaag	agtccactt	tctaaattgg	tgatttctgt	cacatgtcta	420
aggtagaacc	agctgcaggc	agtggggact	tggggactag	aacaggcagg	gaggtggaga	480
gctattctgg	tgggatgtcc	taggggctga	tgaagtggag	ccttgacagc	agctttgttc	540
taaaggagct	taaagagaaa	gcagtggccg	ggcgagtggt	ctcacgcctg	taatcccagc	600
actttgggag	gccgaggcgg	gtggatcacg	aggtcaggag	atcgagacta	tcctggctaa	660
tgtggtgaaa	ccccgtctct	actaaaaata	caaaaaaaaaa	aaaaaa		706

<210> 45

<211> 419

<212> DNA

<213> Homo sapiens

<400> 45

ccacgcgtcc	ggaagaggag	gtttgtgttg	ccttcagatg	actgttgatt	taattgacag	60
gcagcttttc	ttcatttgtt	gctttttgct	tttgctcaca	aaagtgcatt	cactatttcta	120
tagtatgttg	aaacttttaa	atggaaacgg	cttttcatta	acaaaggaag	cattttcttc	180
cccccttcca	tgccttagct	ttctccgcta	agtcttggct	tcttcagcag	ctgtacctcc	240
accagaaatg	acaaaagggt	caatttgtgt	aggagatggc	gatgaagata	tgccagtggg	300
gttttgtgtg	tgggtgctg	gggactgttt	tccttctgtg	tttggttttg	ttttattttt	360
gttattccag	aaagttgaga	gcacatctaa	aaacaaaaaa	aaaaaaaaaa	aaaaaaaaaa	419

<210> 46

<211> 1025

<212> DNA

<213> Homo sapiens

<400> 46

gccaagcttc	aaacatagat	ctcctgactc	cattcatatg	accctataaa	ctgtctcaaa	60
acaaaaagat	aaattaatat	aaatatattat	tgaatatgtc	ttttagagaga	aagcataata	120
agcataaagg	gcaatgcgtt	aacctttatc	acaagcaacc	ctattggaat	gtgtcaactt	180

atcagaatga atcaggccag aatatcaagt ataaatgaag cctgtagtta actgaaagtt	240
gcatatcaat caggcactcc agtttctctc ctcaaaactct gaatattcaa tgaataagat	300
aaagaaatgg ctaatttgat ttacaccttc atttttttga cctaattcta aggtgactac	360
tcactcctca agatttaact aatgttgctt tatttttctc cctctgggga gacagaagag	420
atgattggga aacacatgtt tgaagtttgt aagttctgct gctttcaacc ccacagatgt	480
ctcttactgc ccacttgss cctkgtgatt aagcaactag atttgagacc agtcaggctt	540
ttgtttagac attttaactt tttcttgctt tccttgcaaa ctctcagcc ttcagactgg	600
ttggaaagta aatgtacaat cttacataaa ttttcaggta atagcatttc agctttttcc	660
ccargatttt ttgcttgga ggagacagat tagactggat tcggagtctt gattttgcaa	720
aggtaacaaa agacatgttt ttttataaga cttttcatca taagtttatt ttattcaaca	780
gaagcaaat ctaataataat ggaaaaata aagatctgtg ataaatctga tctgtgtkga	840
taaacacaat tagaaagatt taaagattaa gtattgaaac aaactaccaa aatattttta	900
tactgatttg taaaaatttc agtacatttt tcttctttgc ttaattctac tgggtcctgt	960
ttttcatcaa aacattctat catgttagta tacaatagcc aaaaaaaaaa aaaaagggcg	1020
gccgc	1025

<210> 47

<211> 783

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (14)

<223> n equals a,t,g, or c

<400> 47

gactcactat aggnaaactg gtacgctgca ggtaccggtc cggaattccc gggtcgaacc	60
accgtccgcg gacgcgtggg ttttaagttg tcgtcttgcc tctgtgctct tgaagttttg	120
agccctttgc atttggcagt gtccaggcat tcaggcctgg agcccgtgta gtgccagtgc	180
ctccctccac gctcttgccc tggttgacct cacctaacc ctccaaaaag cagatgggtca	240
gactctcttc ccttcaaact cttctctgcc ctgactcaca cctgggccat ttcattccagt	300
gagactgagg gaggagcgag ggagtccatg tttccctccc atgacgccgg gacaggaaagc	360
tagactcagt ctcttccata tggccaggaa ggggagtacc tgactgcca tcttggtttt	420
gggagagaga aaaaccagtg ctccagtctgg gaaatgaggt tttgggggat tgtgataaaa	480
tagaggacag gactctgcag gtcaaggaca gaggtgcac ctgagggcgc actgcagtca	540
gggccaagtg gctcactctt tgcagcttta gcagcacctt ggaatatagtt gctgtccgt	600
aaacgtgttg attgacagag gtgcaggtaa aaacctcaga acagttgggc ttaaggatgc	660
tacaaaaaaa gctctgggat gggatttaat tttttttttt ttttaactgtt gttatcagcc	720
tggccaacac ggtgaaatcc catctctact aaaaatacaa aaaaaaaaaa aaagggcggc	780
cgc	783

<210> 48

<211> 909

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (909)

<223> n equals a,t,g, or c

<400> 48

```

ggctgggggtt tgccattttg tgtctttttt agtgtctctc caaattaaat tttcaagatc      60
aaatatggga ctcatgtgcc tgtttatgga gctttttttc ttttgggtga ctattctttt      120
gtttcctaca gtcttcaaca actgttagtg cttacaatgg atacatcatg catcttaatt      180
gataggatgg atatgataat accttttttg gtccaaggtc ccgctcatta aaaaaaatag      240
tttataaagc tgaaaagttt ttatttctat tttttgtaaa atgattttca tgataggatt      300
ttatataaag gggaagggtt ttttgtatca tttttataac atttttgaaa tgagtactta      360
ttctctttca tcatctattt tagactcaca gttttatgag taatgcagta aaggtcatgt      420
ggcacattag taaaatatgt tctgaacaca gaaactattc tccttatcac aaattaaatt      480
ttatgttaag ttgaagagc actggcctgg ggtatacttt gctgtgaaaa gatcattttg      540
gtcacttaaa ttacaataga aatatttgtg ttaagaaaaa taagtaaaaa ttaggctggg      600
cacagtggct cgcacctgta attccaggac tttgggaggc ctaggcaggt ggatcacctg      660
aggtcaggag ttcgagaccg gcccgaccaa cacggtgaaa ccctgtctgt actaaaaata      720
caaaaattat ccaggcgtag tggcaggagc ctgtaatccc agctccttgg gaggctgagg      780
caggagaatc gcttgaaccc gggaggcgga ggttgcatg agccgagact gtgccactgt      840
accccgacct gggtgacaag agtgaaactc tgtctcaaaa aaaaaaaaaa aaaaaaaaag      900
ggcggccgn                                     909

```

```

<210> 49
<211> 766
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> SITE
<222> (758)
<223> n equals a,t,g, or c

```

```

<220>
<221> SITE
<222> (759)
<223> n equals a,t,g, or c

```

```

<220>
<221> SITE
<222> (760)
<223> n equals a,t,g, or c

```

```

<220>
<221> SITE
<222> (761)
<223> n equals a,t,g, or c

```

```

<220>
<221> SITE
<222> (762)
<223> n equals a,t,g, or c

```

```

<400> 49
ccacgcgtcc ggtctcagca acaagagcaa aactccatct caaaaaaaaa agaaaaaaga      60
tggcttgccc tggcattggc gtttgtcctt cttgtacaca caaaagcatc atgctttgta      120
cactgttcgt ttttgtttcg ttcactgtct tcacggtaaa gatgaactaa gagccactga      180
gaaggctgtt tatgggaacc agaaggttgt ttgttccct taagtctcgt cttcctggca      240
cctgggtgcc ttgcagccca tggggaatgt gatggtcacg ttttctcgtc tctcctgcct      300
catcccatca gccagttcac tgctgtgcct gaactcctgt acagggtgtc tgggtgcatgt      360

```

ccacataaca	aaaagatggt	actgatcatt	gagtttatct	ccaaaaattt	ttttcaattt	420
gattgtagaa	aatacatttt	taggcggggc	atgatggctc	atacctgtaa	ttccggcatt	480
tggggggccg	aggcggggcg	atcacctgag	gtcaggagtt	ccttacagcc	tggccggcgt	540
ggtgaagccc	cgtctctgct	aaagatacag	agactagcca	agcgtgggtg	catacgctg	600
taattccagc	tactcgggag	gctgaggcag	gagaatcgct	tgggcctggg	aggcggaggt	660
tgcagtgagc	agagattgca	tcattgcact	gcagcctggg	cgataagagt	gaaactccgt	720
ctctaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaann	nnaaaa		766

<210> 50
 <211> 712
 <212> DNA
 <213> Homo sapiens

<400> 50	
ccacgcgtcc	gggcctccac cacaagaaat tggctccaaa cctattccca ttacatcatc 60
accactaat	gtcattacac cagaatgaaa tctgctttga ggcttgtatt gttgttttcc 120
ttccatttct	ttccctcat catttccttt agaaggaaaa gggaaggaaa aaaaaagaaa 180
aaaatcagag	acctgtaaac atttaagagt gaattggaca tccccacagg ctaaaccctag 240
agttgtcagc	tttcattcca ttctgctgtg atagagtgtt ctgcccagta tgtatccaca 300
tggagcctgg	aaagtaacaa gtctggtcat gtgacaactg gctttggtgc tttgctttca 360
gccctcaga	attagttcct aatggctcac atagctccct gccagcatcc cgtcactggc 420
acgttaaacc	ttttcaaata attcctcaga atagagattc ttaggagagc actaaatttc 480
tccattcttt	gtgtttaatt aaactgttct tgcattttta tatttgccag ataagcgctt 540
aagacattac	gatacagggt gagtatcctt aatctgaaaa tctgaaatcc gaaacttttt 600
gacctgaca	tgacgctcaa aggaaatgct catcttcgat tttggatttc agatttgagg 660
tgctcaacca	gtaagtataa tggcaaactt attccaaaaa aaaaaaaaaa aa 712

<210> 51
 <211> 680
 <212> DNA
 <213> Homo sapiens

<400> 51	
cctactttgc	cagtacgac tgttcctctc tctctctcgc tccaaaagac taatctgcac 60
actctgttac	agcacttgct taattgtgct gtagcgttta ttacatggt tgtttctcct 120
ctagccagtg	agcacctctc aagcagaatt atcttccttg tttctctggc accctaaata 180
tttgttgaat	taatagtcct tctctctctt tgtaattttg cttctctgtaa tagaagctta 240
attttaagta	tagttatata agtaatcaaa atgaatcaca cactgagaaa tcaatgtgga 300
tgccctttta	gggttctggt atttttttta ttgccattga gtaaaataag atactctgtg 360
ataaagtata	ttagcattaa agtggtcaaa tctgatcttt attagtaggc ctcaagtgaa 420
tccttgctga	catttaagggt ttatgacatt tccttcacgt tcgttcttga ctggaaggca 480
taaatggctg	acagtaaaga gcaattaata attttccaag taaaacattt tcaggagacat 540
ctgcctttat	tgctccccag atgagagtac agcctgtttc ttatgtgttc caaagatgat 600
ttccctatca	gcttttttgg cagttaacca aaaaaaaaaa aaaaaaaaaa agaaaagaaa 660
aaaaaaaaaa	aagggcggcc 680

<210> 52
 <211> 1004
 <212> DNA
 <213> Homo sapiens

<400> 52

ccacgcgtcc	gcccacgcgt	ccgcagtggt	gtacagcaga	tctctagagc	ttattcatca	60
gttaaggaag	ataagctcag	gaaaatgtat	ttttaatgtg	aaatgtcaaa	aggacttagg	120
aaagaaagtc	atgttttctt	cttgctgttc	agtaatctag	tcattaccaa	acaatgttat	180
caatgcataa	catataaaca	tttcataatt	cattttcatt	tctagaggaa	actagtgagt	240
aaaaacaata	tcttcaggta	gcagtgaag	gaagcagaga	tcatggccgt	ttggatttga	300
gtgaattctt	tcaacacaaa	tcaatgattt	tatagacata	tataggtttt	tgctcaggggt	360
ccttagcttt	tgcatctcaa	aatctatttt	tgctaaggaa	ctgtataata	gctcatctgt	420
cttaccctcc	ttgtctcaga	gcccaggctt	ctgtgccttt	tccaatctga	aaagcctcct	480
ccttgtctct	gtccttttga	gcctcatgta	actttttaaa	ggccacttt	aaattccacc	540
actctataaa	aagcttctcc	aactattcta	cctttcaatg	gctctcagtt	tcctccataa	600
tactttggac	cttttagctg	atcctgactc	tcttgacta	cagtgtaaag	gttctgtatt	660
tatgggctgc	atactggcat	gatgcttctc	agttatgtat	gatatcatag	tcgctttttc	720
aggacaaaac	tgtgtcacag	tgctacagca	aaatctataa	cacagtgtcg	gcatgtaaat	780
tcagctcagt	gaatgcttat	taaacattaa	cgtaatcctt	gttgctttct	ttattcaacc	840
ctacagtga	attttctggg	tgcttgaggt	aactaaaata	tatttgaga	tatttaattt	900
caacttttac	acattaaaca	agtagaaaac	catgtcttca	aaaacatatt	ctttaaaggt	960
gattgtttag	ggcaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaa		1004

<210> 53

<211> 1305

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (329)

<223> n equals a,t,g, or c

<400> 53

gcggaacgst	ggtgtgggtt	tttaaaaaac	aaaacaagat	acatgctgac	atttctgggt	60
tggcaggcag	agcttggtct	gctccccacc	ctcccttttc	ccatagtaac	catttatagg	120
acatctcact	gttgtctact	ctgtgttgcc	tctgcttccc	tgcttggtag	atctaggaat	180
cttaggattt	cttagtttta	gctggtgatc	cgtatctttt	tcttaattcc	attgtaacct	240
cagcttttct	tattgcttgt	aggaaggctg	tttccattga	atacaaaaca	aataaaagct	300
tttattctta	atcttagaga	taggatgtnt	gtatttaaaa	acaattgtgc	tgtcaaaatt	360
ctgtcaagtt	ggcttttacc	acattagttt	tttttaatgt	ggtttatatg	accctgragt	420
accttgtctt	ctcactgtta	aattctcaac	tgagttgtcc	ctattttaaag	tgtgagactg	480
tgccagtttg	attttaaaat	attgcaagtg	cgttatggca	agataaaaact	gcaaagaaag	540
aaccttcatg	tccctttgat	tataaatgct	tttggcactt	gtttctactt	tttcctaatt	600
ttttttgagg	aaagaacctc	caactctcca	gacaggctcg	ggggcaaatg	actaaaacat	660
gaactgaggc	cctgggctgt	ctctgtgagg	atatcccttc	tattctctct	gaaatgtccc	720
agcatgtggt	gcatttcttg	ttagtgtgga	ctcctctgta	tataacacat	cttattttatc	780
ttctgtgcat	aacatgaagt	agtgccctaa	tgcaattcca	ggatgtaatt	cagcattttct	840
ataaaaaatac	agtgtttttc	tacatttgca	tcaaaaaata	accagataat	tatatattatt	900
aagaaaaatag	catttttggc	tgggtgtggt	ggctcacgta	atcccagcac	tttgggaggc	960
cgaggcaggc	agatcacttg	aggtcaggag	tgaggcaggc	agatcacttg	agatcaggag	1020
ttcgagacca	gyctggccaa	catggtgaaa	ccccatctct	actaaaaatg	caaaattagc	1080
ctggcgtagt	ggtgcatgcc	tgtaatccta	gctactcagg	agactgaggc	aggagaatca	1140
cttgaacttg	ggaaggggag	attgcagtga	gctgagattg	tgccactgca	ctccagccta	1200
ggcaacagag	tgagactctg	tttcaaaaaa	aaaaaaaaaag	ggcgcccgct	ctagaggatc	1260
caagcttacc	gtacgccgtg	catgcgacgt	catagctctt	ctata		1305

<210> 54

<211> 813

<212> DNA

<213> Homo sapiens

<400> 54

gaagaaaacg tgaccgtaaa tatctgtagg atataggaac cagagagtag atggaacatg	60
actagtaaga gacttaaatc caggggacct caggaggtaa tacaaaagaa ttttgtatta	120
cggaattgtt tacctagaat ttgaactcgg gagagaatcc ctgtgtagga gtatatttct	180
gcaaagaaag tgcttaaaaga aatgggttctt ccttcaatct gtttctttga tttgtatcag	240
attaggggaag gggaagctat ttgttggaact tttcattttg gtaaaatctg aatgagtatt	300
gagaatggct cttgagacag tagtgcaact tatattgctt tcttacttgg tttttatgta	360
tgatttatta ataggcaaaa atctcattat ggtgagctta atgacaaatc agtttgtttt	420
aaacacattt tattaaaaata catttagttt aaaaagtaaa tttccaaact accagctgaa	480
tacaactgtc cagattcttg caaggaaaac caaatgctag agaaggccag gcgcgatggc	540
tcaactcctgt ggtcctagta ttttgggagg ccgaggcggg tggatcagtt tagctcagga	600
gttcgagatc agcttaggca gcatgggtgg accctgtctc tacagaaaat acaaaaatta	660
gccggagggg aggcattgtc ctgtagtccc agctactcgg gaggctacag cagggggatc	720
gcatgggcct gggaggcgga ggtttagtt agccaagatg acaccattgc attccagcct	780
gggtgacagg agtgaaatcc tgtctcaaaa aaa	813

<210> 55

<211> 1694

<212> DNA

<213> Homo sapiens

<400> 55

ccacgcgtcc gaagtaattt ggaaaaattt aacattccta gtgacttaag atttgattaa	60
tagccttggt ggtagtattt tatatattcc taaatactat tgtaaaatac tccctcaata	120
aatcctgcat gcctttaaaa gtccctctca aaataatctg ttatttcggc aggtaatgac	180
caatgtgttt tttgggtgga atctttcatc ggttttccac attgtttgta cagtgtggt	240
catcaactgta gccacgcttg tgcattgct gattgatgcc tcgggatagt tctagaactc	300
aatgtgagta cgtgcaaaaga ttaccctt cactctaaaa ttctctttaa agataatga	360
ttacatttaa cataagatgt attttcctta acaaaagtgt cacttttgaa gtgggaatca	420
aaatatgttt gtaatagtaa atatttcaat gatgattctg tgcactttgt gggactatat	480
agttttaaag tagtggttgt ttagagacat atggggctgc cacaactggg taggcagtgc	540
tgttggcctc tagtgataaa ggccaggat atacatccca caatgtgcag gtctctcaca	600
caaaaaatta tctgatccaa aatgtcaatt gtgctgaggt tgagaaaccc tggtttagag	660
tactttgcat atctcattta ctataacaca taaatgttac taaaaatagc tataaattaa	720
gtggatttgg actttgctga ataataatat tctagtgaat tttatgagaa atatgaaagg	780
attcaagtta tatccattca ctgtctatga caaaatttct ttttctttaa atatttttct	840
ttctccagat ctttctttaa tagtctgcac tgccatcaac caaatagaag tcctcataat	900
atcacagttg aattaatccc agctctgttt caactatcat gtatttaagt tctgctttca	960
gtttatcggc attttcctac cagagcaagt ataaattccg ttgcttctac ctctctgtct	1020
tctgtgtaaa actatttccc atttacttcc caaaatttat gttcagctct agtcaatcta	1080
attttttggc ctgtgaataa gccatatcaa ttctttccat tattctttgt cctatctgct	1140
ttttatttcc gataatgatt atttttctt catttctgtc caaattttac aaaaacttta	1200
agatccagtg caacttctaa ttcttttatt cacattcact gatcatatat ttattgagta	1260
attactatgt gccacacaaat agaataataa gatgaatgca ataagaaagg acctgtgcat	1320
tctcacaat aaacataaaa gttcaactgc aatatagggt atgaagacag aggaggccag	1380
gaacggtggc tcacgctgtg atctcagcat ttttggaggc caagggtggat ggattgcttg	1440
agcccaggag ttggagacca gcctgggcaa cgcggtgaga cccgcctct acaaaaaatt	1500
agccgggttt ggtggcatgt gcctgtggtc ccggctactt gagaggctga ggtgagagga	1560
ttgcttgagc ccgggaggta gaggtttcag tgagctagat gcaccactgc actccacctg	1620
ggcaacagag tgagaccggg tctcaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa	1680

aaaaaaaaaa aaaa

1694

<210> 56

<211> 865

<212> DNA

<213> Homo sapiens

<400> 56

ggcacgaggt gaggtgtgcc ttttaccttc tgccatgatt gtgaagcctc cccaccacat	60
ggaacggtga cagattgact gggatcccct cgcacatcct caacagctcc ccatcagacc	120
ggcagattaa ccagctggcc cagaggctgg gccctgagtg ggagcccatg gtgctgtctc	180
tgggactgtc ccagacggat atctaccgct gtaaggccaa ccacccccac aacgtgcagt	240
cgcagggtgt ggaggccttc atcctgtggc ggacgcgctt cgggaagcag gccaccttcc	300
agagcctgca caacgggctg cgggctgtgg aggtggaccc ctgcgtgctc ctgcacatgt	360
tggagtgatg gtgcctccag caaccgctgg ggagtgtgtc cctgagtcac gtgggctgaa	420
tcctgacttt cactcagagc aggtgggttt ttgtgtaggt ttgtttttta tttttgatga	480
tcttcagatg gaaggagaaa acagggtttc cactagacat tacttgaaag gccagattac	540
tcagcagatc tcccatgttg gctcaacaat tctttgtttt taattgcttg aagattgcat	600
tgttgaatt gtccagtttt taaatgtgta atggcatttt aatagactag taaatcacag	660
tggttcaaaa tatatatcca tatatatata tatccatata tatatctcat gtcatcacat	720
tacaggcagg tgtctcatat gttaaacatt tacctgaatg ttgtctgagg actgaactgt	780
ggactttact attcataatg ataaaataat aaaatgcgaa ttactattta taatgtgcct	840
cactcatgag aaaaaaaaaa aaaaaa	865

<210> 57

<211> 337

<212> DNA

<213> Homo sapiens

<400> 57

ggcacgagcc agctttaaaa atataaatat aattcttttc atttttagatg gaaatcattt	60
atgtcacatt actcataaat gtcgtggtgg tgcagtccctg taattcttgg agatctctca	120
ggcagatgtc acccaagtat tctacataat atgattttcca tctaaaatga aaagaattat	180
attttatattt ttaaagctgg tatactgctt ttgcaagtta tttttttttt aaagcaaagg	240
tattttaaag gacaagtaac aagggaacaga gtaactatta aattaattat cttcttttaa	300
agtgcacaaa aaaaaaaaaa aaaaaaaaaa aaaaaa	337

<210> 58

<211> 1777

<212> DNA

<213> Homo sapiens

<400> 58

ggcacgagac tccatggatt ttgatgacac gtggcaccct gccacccacc cttctggggc	60
tgctcttctt gtcctcacag ctttagcaga agccctgccca aggagtccaa agttttcttg	120
ccttgacctg ctgctggctt tcaatgttgg tattgaagtg caaggccgat tactgcattt	180
cgccaaggag gccaatgaca tgccaaagag attccatccc ccttccgtgg taggaacgtt	240
gggtagtgtc gctgctgcat ccaagttttt aggaacttagc tcgacaaagt gccgagaagc	300
tctggccatt gctgtttccc atgctggggc acccatggcc aatgctgccca cccagaccaa	360
gcccctccac attggcaatg ctgccaaagca tgggatagaa gctgcatttt tggcaatgtt	420
gggtctccaa ggaacaagc aggtcttggc cttggaggca ggatttgggg ccttttatgc	480
caactattcc ccaaaagtcc ttccaagcat agcttcctac agttggctgc tggaccagca	540

```

ggacgtggcc tttaagcggt ttctgcaca tttatctacc cactgggtgg cagacgcagc      600
tgcattctgtg agaaaagcacc ttgtaccaga gagagccctg cttccaactg actacattaa      660
gagaattgtg ctcaggatac caaatgtcca gtatgtaaac aggccctttc cagtttcgga      720
gcatgaagcc cgtcattcat tccagtatgt ggcctgtgcc atgctgcttg atggtggcat      780
cactgtcccc tcattccatg aatgccagat caacaggcca cagggtgagag agctgctcag      840
taaggtggag ctggagtacc ctccggacaa cttgccaagc ttcaacatac tgtactgtga      900
aataagtgtc accctcaagg atggagccac cttcacagat cgctctgata cttctatgg      960
gcaactggaga aaaccactga gccaggagga cctagaggaa aagttcagag ccaatgcctc     1020
caagatgtcg tcctgggaca cagtggaaag ccttataaaag atagtcaaaa atctagaaga     1080
ctagaagact gttctgtgtt aactacactt ctcaaaggac ctctccacca gaggtagctt     1140
caaactctcc agcatgtaat aattctatca caaatctctc ctgaggctta ccaacatcta     1200
aatgactttg catttgggga gattcaatga tttggtttgt aaagcaaggg tctgctgctt     1260
ggttttccca ggaaaaatga acaaagatgg agagagtcca gaaacagaac tacatatatc     1320
tggaaggagc cttctcctga aaattttgca ggacagttcc acttacctaa atcagatgaa     1380
acacacacac aaaaatgagt ttgtaagcat tcacaagggt gaaattcaac tcacctgtga     1440
tttacttata aaattaatct cttcatagga attatgtgtg gacttcatga gcctcaaggt     1500
tttagaggga tgtgaacctg catgtatatt ttctgacagt ggagaggggt ctggtgcatt     1560
gtgtcaccaa cagatctcct agaccatggc ttattaccaa gccctccaca gtgcaagggg     1620
tgctactggg gaatgggtgg gtttaaatcc tgccctctgc attcactaga ttagccttg     1680
agcatgttac cattagccct ctgcctcagt ttccctattt gtcaagccga agtaaaaagc     1740
agtctggaaa aatcgcaaaa aaaaaaaaaa aaaaaaa      1777

```

<210> 59

<211> 51

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (51)

<223> Xaa equals stop translation

<400> 59

```

Met Trp Asn Ile Phe Ser Tyr Val Cys Trp Leu Leu Val Cys Leu Leu
  1               5               10              15

```

```

Leu Arg Ser Val Cys Ser Cys Leu Leu Pro Ser Phe Lys Trp Asp Leu
      20               25              30

```

```

Phe Phe Ala Cys Ser Leu Val His Thr Phe Phe Phe Phe Ile Asp Ser
  35               40              45

```

```

Gly Cys Xaa
  50

```

<210> 60

<211> 21

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (21)

<223> Xaa equals stop translation

<400> 60

Ser Gly Glu Gly Ala Trp Val Pro Gly Ala Ser Leu Ala Leu His Gln
1 5 10 15

Asp Pro Val Glu Xaa
20

<210> 61

<211> 58

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (58)

<223> Xaa equals stop translation

<400> 61

Met Trp Leu Leu Lys Pro Ser Ala His Ser Pro Val His Val Leu Val
1 5 10 15

Leu Leu Phe Pro Arg Gly Trp Ser Gln Pro Gly Thr His Lys Arg Gln
20 25 30

Ile Leu Val Asn Ala Ala Ser Leu Pro Gly Gly Cys Leu Leu Pro Trp
35 40 45

Ile Trp Ser Gly Ala Ala Leu Arg Phe Xaa
50 55

<210> 62

<211> 12

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (12)

<223> Xaa equals stop translation

<400> 62

Met Val Leu Phe Leu Leu Arg Phe Leu Phe Leu Xaa
1 5 10

<210> 63

<211> 16

<212> PRT

<213> Homo sapiens

<220>

<221> SITE
 <222> (16)
 <223> Xaa equals stop translation

<400> 63

Met Glu Thr His Arg Gln Gln Leu Arg Lys Met Val Cys Gln Gln Xaa
 1 5 10 15

<210> 64
 <211> 37
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (37)
 <223> Xaa equals stop translation

<400> 64

Met Leu Ser Asp Trp Leu Ile Ile Val Leu Gln Cys Tyr Val Gln Val
 1 5 10 15

Thr Leu Ile Leu Leu Ile Val Val Pro Arg Cys Lys Ser Ser Asp Ala
 20 25 30

Asp Ile Leu Leu Xaa
 35

<210> 65
 <211> 39
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (39)
 <223> Xaa equals stop translation

<400> 65

Met Ile Ser Cys Leu Asn Ile Leu Arg Val Leu Tyr Leu Leu Trp Gly
 1 5 10 15

Leu Leu Ala Leu Ser Ala Leu Thr Gln Ile Ile Gly Tyr Ile Thr Trp
 20 25 30

Leu Met Phe Leu Tyr Thr Xaa
 35

<210> 66

<211> 59
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (59)
 <223> Xaa equals stop translation

<400> 66
 Met Leu Arg Gln Glu Ile Cys Leu Ile Arg Thr Gly Ser Ser Val Leu
 1 5 10 15
 Ser Val Thr Leu Val Ala Leu Leu Leu Gln Val Ile Thr Leu Val Met
 20 25 30
 Tyr Met Thr Leu Arg Ser Lys Arg Gly Leu Leu Thr Met Thr Trp Arg
 35 40 45
 Tyr Gln Lys Ser Lys Arg Leu Pro Cys Lys Xaa
 50 55

<210> 67
 <211> 94
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (94)
 <223> Xaa equals stop translation

<400> 67
 Met Trp Ile Arg Val Gly Phe Leu Val Phe Lys Thr Pro Gly Leu Arg
 1 5 10 15
 Thr Pro Ala Ala Gly Glu Arg Ile Tyr Asn Ile Ser Gly Asn Gly Ser
 20 25 30
 Pro Leu Ala Asp Ser Lys Glu Ile Phe Leu Thr Val Pro Val Gly Gly
 35 40 45
 Gly Glu Ser Leu Arg Leu Leu Ala Ser Asp Leu Gln Arg His Ser Ile
 50 55 60
 Ala Gln Leu Asp Pro Glu Ala Leu Gly Asn Ile Lys Lys Leu Ser Asn
 65 70 75 80
 Arg Leu Ala Gln Ile Cys Ser Ser Ile Arg Thr His Lys Xaa
 85 90

<210> 68
 <211> 24

<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (24)
<223> Xaa equals stop translation

<400> 68
Met Leu Leu Leu Gln Ser Leu Phe Phe Ser His Glu Leu Gly Val Gly
1 5 10 15
Trp Gly Arg Glu Arg Glu Gly Xaa
20

<210> 69
<211> 22
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (22)
<223> Xaa equals stop translation

<400> 69
Met Pro Cys Phe Ser Leu Leu Ser Leu Pro Leu Trp Asp Pro Leu Val
1 5 10 15
Ile Leu Val Phe Cys Xaa
20

<210> 70
<211> 34
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (34)
<223> Xaa equals stop translation

<400> 70
Met Ser Asp Arg Trp Ser Pro Phe Ile Pro Phe Leu Leu Leu Ala Pro
1 5 10 15
Val Ser Ser Gly Ser Gly His Leu Thr Phe Ser Cys Pro Ala Gly Ser
20 25 30

Ala Xaa

<210> 71
<211> 53
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (53)
<223> Xaa equals stop translation

<400> 71
Met Ala Met Ala Met Ala Arg Ile Thr Pro Pro Thr Met Gly Met Val
1 5 10 15
Trp Pro Leu His Thr Leu Gly Lys Cys Leu Ala Leu Thr Gln Met Gln
20 25 30
Thr Leu Val Pro Arg Val Ala Pro Val Pro Ile Pro Phe Tyr Pro Glu
35 40 45
Leu Thr Ser Ala Xaa
50

<210> 72
<211> 36
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (36)
<223> Xaa equals stop translation

<400> 72
Met Leu Ser Ile Met Leu Cys Phe Leu Trp Asn Met Ile Ile Leu Leu
1 5 10 15
Val Ala Ser Ser Ala Tyr Ser Gly Cys Asp Leu Ala Leu Pro Gly Thr
20 25 30
Ser Ala Leu Xaa
35

<210> 73
<211> 32
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (32)
<223> Xaa equals stop translation

<400> 73

Met Tyr Asn Leu Ser Ser Leu Phe Met Ile Ser Phe Leu Val Cys His
 1 5 10 15
 Val Thr Pro Ser Gln Thr Leu Lys Gly Pro Pro Leu Ser Trp Ser Xaa
 20 25 30

<210> 74

<211> 42

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (42)

<223> Xaa equals stop translation

<400> 74

Met Tyr Met Lys Met Met Phe Met Leu Phe Ile Ile Leu Pro Phe Ile
 1 5 10 15
 Ile Ser Phe Phe Ile Val Leu Ile Ala Met Ser Phe Ser Ser Leu Ile
 20 25 30
 Phe Phe Pro Gln Cys Leu Ile Cys His Xaa
 35 40

<210> 75

<211> 25

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (25)

<223> Xaa equals stop translation

<400> 75

Met Ala Ala Ser Ala Leu Leu Leu Cys Val Val Thr Leu Ile Leu Phe
 1 5 10 15
 Leu Val Leu His Tyr Ile Val Ser Xaa
 20 25

<210> 76

<211> 9

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (9)

<223> Xaa equals stop translation

<400> 76

Met Ser Asn Val Gln Leu Gln Arg Xaa

1 5

<210> 77

<211> 25

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (25)

<223> Xaa equals stop translation

<400> 77

Ser Phe Phe Val Phe Leu Gly Asp Leu Tyr Phe Phe Phe Gly Glu Met

1 5 10 15

Ser Ile Pro Ile Leu Ala His Phe Xaa

20 25

<210> 78

<211> 66

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (66)

<223> Xaa equals stop translation

<400> 78

Gly Ser Phe Pro Ser Pro Lys His Arg Gln Arg Gly Gly Glu Gln Phe

1 5 10 15

Leu Val Leu Phe Leu Phe Leu Lys Trp Cys Leu Tyr Leu Gln Pro Pro

20 25 30

Gly Gly Leu Pro Trp Pro His Phe Ser Ala Pro Pro Arg His Arg His

35 40 45

Pro Ser Thr Leu Leu His Val Thr Arg Lys Met Pro Phe Ala Glu Cys

50 55 60

Thr Xaa

65

<210> 79
 <211> 91
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (91)
 <223> Xaa equals stop translation

<400> 79
 Met Cys Pro Val Ser Gln Phe Pro Gly Ser Ser Ser Val Cys Cys Pro
 1 5 10 15
 Phe Ser Ser Ser Gly Ser Ile Val Arg Glu Pro Arg Met Glu Ala Lys
 20 25 30
 Cys Thr Gly His Trp Leu Phe Phe Gln Cys Pro Ser Asp Ser Pro Cys
 35 40 45
 Pro Gly Gly Leu Val Pro Ser Leu Ser Val Trp Cys Leu Phe Tyr Lys
 50 55 60
 Leu Val Met Thr Ser Gly Asn Gly Pro Gly Phe Gln Ile Ala Ile Pro
 65 70 75 80
 Gly Asp Ile Leu Ile Leu Trp Phe Lys Pro Xaa
 85 90

<210> 80
 <211> 7
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (7)
 <223> Xaa equals stop translation

<400> 80
 Met Phe Leu Arg Glu Gln Xaa
 1 5

<210> 81
 <211> 34
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (34)
 <223> Xaa equals stop translation

<400> 81

Met Tyr Tyr Phe Ile Phe Leu Phe Phe Ser Tyr Val Leu Cys Phe Arg
1 5 10 15

Ile Phe Leu Glu Ala Ser Ser Lys Ser Cys Phe Val Gly Asn Lys Gln
20 25 30

Ser Xaa

<210> 82

<211> 33

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (33)

<223> Xaa equals stop translation

<400> 82

Thr Ser Cys Cys Ile Ile Ile Phe Leu Cys Ala Glu Gln Leu Lys Thr
1 5 10 15

Gln Pro Gln Tyr Ala Asn Cys Phe Asn Ala Ile Leu Arg Gln Leu Ile
20 25 30

Xaa

<210> 83

<211> 11

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (11)

<223> Xaa equals stop translation

<400> 83

Met Leu Leu Tyr Leu Thr Val Gly Tyr Trp Xaa
1 5 10

<210> 84

<211> 35

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (35)

<223> Xaa equals stop translation

<400> 84

Met Gly Leu Leu Trp Leu Ser Val Gly His Thr Ser Pro Gln Lys
1 5 10 15

Ala Pro Ala Lys Glu Leu Arg Phe Trp Leu Gly Lys Asn Ile Thr Pro
20 25 30

Leu Gln Xaa
35

<210> 85

<211> 38

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (38)

<223> Xaa equals stop translation

<400> 85

Met Leu Leu Ile Phe Thr Phe Ser Ala Leu Val Leu Ser Tyr Pro Leu
1 5 10 15

Leu Ile Leu Gly Asp Trp Val Glu Met Ala Ile Glu His His Thr Leu
20 25 30

Leu Thr Lys Thr Ile Xaa
35

<210> 86

<211> 31

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (31)

<223> Xaa equals stop translation

<400> 86

Met Ala Leu Val Gly Leu Leu Ser Ala Gly Val Pro Gly Val Ser Leu
1 5 10 15

Cys Val Gln Ile Phe Ser Tyr Lys Asp Thr Gly Glu Ile Gly Xaa
20 25 30

<210> 87

<211> 3

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (3)

<223> Xaa equals stop translation

<400> 87

Met Leu Xaa

1

<210> 88

<211> 13

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (13)

<223> Xaa equals stop translation

<400> 88

Val Leu Leu Leu Pro His Val Leu Ser Gly Gly Leu Xaa

1

5

10

<210> 89

<211> 36

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (36)

<223> Xaa equals stop translation

<400> 89

Leu Ile Val Phe His Ile Ile Phe Ile Pro Trp Ile Thr Leu Leu Cys

1

5

10

15

Val Phe Ile Gly Val Arg Leu Leu Ala Val Ser Tyr Glu Thr Thr Tyr

20

25

30

Tyr Ile Phe Xaa

35

<210> 90

<211> 20

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (20)

<223> Xaa equals stop translation

<400> 90

Met	Thr	Ser	Tyr	Cys	Ser	Phe	Asp	Leu	His	Phe	Ser	Asp	Asp	Asn	Tyr
1				5					10					15	

Val	Glu	His	Xaa
			20

<210> 91

<211> 35

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (35)

<223> Xaa equals stop translation

<400> 91

Met	Asn	Phe	Trp	His	Ser	Cys	Tyr	Leu	Leu	Arg	Asn	Cys	Thr	Val	Cys
1				5					10					15	

Phe	Leu	Cys	Ser	Ile	Phe	Phe	Phe	Phe	Pro	Gly	Met	Arg	Met	Tyr	Leu
			20					25						30	

Ser	Ser	Xaa
		35

<210> 92

<211> 35

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (35)

<223> Xaa equals stop translation

<400> 92

Met	Lys	Leu	Thr	Leu	Gly	Thr	Ala	Gly	Ser	Leu	Phe	Pro	Gln	Ala	Leu
1				5					10					15	

Tyr	Ile	Leu	Leu	Asp	Phe	Ile	Trp	Val	Asn	Phe	Ile	Asp	Gly	Ser	His
			20					25					30		

Tyr	Ile	Xaa
		35

<210> 93

<211> 48

44

<212> PRT

<213> Homo sapiens

<400> 93

Met Ala Met Lys Ile Cys Gln Trp Ser Phe Val Cys Gly Leu Leu Gly
1 5 10 15

Thr Val Phe Leu Leu Cys Leu Val Leu Phe Tyr Phe Cys Tyr Ser Arg
20 25 30

Lys Leu Arg Ala His Leu Lys Thr Lys Lys Lys Lys Lys Lys Lys
35 40 45

<210> 94

<211> 10

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (10)

<223> Xaa equals stop translation

<400> 94

Met Asn Lys Ile Lys Lys Trp Leu Ile Xaa
1 5 10

<210> 95

<211> 23

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (23)

<223> Xaa equals stop translation

<400> 95

Met Val Arg Leu Ser Ser Leu Gln Thr Leu Leu Cys Pro Asp Ser His
1 5 10 15

Leu Gly His Phe Ile Gln Xaa
20

<210> 96

<211> 21

<212> PRT

<213> Homo sapiens

45

<220>
 <221> SITE
 <222> (21)
 <223> Xaa equals stop translation

<400> 96
 Met Glu Leu Phe Phe Phe Trp Leu Thr Ile Leu Leu Phe Pro Thr Val
 1 5 10 15
 Phe Asn Asn Cys Xaa
 20

<210> 97
 <211> 42
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (42)
 <223> Xaa equals stop translation

<400> 97
 Met Gly Asn Val Met Val Thr Phe Ser Arg Leu Ser Cys Leu Ile Pro
 1 5 10 15
 Ser Ala Ser Ser Leu Leu Cys Leu Asn Ser Cys Thr Gly Cys Leu Val
 20 25 30
 His Val His Ile Thr Lys Arg Trp Tyr Xaa
 35 40

<210> 98
 <211> 38
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (38)
 <223> Xaa equals stop translation

<400> 98
 Met Lys Ser Ala Leu Arg Leu Val Leu Leu Phe Ser Phe His Phe Phe
 1 5 10 15
 Pro Leu Ile Ile Ser Phe Arg Arg Lys Arg Glu Gly Lys Lys Lys Lys
 20 25 30
 Lys Ile Arg Asp Leu Xaa
 35

46

<210> 99
 <211> 45
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (45)
 <223> Xaa equals stop translation

<400> 99
 Met Asn His Thr Leu Arg Asn Gln Cys Gly Cys Pro Leu Arg Val Leu
 1 5 10 15
 Leu Phe Phe Leu Leu Pro Leu Ser Lys Ile Arg Tyr Ser Val Ile Lys
 20 25 30
 Tyr Ile Ser Ile Lys Val Phe Lys Ser Asp Leu Tyr Xaa
 35 40 45

<210> 100
 <211> 41
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (41)
 <223> Xaa equals stop translation

<400> 100
 Met Ser Lys Gly Leu Arg Lys Glu Ser His Val Phe Phe Leu Leu Phe
 1 5 10 15
 Ser Asn Leu Val Ile Thr Lys Gln Cys Tyr Gln Cys Ile Thr Tyr Lys
 20 25 30
 His Phe Ile Ile His Phe His Phe Xaa
 35 40

<210> 101
 <211> 51
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (51)
 <223> Xaa equals stop translation

<400> 101
 Met Leu Thr Phe Leu Val Trp Gln Ala Glu Leu Val Leu Leu Pro Thr
 1 5 10 15

47

Leu Pro Phe Pro Ile Val Thr Ile Tyr Arg Thr Ser His Cys Cys Leu
20 25 30

Leu Cys Val Ala Ser Ala Ser Leu Pro Gly Arg Ser Arg Asn Leu Arg
35 40 45

Ile Ser Xaa
50

<210> 102
<211> 43
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (43)
<223> Xaa equals stop translation

<400> 102
Ala Leu Glu Thr Val Val His Phe Ile Leu Leu Ser Leu Leu Val Phe
1 5 10 15

Met Tyr Asp Leu Leu Ile Gly Lys Asn Leu Ile Met Val Ser Leu Met
20 25 30

Thr Asn Gln Phe Val Leu Asn Thr Phe Tyr Xaa
35 40

<210> 103
<211> 18
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (18)
<223> Xaa equals stop translation

<400> 103
Met Val Ile Thr Val Ala Thr Leu Val Ser Leu Leu Ile Asp Ala Ser
1 5 10 15

Gly Xaa

<210> 104
<211> 7
<212> PRT
<213> Homo sapiens

Asn Val Gln Lys Ser Gln Ala Phe Leu Tyr Thr Asn Asn Arg Gln Thr
 1 5 10 15
 Glu Ser Gln Ile Met Ser Glu Leu Pro Phe Thr Ile Ala Ser Lys Arg
 20 25 30
 Ile Lys Tyr Leu Gly Ile Gln Leu Thr Arg Asp Val Lys Asp Leu Phe
 35 40 45
 Lys Glu Asn Tyr Lys Pro Leu Leu Lys Glu Ile Lys Glu Asp Thr Asn
 50 55 60
 Lys Trp Lys Asn Ile Pro Cys Ser Trp Val Gly Arg Ile Asn Ile Val
 65 70 75 80
 Lys Met Ala Ile Leu Pro Lys Val Ile Tyr Arg Phe Asn Ala Ile Pro
 85 90 95
 Ile Lys Leu Pro Met Thr Phe Phe Thr Glu Leu Glu Lys Thr Thr Leu
 100 105 110
 Lys Phe Ile Trp Asn Gln Lys Arg Ala Arg Ile Ala Lys Ser Ile Leu
 115 120 125
 Ser Gln Lys Asn Lys Ala Gly Gly Ile Thr Leu Pro Asp Phe Lys Leu
 130 135 140
 Tyr Tyr Lys Ala Thr Val Thr Lys Thr Ala Trp Tyr Trp Tyr Gln Asn
 145 150 155 160
 Arg Asp Ile Asp Gln Trp Asn Arg Thr Glu Pro Ser Glu Ile Met Pro
 165 170 175
 His Ile Tyr Asn Tyr Leu Ile Phe Asp Lys Pro Glu Lys Asn Lys Gln
 180 185 190
 Trp Gly Lys Asp Ser Leu Phe Asn Lys Trp Cys Trp Glu Asn Trp Leu
 195 200 205
 Ala Ile Cys Arg Lys Leu Lys Leu Asp Pro Phe Leu Thr Pro Tyr Thr
 210 215 220
 Lys Ile Asn Ser Arg Trp Ile Lys Asp Leu Asn Val Arg Pro Lys Thr
 225 230 235 240
 Ile Lys Thr Leu Glu Glu Asn Leu Gly Ile Thr Ile Glu Asp Ile Gly
 245 250 255
 Val Gly Lys Asp Phe Met Ser Lys Thr Pro Lys Ala Met Ala Thr Lys
 260 265 270
 Ala Lys Ile Asp Lys Trp Asp Leu Ile Lys Leu Lys Ser Phe Cys Thr
 275 280 285
 Ala Lys Glu Thr Thr Ile Arg Val Asn Arg Gln Pro Thr Thr Trp Glu

290	295	300
Lys Ile Phe Ala Thr Tyr Ser Ser Asp Lys Gly Leu Ile Ser Arg Ile		
305	310	315 320

Tyr

<210> 108
<211> 327
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (7)
<223> Xaa equals stop translation

<220>
<221> SITE
<222> (53)
<223> Xaa equals stop translation

<220>
<221> SITE
<222> (100)
<223> Xaa equals stop translation

<220>
<221> SITE
<222> (151)
<223> Xaa equals stop translation

<220>
<221> SITE
<222> (175)
<223> Xaa equals stop translation

<220>
<221> SITE
<222> (205)
<223> Xaa equals stop translation

<220>
<221> SITE
<222> (223)
<223> Xaa equals stop translation

<220>
<221> SITE
<222> (286)
<223> Xaa equals stop translation

<400> 108

Ser Met Tyr Lys Asn Gln Xaa His Phe Tyr Thr Pro Ile Thr Ser Lys
 1 5 10 15
 Leu Glu Ser Gln Met Lys Asn Thr Ile Leu Phe Thr Ile Ala Ile Lys
 20 25 30
 Lys Thr Lys Tyr Leu Glu Ile His Leu Thr Lys Glu Val Lys Asp Leu
 35 40 45
 Tyr Lys Glu Asn Xaa Glu Thr Leu Leu Lys Glu Ile Thr Asp Asp Thr
 50 55 60
 Asp Lys Trp Lys Asn Ile Pro Cys Ser Trp Ile Thr Arg Ile Asn Ile
 65 70 75 80
 Val Lys Met Ala Ile Leu Leu Lys Ala Ile Tyr Arg Phe Asn Val Ile
 85 90 95
 Pro Ile Lys Xaa Pro Val Ser Phe Phe Thr Glu Leu Glu Lys Thr Ile
 100 105 110
 Leu Lys Phe Ile Trp Asn Gln Lys Arg Thr Gln Ile Thr Lys Ala Ile
 115 120 125
 Pro Ser Lys Lys Asn Lys Ala Glu Ala Ser His Tyr Leu Thr Leu Tyr
 130 135 140
 Tyr Thr Ile Lys Leu Gln Xaa Pro Lys Gln His Gly Thr Gly Thr Lys
 145 150 155 160
 Ala Asp Lys Gln Thr Lys Gly Thr Glu Leu Arg Thr Gln Lys Xaa Ser
 165 170 175
 Cys Thr Pro Thr Ala Thr Cys Ser Leu Thr Lys Ser Thr Lys Asn Arg
 180 185 190
 Gln Trp Gly Lys Asp Ser Leu Phe Ser Lys Trp Cys Xaa Asp Ser Trp
 195 200 205
 Leu Ala Ile Cys Lys Arg Met Lys Gln Arg Phe Ile Leu Trp Xaa Lys
 210 215 220
 Asp Pro Tyr Leu Ser Pro Tyr Thr Lys Ile Asn Ser Arg Trp Ile Lys
 225 230 235 240
 Tyr Leu Ser Val Arg Pro Gln Thr Ile Arg Ile Leu Gln Glu Asn Leu
 245 250 255
 Gly Asn Thr Ile Leu Asp Ile Gly Ser Gly Lys Glu Phe Met Thr Lys
 260 265 270
 Cys Leu Lys Ala Thr Ala Thr Lys Ile Lys Ile Asp Lys Xaa Asp Leu
 275 280 285
 Ile Lys Glu Leu Leu His Tyr Lys Arg Asn Phe Gln Trp Ser Lys Gln

290 295 300
 Thr Thr Tyr Lys Ile Glu Lys Met Phe Ala Asn Tyr Ala Ser Asp Lys
 305 310 315 320
 Gly Leu Thr Ser Arg Ile Tyr
 325

 <210> 109
 <211> 286
 <212> PRT
 <213> Homo sapiens

 <400> 109
 Leu Phe Ile Glu Arg Gly Arg His Glu Thr Thr Trp Ala Val Leu Arg
 1 5 10 15
 Lys Phe Gly Tyr Glu Thr Ser Leu Lys Leu Ser Glu Asp Tyr Leu Tyr
 20 25 30
 Pro Arg Ile Thr Ile Pro Val Gly Cys Ser Thr Glu Leu Ser Pro Glu
 35 40 45
 Gly Val Gln Phe Val Ser Ala Leu Phe Glu Lys Tyr Asp Glu Asp Lys
 50 55 60
 Asp Gly Cys Leu Ser Pro Ser Glu Leu Gln Asn Leu Phe Ser Val Cys
 65 70 75 80
 Pro Val Pro Val Ile Thr Lys Asp Asn Ile Leu Ala Leu Glu Thr Asn
 85 90 95
 Gln Arg Gly Trp Leu Thr Tyr Asn Gly Tyr Met Ala Tyr Trp Asn Met
 100 105 110
 Thr Thr Leu Ile Asn Leu Thr Gln Thr Phe Glu Gln Leu Ala Tyr Leu
 115 120 125
 Gly Phe Pro Val Gly Arg Ser Gly Pro Gly Arg Ala Gly Asn Thr Leu
 130 135 140
 Asp Ser Ile Arg Val Thr Arg Glu Arg Lys Lys Asp Leu Glu Asn His
 145 150 155 160
 Gly Thr Asp Arg Lys Val Phe Gln Cys Leu Val Val Gly Ala Lys Asp
 165 170 175
 Ala Gly Lys Thr Val Phe Met Gln Ser Leu Ala Gly Arg Gly Met Ala
 180 185 190
 Asp Val Ala Gln Ile Gly Arg Arg His Ser Pro Phe Val Ile Asn Arg
 195 200 205
 Val Arg Val Lys Glu Glu Ser Lys Tyr Leu Leu Leu Arg Glu Val Asp

210	215	220
Val Leu Ser Pro Gln Asp Ala Leu Gly Ser Gly Glu Thr Ser Ala Asp		
225	230	235 240
Val Val Ala Phe Leu Tyr Asp Ile Ser Asn Pro Asp Ser Phe Ala Phe		
	245	250 255
Cys Ala Thr Val Tyr Gln Lys Tyr Phe Tyr Arg Thr Lys Thr Pro Cys		
	260	265 270
Val Met Ile Ala Thr Lys Val Glu Arg Glu Glu Val Asp Gln		
	275	280 285

<210> 110
 <211> 277
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (159)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (160)
 <223> Xaa equals any of the naturally occurring L-amino acids

<400> 110
Leu Phe Ile Gln Arg Gly Arg His Glu Thr Thr Trp Thr Ile Leu Arg
1 5 10 15
Arg Phe Gly Tyr Ser Asp Ala Leu Glu Leu Thr Ala Asp Tyr Leu Ser
20 25 30
Pro Leu Ile His Val Pro Pro Gly Cys Ser Thr Glu Leu Asn His Leu
35 40 45
Gly Tyr Gln Phe Val Gln Arg Val Phe Glu Lys His Asp Gln Asp Arg
50 55 60
Asp Gly Ala Leu Ser Pro Val Glu Leu Gln Ser Leu Phe Ser Val Phe
65 70 75 80
Pro Ala Ala Pro Trp Gly Pro Glu Leu Pro Arg Thr Val Arg Thr Glu
85 90 95
Ala Gly Arg Leu Pro Leu His Gly Tyr Leu Cys Gln Trp Thr Leu Val
100 105 110
Thr Tyr Leu Asp Val Arg Ser Cys Leu Gly His Leu Gly Tyr Leu Gly
115 120 125

Tyr Pro Thr Leu Cys Glu Gln Asp Gln Ala His Ala Ile Thr Val Thr
 130 135 140
 Arg Glu Lys Arg Leu Asp Gln Glu Lys Gly Gln Thr Gln Arg Xaa Xaa
 145 150 155 160
 Leu Leu Cys Lys Val Val Gly Ala Arg Gly Val Gly Lys Ser Ala Phe
 165 170 175
 Leu Gln Ala Phe Leu Gly Arg Gly Leu Gly His Gln Asp Thr Arg Glu
 180 185 190
 Gln Pro Pro Gly Tyr Ala Ile Asp Thr Val Gln Val Asn Gly Gln Glu
 195 200 205
 Lys Tyr Leu Ile Leu Cys Glu Val Gly Thr Asp Gly Leu Leu Ala Thr
 210 215 220
 Ser Leu Asp Ala Thr Cys Asp Val Ala Cys Leu Met Phe Asp Gly Ser
 225 230 235 240
 Asp Pro Lys Ser Phe Ala His Cys Ala Ser Val Tyr Lys His His Tyr
 245 250 255
 Met Asp Gly Gln Thr Pro Cys Leu Phe Val Ser Ser Lys Ala Asp Leu
 260 265 270
 Pro Glu Val Ser Arg
 275

<210> 111

<211> 159

<212> PRT

<213> Homo sapiens

<400> 111

Met Trp Lys Val Ser Ala Leu Leu Phe Val Leu Gly Ser Ala Ser Leu
 1 5 10 15
 Trp Val Leu Ala Glu Gly Ala Ser Thr Gly Gln Pro Glu Asp Asp Thr
 20 25 30
 Glu Thr Thr Gly Leu Glu Gly Gly Val Ala Met Pro Gly Ala Glu Asp
 35 40 45
 Asp Val Val Thr Pro Gly Thr Ser Glu Asp Arg Tyr Lys Ser Gly Leu
 50 55 60
 Thr Thr Leu Val Ala Thr Ser Val Asn Ser Val Thr Gly Ile Arg Ile
 65 70 75 80
 Glu Asp Leu Pro Thr Ser Glu Ser Thr Val His Ala Gln Glu Gln Ser
 85 90 95

55

Pro Ser Ala Thr Ala Ser Asn Val Ala Thr Ser His Ser Thr Glu Lys
 100 105 110

Val Asp Gly Asp Thr Gln Thr Thr Val Glu Lys Asp Gly Leu Ser Thr
 115 120 125

Val Thr Leu Val Gly Ile Ile Val Gly Val Leu Leu Ala Ile Gly Phe
 130 135 140

Ile Gly Gly Ile Ile Val Val Val Met Arg Lys Met Ser Gly Arg
 145 150 155

<210> 112

<211> 159

<212> PRT

<213> Homo sapiens

<400> 112

Met Trp Lys Val Ser Ala Leu Leu Phe Val Leu Gly Ser Ala Ser Leu
 1 5 10 15

Trp Val Leu Ala Glu Gly Ala Ser Thr Gly Gln Pro Glu Asp Asp Thr
 20 25 30

Glu Thr Thr Gly Leu Glu Gly Gly Val Ala Met Pro Gly Ala Glu Asp
 35 40 45

Asp Val Val Thr Pro Gly Thr Ser Glu Asp Arg Tyr Lys Ser Gly Leu
 50 55 60

Thr Thr Leu Val Ala Thr Ser Val Asn Ser Val Thr Gly Ile Arg Ile
 65 70 75 80

Glu Asp Leu Pro Thr Ser Glu Ser Thr Val His Ala Gln Glu Gln Ser
 85 90 95

Pro Ser Ala Thr Ala Ser Asn Val Ala Thr Ser His Ser Thr Glu Lys
 100 105 110

Val Asp Gly Asp Thr Gln Thr Thr Val Glu Lys Asp Gly Leu Ser Thr
 115 120 125

Val Thr Leu Val Gly Ile Ile Val Gly Val Leu Leu Ala Ile Gly Phe
 130 135 140

Ile Gly Gly Ile Ile Val Val Val Met Arg Lys Met Ser Gly Arg
 145 150 155

<210> 113

<211> 64

<212> PRT

<213> Homo sapiens

56

<400> 113

Ala Glu Glu Leu Lys Arg Asn Ala Glu Thr Gly Asn Leu Pro His Ser
 1 5 10 15

Tyr Arg Leu Ile Ser Val Val Ser His Ile Gly Ser Thr Ser Ser Ser
 20 25 30

Gly His Tyr Ile Ser Asp Val Tyr Asp Ile Lys Lys Gln Ala Trp Phe
 35 40 45

Thr Tyr Asn Asp Leu Glu Val Ser Lys Ile Gln Glu Ala Ser Val Gln
 50 55 60

<210> 114

<211> 64

<212> PRT

<213> Homo sapiens

<400> 114

Ala Glu Glu Leu Lys Arg Asn Ala Glu Thr Gly Asn Leu Pro His Ser
 1 5 10 15

Tyr Arg Leu Ile Ser Val Val Ser His Ile Gly Ser Thr Ser Ser Ser
 20 25 30

Gly His Tyr Ile Ser Asp Val Tyr Asp Ile Lys Lys Gln Ala Trp Phe
 35 40 45

Thr Tyr Asn Asp Leu Glu Val Ser Lys Ile Gln Glu Ala Ala Val Gln
 50 55 60

<210> 115

<211> 287

<212> PRT

<213> Homo sapiens

<400> 115

Glu Gln Gln Lys Pro Ala Ser Ser Asp Val Val Leu Pro Ala Thr Met
 1 5 10 15

Ser Tyr Thr Gly Phe Val Gln Gly Ser Glu Thr Thr Leu Gln Ser Thr
 20 25 30

Tyr Ser Asp Thr Ser Ala Gln Pro Thr Cys Asp Tyr Gly Tyr Gly Thr
 35 40 45

Trp Asn Ser Gly Thr Asn Arg Gly Tyr Glu Gly Tyr Gly Tyr Gly Tyr

50 55 60
 Gly Tyr Gly Gln Asp Asn Thr Thr Asn Tyr Gly Tyr Gly Met Ala Thr
 65 70 75 80
 Ser His Ser Trp Glu Met Pro Ser Ser Asp Thr Asn Ala Asn Thr Ser
 85 90 95
 Ala Ser Gly Ser Ala Ser Ala Asp Ser Val Leu Ser Arg Ile Asn Gln
 100 105 110
 Arg Leu Asp Met Val Pro His Leu Glu Thr Asp Met Met Gln Gly Gly
 115 120 125
 Val Tyr Gly Ser Gly Gly Glu Arg Tyr Asp Ser Tyr Glu Ser Cys Asp
 130 135 140
 Ser Arg Ala Val Leu Ser Glu Arg Asp Leu Tyr Arg Ser Gly Tyr Asp
 145 150 155 160
 Tyr Ser Glu Leu Asp Pro Glu Met Glu Met Ala Tyr Glu Gly Gln Tyr
 165 170 175
 Asp Ala Tyr Arg Asp Gln Phe Arg Met Arg Gly Asn Asp Thr Phe Gly
 180 185 190
 Pro Arg Ala Gln Gly Trp Ala Arg Asp Ala Arg Ser Gly Arg Pro Met
 195 200 205
 Ala Ser Gly Tyr Gly Arg Met Trp Glu Asp Pro Met Gly Ala Arg Gly
 210 215 220
 Gln Cys Met Ser Gly Ala Ser Arg Leu Pro Ser Leu Phe Ser Gln Asn
 225 230 235 240
 Ile Ile Pro Glu Tyr Gly Met Phe Gln Gly Met Arg Gly Gly Gly Ala
 245 250 255
 Phe Pro Gly Gly Ser Arg Phe Gly Phe Gly Phe Gly Asn Gly Met Lys
 260 265 270
 Gln Met Arg Arg Thr Trp Lys Thr Trp Thr Thr Ala Asp Phe Arg
 275 280 285

 <210> 116
 <211> 287
 <212> PRT
 <213> Homo sapiens

 <400> 116
 Glu Gln Gln Lys Pro Ala Ser Ser Asp Val Val Leu Pro Ala Thr Met
 1 5 10 15
 Ser Tyr Thr Gly Phe Val Gln Gly Ser Glu Thr Thr Leu Gln Ser Thr

58

20	25	30
Tyr Ser Asp Thr Ser Ala Gln Pro Thr Cys Asp Tyr Gly Tyr Gly Thr		
35	40	45
Trp Asn Ser Gly Thr Asn Arg Gly Tyr Glu Gly Tyr Gly Tyr Gly Tyr		
50	55	60
Gly Tyr Gly Gln Asp Asn Thr Thr Asn Tyr Gly Tyr Gly Met Ala Thr		
65	70	75
Ser His Ser Trp Glu Met Pro Ser Ser Asp Thr Asn Ala Asn Thr Ser		
85	90	95
Ala Ser Gly Ser Ala Ser Ala Asp Ser Val Leu Ser Arg Ile Asn Gln		
100	105	110
Arg Leu Asp Met Val Pro His Leu Glu Thr Asp Met Met Gln Gly Gly		
115	120	125
Val Tyr Gly Ser Gly Gly Glu Arg Tyr Asp Ser Tyr Glu Ser Cys Asp		
130	135	140
Ser Arg Ala Val Leu Ser Glu Arg Asp Leu Tyr Arg Ser Gly Tyr Asp		
145	150	155
Tyr Ser Glu Leu Asp Pro Glu Met Glu Met Ala Tyr Glu Gly Gln Tyr		
165	170	175
Asp Ala Tyr Arg Asp Gln Phe Arg Met Arg Gly Asn Asp Thr Phe Gly		
180	185	190
Pro Arg Ala Gln Gly Trp Ala Arg Asp Ala Arg Ser Gly Arg Pro Met		
195	200	205
Ala Ser Gly Tyr Gly Arg Met Trp Glu Asp Pro Met Gly Ala Arg Gly		
210	215	220
Gln Cys Met Ser Gly Ala Ser Arg Leu Pro Ser Leu Phe Ser Gln Asn		
225	230	235
Ile Ile Pro Glu Tyr Gly Met Phe Gln Gly Met Arg Gly Gly Gly Ala		
245	250	255
Phe Pro Gly Gly Ser Arg Phe Gly Phe Gly Phe Gly Asn Gly Met Lys		
260	265	270
Gln Met Arg Arg Thr Trp Lys Thr Trp Thr Thr Ala Asp Phe Arg		
275	280	285

<210> 117

<211> 56

<212> PRT

<213> Homo sapiens

<400> 117

Thr Pro Val Phe Ser Lys Ala Arg Tyr Thr Val Arg Ser Phe Gly Ile
 1 5 10 15

Arg Arg Asn Glu Lys Ile Ala Val His Cys Thr Val Arg Gly Ala Lys
 20 25 30

Ala Glu Glu Ile Leu Glu Lys Gly Leu Lys Val Arg Glu Tyr Glu Leu
 35 40 45

Arg Lys Asn Asn Phe Ser Asp Thr
 50 55

<210> 118

<211> 62

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (50)

<223> Xaa equals stop translation

<400> 118

Thr Pro Cys Tyr Pro Leu Pro Ala Ala Arg Tyr Thr Val Arg Ser Phe
 1 5 10 15

Gly Ile Arg Arg Asn Glu Lys Ile Ala Val His Cys Thr Val Arg Gly
 20 25 30

Ala Lys Ala Glu Glu Ile Leu Glu Lys Gly Leu Lys Val Ser Leu Ile
 35 40 45

Pro Xaa Trp Ser Asp Ile Asp Gln His Ser Phe Ser Asn Thr
 50 55 60

<210> 119

<211> 281

<212> PRT

<213> Homo sapiens

<400> 119

Ala Glu Asp Gly Pro Glu Val Leu Asp Glu Glu Gly Thr Gln Glu Asp
 1 5 10 15

Leu Glu Tyr Lys Leu Lys Gly Leu Ile Asp Leu Thr Leu Asp Lys Ser
 20 25 30

Ala Lys Thr Arg Gln Ala Ala Leu Glu Gly Val Lys Asn Ala Leu Ser
 35 40 45

Ser Lys Val Leu Tyr Glu Phe Val Leu Glu Arg Arg Met Thr Leu Thr

60

50 55 60
 Asp Ser Ile Glu Arg Cys Leu Lys Lys Gly Lys Ser Asp Glu Gln Arg
 65 70 75 80
 Ala Ala Ala Ala Val Ala Ser Val Leu Cys Ile Gln Leu Gly Pro Gly
 85 90 95
 Phe Glu Ser Glu Glu Ile Leu Lys Thr Leu Gly Pro Ile Leu Lys Lys
 100 105 110
 Ile Ile Cys Asp Gly Ala Ala Ser Ile Gln Ala Arg Gln Thr Cys Ala
 115 120 125
 Thr Cys Phe Gly Val Cys Cys Phe Ile Ala Thr Asp Asp Ile Thr Glu
 130 135 140
 Leu Tyr Ser Thr Leu Glu Cys Phe Glu Asn Ile Phe Thr Lys Ser Tyr
 145 150 155 160
 Leu Lys Glu Lys Asp Thr Asn Val Thr Cys Ser Thr Pro Asn Thr Val
 165 170 175
 Leu His Ile Ser Ser Leu Leu Ala Trp Thr Leu Leu Leu Thr Ile Cys
 180 185 190
 Pro Ile Asn Glu Val Lys Lys Lys Leu Glu Leu His Phe His Lys Leu
 195 200 205
 Pro Ser Leu Leu Ser Cys Asp Asp Val Asn Met Arg Ile Ala Ala Gly
 210 215 220
 Glu Ser Leu Ala Leu Leu Phe Glu Leu Ala Arg Gly Met Glu Ser Asp
 225 230 235 240
 Phe Phe Tyr Glu Asp Met Asp Ser Leu Thr Gln Met Leu Arg Ala Leu
 245 250 255
 Ala Thr Asp Gly Asn Lys His Arg Ala Lys Val Asp Lys Arg Lys Gln
 260 265 270
 Arg Ser Val Phe Arg Asp Val Leu Arg
 275 280

 <210> 120
 <211> 281
 <212> PRT
 <213> Homo sapiens

 <400> 120
 Ser Glu Asp Gly Pro Glu Val Leu Asp Glu Glu Gly Thr Gln Glu Asp
 1 5 10 15
 Leu Glu Tyr Lys Leu Lys Gly Leu Ile Asp Leu Thr Leu Asp Lys Ser

20					25					30					
Ala	Lys	Thr	Arg	Gln	Ala	Ala	Leu	Glu	Gly	Ile	Lys	Asn	Ala	Leu	Ala
	35						40					45			
Ser	Lys	Met	Leu	Tyr	Glu	Phe	Ile	Leu	Glu	Arg	Arg	Met	Thr	Leu	Thr
	50					55					60				
Asp	Ser	Ile	Glu	Arg	Cys	Leu	Lys	Lys	Gly	Lys	Ser	Asp	Glu	Gln	Arg
	65					70					75				80
Ala	Ala	Ala	Ala	Leu	Ala	Ser	Val	Leu	Cys	Ile	Gln	Leu	Gly	Pro	Gly
				85					90					95	
Ile	Glu	Ser	Glu	Glu	Ile	Leu	Lys	Thr	Leu	Gly	Pro	Ile	Leu	Lys	Lys
			100					105					110		
Ile	Ile	Cys	Asp	Gly	Ser	Ala	Ser	Met	Gln	Ala	Arg	Gln	Thr	Cys	Ala
		115					120					125			
Thr	Cys	Phe	Gly	Val	Cys	Cys	Phe	Ile	Ala	Thr	Asp	Asp	Ile	Thr	Glu
	130						135					140			
Leu	Tyr	Ser	Thr	Leu	Glu	Cys	Leu	Glu	Asn	Ile	Phe	Thr	Lys	Ser	Tyr
	145					150					155				160
Leu	Lys	Glu	Lys	Asp	Thr	Thr	Val	Ile	Cys	Ser	Thr	Pro	Asn	Thr	Val
			165						170					175	
Leu	His	Ile	Ser	Ser	Leu	Leu	Ala	Trp	Thr	Leu	Leu	Leu	Thr	Ile	Cys
			180					185					190		
Pro	Ile	Asn	Glu	Val	Lys	Lys	Lys	Leu	Glu	Met	His	Phe	His	Lys	Leu
			195				200					205			
Pro	Ser	Leu	Leu	Ser	Cys	Asp	Asp	Val	Asn	Met	Arg	Ile	Ala	Ala	Gly
	210					215					220				
Glu	Ser	Leu	Ala	Leu	Leu	Phe	Glu	Leu	Ala	Arg	Gly	Ile	Glu	Ser	Asp
	225					230					235				240
Phe	Phe	Tyr	Glu	Asp	Met	Glu	Ser	Leu	Thr	Gln	Met	Leu	Arg	Ala	Leu
			245						250				255		
Ala	Thr	Asp	Gly	Asn	Lys	His	Arg	Ala	Lys	Val	Asp	Lys	Arg	Lys	Gln
			260					265					270		
Arg	Ser	Val	Phe	Arg	Asp	Val	Leu	Arg							
		275					280								

<210> 121

<211> 74

<212> PRT

<213> Homo sapiens

<400> 121

Phe Arg Asp Val Leu Arg Ala Val Glu Glu Arg Asp Phe Pro Thr Glu
 1 5 10 15

Thr Val Lys Phe Gly Pro Glu Arg Met Tyr Ile Asp Ser Trp Val Lys
 20 25 30

Lys His Thr Tyr Asp Thr Phe Lys Glu Val Leu Gly Ser Gly Met Gln
 35 40 45

Tyr His Leu Gln Thr Asn Glu Phe Leu Arg Asn Val Phe Glu Leu Gly
 50 55 60

Pro Pro Val Met Leu Asp Ala Ala Thr Leu
 65 70

<210> 122

<211> 74

<212> PRT

<213> Homo sapiens

<400> 122

Phe Gln Arg Cys Pro Glu Ala Val Glu Glu Arg Asp Phe Pro Thr Glu
 1 5 10 15

Thr Ile Lys Phe Gly Pro Glu Arg Met Tyr Ile Asp Cys Trp Val Lys
 20 25 30

Lys His Thr Tyr Asp Thr Phe Lys Glu Val Leu Gly Ser Gly Met Gln
 35 40 45

Tyr His Leu Gln Ser Asn Glu Phe Leu Arg Asn Val Phe Glu Leu Gly
 50 55 60

Pro Pro Val Met Leu Asp Ala Ala Arg Leu
 65 70

<210> 123

<211> 40

<212> PRT

<213> Homo sapiens

<400> 123

Thr Leu Lys Thr Met Lys Ile Ser Arg Phe Glu Arg His Leu Tyr Asn
 1 5 10 15

Ser Ala Ala Phe Lys Ala Arg Thr Lys Ala Arg Ser Lys Cys Arg Asp
 20 25 30

Lys Arg Ala Asp Val Gly Glu Phe
 35 40

<210> 124
 <211> 39
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (5)
 <223> Xaa equals stop translation

<400> 124
 Thr Leu Lys Arg Xaa Arg Phe Ser Phe Glu Arg His Leu Tyr Asn Ser
 1 5 10 15
 Ala Ala Phe Lys Ala Arg Thr Lys Ala Arg Ser Lys Cys Arg Asp Lys
 20 25 30
 Arg Ala Asp Val Gly Glu Phe
 35

<210> 125
 <211> 92
 <212> PRT
 <213> Homo sapiens

<400> 125
 Lys Trp Ala Lys Asp Met Asn Arg His Phe Ser Lys Glu Asp Ile Tyr
 1 5 10 15
 Ala Ala Lys Lys His Met Lys Lys Cys Ser Pro Ser Leu Ala Ile Arg
 20 25 30
 Glu Met Gln Ile Lys Thr Thr Met Arg Tyr His Leu Thr Pro Val Arg
 35 40 45
 Met Ala Ile Ile Lys Lys Ser Gly Asn Asn Arg Cys Trp Arg Gly Cys
 50 55 60
 Gly Glu Ile Gly Thr Leu Leu His Cys Trp Trp Asn Cys Lys Leu Val
 65 70 75 80
 Gln Pro Leu Trp Lys Ser Val Trp Arg Phe Leu Arg
 85 90

<210> 126
 <211> 90
 <212> PRT
 <213> Homo sapiens

<400> 126
 Lys Val Gly Lys Asn Arg Asn Arg His Phe Ser Lys Glu Glu Ile Gln
 1 5 10 15

Ile Ala Lys Lys Asp Lys Lys Arg Cys Ser Thr Leu Ser Val Ile Met
 20 25 30
 Lys Met Gln Met Lys Thr Thr Arg His Cys Phe Thr Pro Asn Met Gly
 35 40 45
 Trp Lys Gln Lys Val Arg Glu Lys Gln Val Met Ala Lys Met Trp Arg
 50 55 60
 Asn Gln Asn Ser Gln Gly Cys Arg Lys Gly Arg Glu Met Val Gln Thr
 65 70 75 80
 Leu Gly Lys Ala Gly Trp Arg Leu Leu Lys
 85 90

<210> 127
 <211> 203
 <212> PRT
 <213> Homo sapiens

<400> 127
 Ser Gln Val Val Asn Ala Lys Glu Lys Phe Leu Lys Glu Ile Lys Ser
 1 5 10 15
 Ala Thr Pro Val Asn Thr Arg Met Ile Arg Lys Gln Asn Ser Leu Ile
 20 25 30
 Ala Asp Met Glu Lys Val Leu Val Val Trp Ile Glu Asp Gln Thr Ser
 35 40 45
 His Asn Ile Pro Leu Ser Gln Ser Leu Ile Gln Ser Lys Ala Leu Thr
 50 55 60
 Leu Phe Asn Ser Met Lys Ala Glu Arg Gly Glu Glu Ala Ala Glu Glu
 65 70 75 80
 Lys Leu Glu Ala Ser Arg Gly Trp Phe Met Arg Phe Lys Glu Arg Ser
 85 90 95
 His Leu His Asn Ile Lys Val Gln Gly Glu Ala Ala Ser Ala Asp Val
 100 105 110
 Glu Ala Ala Ala Ser Tyr Pro Glu Asp Leu Ala Lys Ile Ile Asp Glu
 115 120 125
 Gly Gly Tyr Thr Lys Gln Gln Ile Phe Asn Val Asp Glu Thr Ala Phe
 130 135 140
 Tyr Trp Lys Lys Met Pro Ser Arg Thr Phe Ile Ala Arg Glu Glu Lys
 145 150 155 160
 Ser Met Pro Gly Phe Lys Ala Ser Lys Asp Arg Leu Thr Leu Leu Leu
 165 170 175

Gly Ala Asn Ala Ala Gly Asp Phe Lys Leu Lys Pro Met Leu Ile Tyr
 180 185 190
His Ser Glu Asn Pro Arg Ala Leu Lys Asn Tyr
 195 200

<210> 128
<211> 198
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (11)
<223> Xaa equals stop translation

<220>
<221> SITE
<222> (24)
<223> Xaa equals stop translation

<220>
<221> SITE
<222> (25)
<223> Xaa equals stop translation

<220>
<221> SITE
<222> (82)
<223> Xaa equals stop translation

<220>
<221> SITE
<222> (126)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (138)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (141)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (143)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE

<222> (151)

<223> Xaa equals stop translation

<220>

<221> SITE

<222> (168)

<223> Xaa equals stop translation

<400> 128

Ser Gln Val Val Asn Ala Met Lys Ser Ser Xaa Arg Lys Leu Lys Val
 1 5 10 15

Leu Leu His Arg Thr His Glu Xaa Xaa Gly Lys His Leu Ile Ser Asp
 20 25 30

Ile Glu Arg Val Leu Met Val Trp Ile Glu Asp Gln Ile Ser His Asn
 35 40 45

Val Ser Leu Ser Arg Ser Gln Ser Lys Ala Leu Thr Leu Phe Asn Ser
 50 55 60

Leu Lys Ala Glu Thr Gly Glu Gly Ala Ala Ala Lys Lys Leu Glu Ala
 65 70 75 80

Ser Xaa Gly Trp Phe Met Arg Phe Lys Lys Arg Ser Tyr Leu Arg Asn
 85 90 95

Ile Lys Val Leu Asp Glu Ala Ala Ser Ala Asp Val Glu Ala Ala Val
 100 105 110

Ser Tyr Pro Glu Asp Leu Ala Gly Ile Ile Asp Lys Ala Xaa Tyr Thr
 115 120 125

Lys Gln His Ile Phe Thr Val Asn Gln Xaa Ala Ile Xaa Glu Xaa Arg
 130 135 140

Cys Cys Leu Gly Leu Ser Xaa Leu Glu Ser Thr Arg His Arg Leu Phe
 145 150 155 160

Leu Lys Leu Glu Arg Thr Ser Xaa Pro Leu Val Gly Ala Asn Ala Ala
 165 170 175

Ser Asp Thr Lys Leu Lys Leu Val Leu Ile Phe Pro Ser Gln Asn Pro
 180 185 190

Thr Ala Leu Gln Asn Tyr
 195

<210> 129

<211> 211

<212> PRT

<213> Homo sapiens

<400> 129

67

Pro Ile Ala Ser Trp Gly Val Pro Tyr Asp Gln Leu Thr Glu Glu Glu
 1 5 10 15
 Lys Thr Arg Ala Trp Phe Thr Asp Gly Ser Ala Arg Tyr Ala Gly Thr
 20 25 30
 Thr Gln Lys Trp Thr Ala Ala Ala Leu Gln Pro Leu Ser Gly Thr Thr
 35 40 45
 Leu Lys Asp Thr Gly Glu Arg Lys Ser Ser Gln Trp Ala Glu Leu Arg
 50 55 60
 Ala Val His Met Val Leu Gln Phe Val Cys Lys Lys Lys Trp Pro Asp
 65 70 75 80
 Val Arg Leu Phe Thr Asp Ser Trp Ala Val Ala Asn Gly Leu Ala Gly
 85 90 95
 Trp Ser Gly Thr Trp Lys Asp His Asn Trp Lys Ile Gly Glu Lys Asp
 100 105 110
 Ile Trp Gly Arg Ser Met Trp Ile Asp Ile Ser Lys Trp Ala Lys Asp
 115 120 125
 Val Lys Ile Phe Val Ser His Val Asn Ala His Gln Lys Val Thr Ser
 130 135 140
 Ala Glu Glu Glu Phe Asn Asn Gln Val Asp Lys Met Thr Arg Ser Val
 145 150 155 160
 Asp Ser Gln Thr Leu Ser Pro Ala Ile Pro Val Ile Ala Gln Trp Ala
 165 170 175
 His Glu Gln Ser Gly His Gly Gly Arg Asp Gly Gly Tyr Pro Trp Ala
 180 185 190
 Gln Gln His Gly Leu Pro Leu Thr Lys Ala Asp Leu Ala Thr Ala Ala
 195 200 205
 Ala Asp Cys
 210

<210> 130

<211> 195

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (15)

<223> Xaa equals stop translation

<220>

<221> SITE

```

<222> (24)
<223> Xaa equals stop translation

<220>
<221> SITE
<222> (28)
<223> Xaa equals stop translation

<220>
<221> SITE
<222> (34)
<223> Xaa equals stop translation

<220>
<221> SITE
<222> (78)
<223> Xaa equals stop translation

<220>
<221> SITE
<222> (96)
<223> Xaa equals stop translation

<220>
<221> SITE
<222> (104)
<223> Xaa equals stop translation

<220>
<221> SITE
<222> (113)
<223> Xaa equals stop translation

<220>
<221> SITE
<222> (168)
<223> Xaa equals stop translation

<220>
<221> SITE
<222> (177)
<223> Xaa equals stop translation

<400> 130
Pro Met Ala Pro Trp Gly Phe Leu His Arg Gln Leu Thr Glu Xaa Lys
 1             5             10             15

Arg Ser Arg Lys Arg Ala Leu Xaa Gly Leu Gly Xaa His Asp Met Leu
      20             25             30

Ala Xaa Ala Lys Lys Met Asp Trp Cys Cys Thr Lys Asp Thr Pro Lys
 35             40             45

Ile Thr Leu Gly Ala Asn Ser Ser Pro Val Val Ile Cys Phe Val Ser
 50             55             60

```

Gly Glu Lys Arg Pro Lys Val Arg Ile Tyr Thr Asp Phe Xaa Val Val
 65 70 75 80
 Glu Asn Gly Leu Val Ala Trp Ser Arg Ala Ser Arg Glu Gln His Xaa
 85 90 95
 Arg Ile Glu Thr Lys His Val Xaa Gly Arg Gly Met Trp Leu Glu Leu
 100 105 110
 Xaa Glu Trp Ala His Asn Val Leu Ile Phe Val Ser His Thr Arg Ala
 115 120 125
 His Gln Arg Ala Arg Val Ala Glu Glu Ala Leu Asn Lys His Met Asp
 130 135 140
 Arg Met Thr Gly Leu Met Asn Ala Cys Gln Pro Leu Ser Ser Ala Ile
 145 150 155 160
 Ser Ile Leu Thr Lys Gly Pro Xaa Met Asn Gly Leu Val Val Met Ala
 165 170 175
 Xaa Asn Asp Gly Leu Ser Leu Ser Lys Ala Asp Leu Ala Thr Thr Thr
 180 185 190
 Thr Glu Cys
 195

<210> 131
 <211> 96
 <212> PRT
 <213> Homo sapiens

<400> 131
 Lys Lys Thr Asn Asn Pro Ile Lys Lys Trp Ala Lys Asp Met Asn Arg
 1 5 10 15
 His Phe Ser Lys Glu Asp Ile Tyr Ala Ala Lys Lys His Met Lys Lys
 20 25 30
 Cys Ser Pro Ser Leu Ala Ile Arg Glu Met Gln Ile Lys Thr Thr Met
 35 40 45
 Arg Tyr His Leu Thr Pro Val Arg Met Ala Ile Ile Lys Lys Ser Gly
 50 55 60
 Asn Asn Arg Cys Trp Arg Gly Cys Gly Glu Ile Gly Thr Leu Leu His
 65 70 75 80
 Cys Trp Trp Asn Cys Lys Leu Val Gln Pro Leu Trp Lys Ser Val Trp
 85 90 95

<210> 132
 <211> 97
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (21)
 <223> Xaa equals stop translation

<400> 132
 Glu Lys Ser Asn His Pro Pro Ala Asp Pro Thr Ser Asn Thr Gly Ser
 1 5 10 15
 Tyr Asn Ser Thr Xaa Asp Leu Val Gly Thr Gln Ile Gln Thr Ile Ser
 20 25 30
 Val Leu Asn Ile Ile Ile Ile Arg Glu Met Gln Ile Lys Thr Thr Ile
 35 40 45
 Arg Cys His Leu Thr Leu Val Gln Met Ala Phe Ile Gln Lys Thr Gly
 50 55 60
 Asn Asn Lys Cys Trp Gln Glu Cys Gly Glu Lys Gly Thr Leu Ile His
 65 70 75 80
 Ile Ser Trp Trp Lys Ser Lys Ser Val Gln Pro Leu Trp Lys Thr Val
 85 90 95

Trp

<210> 133
 <211> 59
 <212> PRT
 <213> Homo sapiens

<400> 133
 Trp Asp Tyr Arg His Val Pro Pro Arg Gln Val His Phe Val Phe Ser
 1 5 10 15
 Val Glu Thr Gly Phe His Arg Ala Gly Gln Ala Gly Leu Glu Leu Leu
 20 25 30
 Thr Ser Ser Val Pro Pro Thr Ser Ala Phe Pro Lys Cys Trp Asp Tyr
 35 40 45
 Arg Arg Asp Asp Gln Ala Trp Pro Thr Leu Ser
 50 55

<210> 134

<211> 59
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (26)
 <223> Xaa equals stop translation

<220>
 <221> SITE
 <222> (34)
 <223> Xaa equals stop translation

<400> 134
 Trp Asp Tyr Arg His Met Pro Pro Leu Arg Leu Ile Phe Val Phe Ser
 1 5 10 15
 Val Glu Thr Gly Ser His His Ala Ala Xaa Ala Asp Leu Glu Leu Leu
 20 25 30
 Ser Xaa Thr Asp Pro Pro Ala Ser Ala Ser Gln Asn Thr Arg Thr Thr
 35 40 45
 Gly Val Ser His Arg Ala Trp Pro Ser Leu Ala
 50 55

<210> 135
 <211> 26
 <212> PRT
 <213> Homo sapiens

<400> 135
 Asp Arg Leu Ser Leu Leu Ser Pro Arg Leu Glu Cys Asn Gly Met Ile
 1 5 10 15
 Leu Ala His Cys Lys Leu Arg Leu Pro Gly
 20 25

<210> 136
 <211> 26
 <212> PRT
 <213> Homo sapiens

<400> 136
 Asp Arg Ile Ser Leu Leu Ser Pro Arg Leu Glu Cys Asn Gly Val Ile
 1 5 10 15
 Leu Ala Asn Tyr Asn Leu Arg Leu Pro Gly
 20 25

<210> 137

<211> 34
<212> PRT
<213> Homo sapiens

<400> 137
Ala His Cys Asn Leu Cys Leu Pro Gly Ser Ser Asn Ser Pro Ala Ser
1 5 10 15

Ala Ser Arg Val Ala Gly Thr Ala Gly Thr Cys Arg Arg Ala Gln Leu
20 25 30

Ile Phe

<210> 138
<211> 34
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (3)
<223> Xaa equals stop translation

<400> 138
Ala His Xaa Asn Leu Tyr Leu Pro Gly Ser Ser Asn Pro Leu Thr Ser
1 5 10 15

Ala Ser Gln Val Ala Gly Thr Thr Gly Thr Cys His Gln Thr Arg Leu
20 25 30

Ile Phe

<210> 139
<211> 18
<212> PRT
<213> Homo sapiens

<400> 139
Glu Thr Gln Ser His Ser Val Thr Arg Leu Glu Cys Ser Gly Thr Ile
1 5 10 15

Ser Ala

<210> 140
<211> 18
<212> PRT
<213> Homo sapiens

<400> 140

Glu Thr Gly Ser His Ser Val Ala Gln Val Glu Cys Ser Gly Ala Ser
 1 5 10 15

Ser Ser

<210> 141
 <211> 100
 <212> PRT
 <213> Homo sapiens

<400> 141
 Gly Asp Arg Leu Thr Gly Ile Pro Ser His Ile Leu Asn Ser Ser Pro
 1 5 10 15

Ser Asp Arg Gln Ile Asn Gln Leu Ala Gln Arg Leu Gly Pro Glu Trp
 20 25 30

Glu Pro Met Val Leu Ser Leu Gly Leu Ser Gln Thr Asp Ile Tyr Arg
 35 40 45

Cys Lys Ala Asn His Pro His Asn Val Gln Ser Gln Val Val Glu Ala
 50 55 60

Phe Ile Arg Trp Arg Gln Arg Phe Gly Lys Gln Ala Thr Phe Gln Ser
 65 70 75 80

Leu His Asn Gly Leu Arg Ala Val Glu Val Asp Pro Ser Leu Leu Leu
 85 90 95

His Met Leu Glu
 100

<210> 142
 <211> 100
 <212> PRT
 <213> Homo sapiens

<400> 142
 Gly Asp Arg Leu Thr Gly Ile Pro Ser His Ile Leu Asn Ser Ser Pro
 1 5 10 15

Ser Asp Arg Gln Ile Asn Gln Leu Ala Gln Arg Leu Gly Pro Glu Trp
 20 25 30

Glu Pro Met Val Leu Ser Leu Gly Leu Ser Gln Thr Asp Ile Tyr Arg
 35 40 45

Cys Lys Ala Asn His Pro His Asn Val Gln Ser Gln Val Val Glu Ala
 50 55 60

Phe Ile Arg Trp Arg Gln Arg Phe Gly Lys Gln Ala Thr Phe Gln Ser
 65 70 75 80

Leu His Asn Gly Leu Arg Ala Val Glu Val Asp Pro Ser Leu Leu Leu
85 90 95

His Met Leu Glu
100

<210> 143
<211> 448
<212> PRT
<213> Homo sapiens

<400> 143
Ser Met Asp Phe Asp Asp Thr Trp His Pro Ala Thr His Pro Ser Gly
1 5 10 15
Ala Val Leu Pro Val Leu Thr Ala Leu Ser Glu Ala Leu Pro Gln Ile
20 25 30
Pro Lys Phe Ser Gly Leu Asp Leu Leu Leu Ala Phe Asn Val Gly Ile
35 40 45
Glu Val Gln Gly Arg Leu Met His Phe Ser Lys Glu Ala Lys Asp Ile
50 55 60
Pro Lys Arg Phe His Pro Pro Ser Val Val Gly Thr Leu Gly Ser Ala
65 70 75 80
Ala Ala Ala Ser Lys Phe Leu Gly Leu Ser Leu Thr Lys Cys Arg Glu
85 90 95
Ala Leu Ala Ile Ala Val Ser His Ala Gly Ala Pro Ile Ala Asn Ala
100 105 110
Ala Thr Gln Thr Lys Pro Leu His Ile Gly Asn Ala Ala Lys His Gly
115 120 125
Met Glu Ala Thr Phe Leu Ala Met Leu Gly Leu Gln Gly Asn Lys Gln
130 135 140
Ile Leu Asp Leu Gly Ser Gly Phe Gly Ala Phe Tyr Ala Asn Tyr Ser
145 150 155 160
Pro Glu Asp Leu Pro Ser Leu Asp Ser His Ile Trp Leu Leu Asp Gln
165 170 175
Gln Asp Val Ala Phe Lys Ser Phe Pro Ala His Leu Ala Thr His Trp
180 185 190
Val Ala Asp Ala Ala Ala Ala Val Arg Lys His Leu Val Thr Pro Glu
195 200 205
Arg Ala Leu Phe Pro Ala Asp His Ile Glu Arg Ile Val Leu Arg Ile
210 215 220

Pro Asp Val Gln Tyr Val Asn Arg Pro Phe Pro Asp Ser Glu His Glu
 225 230 235 240
 Ala Arg His Ser Phe Gln Tyr Val Ala Cys Ala Ser Leu Leu Asp Gly
 245 250 255
 Ser Ile Thr Val Pro Ser Phe His Ser Gln Gln Val Asn Arg Pro Gln
 260 265 270
 Val Arg Glu Leu Leu Lys Lys Val Lys Leu Glu His Pro Pro Asp Asn
 275 280 285
 Pro Pro Ser Phe Asp Thr Leu Tyr Cys Glu Ile Ser Ile Thr Leu Lys
 290 295 300
 Asp Gly Thr Thr Phe Thr Glu Arg Ser Asp Thr Phe Tyr Gly His Trp
 305 310 315 320
 Arg Lys Pro Leu Ser Gln Glu Asp Leu Arg Asn Lys Phe Arg Ala Asn
 325 330 335
 Ala Ser Lys Met Leu Cys Arg Asp Thr Val Glu Ser Leu Ile Thr Val
 340 345 350
 Val Glu Lys Leu Glu Asp Leu Glu Asp Cys Ser Val Leu Thr Arg Leu
 355 360 365
 Leu Lys Asp Pro Leu Ser Lys Met Lys Leu Gln Asn Tyr Pro Ala Cys
 370 375 380
 Pro His Ser Ile Thr Gln Arg Cys Pro Gly Leu Pro Ile Ser Lys Arg
 385 390 395 400
 Ala Leu Ala Leu Glu Glu Ile His Cys Phe Gly Phe Leu Leu His Leu
 405 410 415
 Pro Ser Asn Glu Leu Ser Lys Val Glu Ser Pro Glu Thr Glu Thr Ser
 420 425 430
 Val Tyr Gly Lys Arg Leu Val Tyr Asn Leu Ile Lys Thr Ser Pro Thr
 435 440 445

<210> 144

<211> 451

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (358)

<223> Xaa equals stop translation

<220>

<221> SITE

<222> (364)

<223> Xaa equals stop translation

<220>

<221> SITE

<222> (383)

<223> Xaa equals stop translation

<220>

<221> SITE

<222> (384)

<223> Xaa equals stop translation

<220>

<221> SITE

<222> (397)

<223> Xaa equals stop translation

<400> 144

Ser Met Asp Phe Asp Asp Thr Trp His Pro Ala Thr His Pro Ser Gly
1 5 10 15

Ala Val Leu Pro Val Leu Thr Ala Leu Ala Glu Ala Leu Pro Arg Ser
20 25 30

Pro Lys Phe Ser Gly Leu Asp Leu Leu Ala Phe Asn Val Gly Ile
35 40 45

Glu Val Gln Gly Arg Leu Leu His Phe Ala Lys Glu Ala Asn Asp Met
50 55 60

Pro Lys Arg Phe His Pro Pro Ser Val Val Gly Thr Leu Gly Ser Ala
65 70 75 80

Ala Ala Ala Ser Lys Phe Leu Gly Leu Ser Ser Thr Lys Cys Arg Glu
85 90 95

Ala Leu Ala Ile Ala Val Ser His Ala Gly Ala Pro Met Ala Asn Ala
100 105 110

Ala Thr Gln Thr Lys Pro Leu His Ile Gly Asn Ala Ala Lys His Gly
115 120 125

Ile Glu Ala Ala Phe Leu Ala Met Leu Gly Leu Gln Gly Asn Lys Gln
130 135 140

Val Leu Asp Leu Glu Ala Gly Phe Gly Ala Phe Tyr Ala Asn Tyr Ser
145 150 155 160

Pro Lys Val Leu Pro Ser Ile Ala Ser Tyr Ser Trp Leu Leu Asp Gln
165 170 175

Gln Asp Val Ala Phe Lys Arg Phe Pro Ala His Leu Ser Thr His Trp
 180 185 190
 Val Ala Asp Ala Ala Ala Ser Val Arg Lys His Leu Val Pro Glu Arg
 195 200 205
 Ala Leu Leu Pro Thr Asp Tyr Ile Lys Arg Ile Val Leu Arg Ile Pro
 210 215 220
 Asn Val Gln Tyr Val Asn Arg Pro Phe Pro Val Ser Glu His Glu Ala
 225 230 235 240
 Arg His Ser Phe Gln Tyr Val Ala Cys Ala Met Leu Leu Asp Gly Gly
 245 250 255
 Ile Thr Val Pro Ser Phe His Glu Cys Gln Ile Asn Arg Pro Gln Val
 260 265 270
 Arg Glu Leu Leu Ser Lys Val Glu Leu Glu Tyr Pro Pro Asp Asn Leu
 275 280 285
 Pro Ser Phe Asn Ile Leu Tyr Cys Glu Ile Ser Val Thr Leu Lys Asp
 290 295 300
 Gly Ala Thr Phe Thr Asp Arg Ser Asp Thr Phe Tyr Gly His Trp Arg
 305 310 315 320
 Lys Pro Leu Ser Gln Glu Asp Leu Glu Glu Lys Phe Arg Ala Asn Ala
 325 330 335
 Ser Lys Met Leu Ser Trp Asp Thr Val Glu Ser Leu Ile Lys Ile Val
 340 345 350
 Lys Asn Leu Glu Asp Xaa Lys Thr Val Leu Cys Xaa Leu His Phe Ser
 355 360 365
 Lys Asp Leu Ser Thr Arg Gly Ser Phe Lys Leu Ser Ser Met Xaa Xaa
 370 375 380
 Phe Tyr His Lys Ser Leu Leu Arg Leu Thr Asn Ile Xaa Met Thr Leu
 385 390 395 400
 His Leu Gly Arg Phe Asn Asp Leu Val Cys Lys Ala Arg Val Cys Cys
 405 410 415
 Leu Val Phe Pro Gly Lys Met Asn Lys Asp Gly Glu Ser Pro Glu Thr
 420 425 430
 Glu Leu His Ile Ser Gly Arg Ser Leu Leu Leu Lys Ile Leu Gln Asp
 435 440 445
 Ser Ser Thr
 450

PATENT COOPERATION TREATY

PCT

DECLARATION OF NON-ESTABLISHMENT OF INTERNATIONAL SEARCH REPORT

(PCT Article 17(2)(a), Rules 13ter and 39)

Applicant's or agent's file reference PS526PCT	IMPORTANT DECLARATION	Date of mailing (day/month/year) 13 AUG 2000
International application No. PCT/US00/07525	International filing date (day/month/year) 22 MARCH 2000	(Earliest) Priority Date (day/month/year) 26 MARCH 1999
International Patent Classification (IPC) or both national classification and IPC Please See Continuation Sheet.		
Applicant HUMAN GENOME SCIENCES, INC.		

This International Searching Authority hereby declares, according to Article 17(2)(a), that no international search report will be established on the international application for the reasons indicated below.

1. ☐ The subject matter of the international application relates to:
 - a. ☐ scientific theories.
 - b. ☐ mathematical theories.
 - c. ☐ plant varieties.
 - d. ☐ animal varieties.
 - e. ☐ essentially biological processes for the production of plants and animals, other than microbiological processes and the products of such processes.
 - f. ☐ schemes, rules or methods of doing business.
 - g. ☐ schemes, rules or methods of performing purely mental acts.
 - h. ☐ schemes, rules or methods of playing games.
 - i. ☐ methods for treatment of the human body by surgery or therapy.
 - j. ☐ methods for treatment of the animal body by surgery or therapy.
 - k. ☐ diagnostic methods practiced on the human or animal body.
 - l. ☐ mere presentations of information.
 - m. ☐ computer programs for which this International Searching Authority is not equipped to search prior art.
2. ☐ The failure of the following parts of the international application to comply with prescribed requirements prevents a meaningful search from being carried out:

☐ the description
 ☐ the claims
 ☐ the drawings
3. ☒ The failure of the nucleotide and/or amino acid sequence listing to comply with the standard provided for in Annex C of the Administrative Instructions prevents a meaningful search from being carried out.

☐ the written form has not been furnished or does not comply with the standard.
☒ the computer readable form has not been furnished or does not comply with the standard.
4. Further comments:
Please See Continuation Sheet.

Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231	Authorized officer <i>my mas</i> MARY K ZEMAN
Facsimile No. (703) 305-3230	Telephone No. (703) 308-0196

**DECLARATION OF NON-ESTABLISHMENT OF
INTERNATIONAL SEARCH REPORT**

International application No.
PCT/US00/07525

The International Patent Classification (IPC) or National Classification and IPC are as listed below:

A61K 39/00; G01N 33/53; C12P 15/09

435/7.1, 69.3, 320.1; 530/350

4. Further Comments (Continued):

The computer readable format did not comply with the rules, and did not match the specification such that a search could not be performed. All claims required a search of at least one sequence.